arXiv:1205.7057v1 [math.AT] 31 May 2012

RATIONAL HOMOTOPY AND INTERSECTION COHOMOLOGY

DAVID CHATAUR, MARTINTXO SARALEGI-ARANGUREN,
AND DANIEL TANRÉ

Abstract. In this text, we extend Sullivan's presentation of rational homotopy type to Goresky and MacPherson's intersection cohomology, providing also a homotopical framework where intersection cohomology is a representable functor.

We choose the context of face sets, also called simplicial sets without degeneracies, introduced by Rourke and Sanderson. We define filtered face sets, perverse local systems over them and intersection cohomology with coefficients in a perverse local system. In particular, we get a perverse local system of cochains quasi-isomorphic to the intersection cochains of Goresky and MacPherson, over $\mathbb{Z}$. We show also that these two complexes are quasi-isomorphic to a filtered version of Sullivan's differential forms over the field $\mathbb{Q}$.

We construct a pair of contravariant adjoint functors between the category of filtered face sets and a category of perverse commutative differential graded algebras (cdga's) due to Hovey. This correspondence gives bijections between homotopy classes, with the usual restriction to fibrant and cofibrant objects. We establish also the existence and unicity of a positively graded, minimal model of some perverse cdga's, including the perverse forms over a filtered face set and their intersection cohomology. This brings a definition of formality in the intersection setting. Finally, we prove the topological invariance of the minimal model of a locally conelike space and this theory creates new topological invariants.

Intersection cohomology of locally conelike singular manifolds has been introduced by Goresky and MacPherson in [16] and [17]. The main feature of their theory concerns the transversality of a simplex relatively to a stratum, the notion of general position being replaced by a less restrictive one depending on a parameter called perversity. We supply the necessary definitions all along the text but, for a more complete background, the reader can consult the original papers or one of the books or proceedings that appeared in the last years as, for instance, [3], [24], [18], [2], [1], or the historical development presented in [25].

Our work is motivated by a presentation of the intersection cohomology, similar to Sullivan's presentation of the rational homotopy type, including the notion of minimal model, with its topological invariance (for locally conelike spaces) and a proof of the representability of intersection cohomology. This rational intersection theory answers a question raised by Goresky ([19,

---

Date: March 6, 2013.

2000 *Mathematics Subject Classification.* 55N33, 55P62, 57N80.

The third author is partially supported by the MICINN grant MTM2010-18089 and the ANR-11-BS01-002-01 "HOGT".

Introduction]) in view of a treatment of formality. In fact, we offer more than a rational point of view, by providing a homotopical framework for intersection cohomology and many sections of this paper are concerned with homotopy type without localization over the field of rational numbers. This brings a new approach for the study of Steenrod operations in intersection cohomology (see [19]) as quoted by MacPherson in [19, Question 4.3]. We will come back to this aspect in a forthcoming paper.

Our work is developed along the lines of Sullivan's theory. (The reader can consult [35], [27], [8], [9], [21] or [36] for a presentation of rational homotopy.) This theory begins with a functor from simplicial sets to the category of commutative graded differential algebras (henceforth CDGA's), denoted by A_{PL}, such that the cohomology of $A_{PL}(\underline{K})$ is isomorphic to the cohomology algebra of the simplicial set $\underline{K}$, with rational coefficients. One important step is thus the determination of a simplicial setting providing a homotopical context to intersection cohomology.

This work is organized as follows. We start from the topological situation, which covers *the first three sections.* The original work of Goresky and MacPherson is designed for locally conelike manifolds filtered by PL-subspaces, $(M_i)_{0\leq i\leq n}$, of respective dimension i. As in King's paper ([23]), we are concerned with general topological spaces, X, together with a filtration by closed subspaces, $X_0 \subseteq X_1 \subseteq \cdots \subseteq X_n = X$. We consider singular simplices, $\sigma\colon \Delta = \Delta^{j_0} * \cdots \Delta^{j_n} \to X$, whose domain is decomposed as a join product such that $\sigma^{-1}(X_i) = \Delta^{j_0} * \cdots * \Delta^{j_i}$. To any such simplex, called *filtered simplex,* and to any number $i \in \{0, \ldots, n\}$ we associate the perverse degree $\|\sigma\|_i = \dim(\Delta^{j_0} * \cdots * \Delta^{j_{n-i}})$. Now comes the fundamental notion of perversity. As it will be justified later (see Example 8.9 for instance), the lack of relation between the filtration degree and a geometrical dimension leads us to consider a notion of perversity with less constraints than in the Goresky and MacPherson original work. We call *loose perversity* (see [23]) any map $\overline{p}\colon \mathbb{N} \to \mathbb{Z}$ such that $\overline{p}(0) = 0$. A filtered simplex as above is said $\overline{p}$-admissible if the inequality $\|\sigma\|_i \leq \dim \Delta - \overline{p}(i) + i$ is satisfied for all $i \in \{0, \ldots, n\}$ and a chain c is $\overline{p}$-admissible if there exist $\overline{p}$-admissible simplices, σ_j, so that $c = \sum_j \lambda_j \sigma_j$, $\lambda_j \in \mathbb{Z}$. We denote by $C_*^{\overline{p}}(X)$ the complex of $\overline{p}$-admissible chains, c, whose boundary ∂c is $\overline{p}$-admissible also. There come our two first results.

- Theorem A: For any loose perversity $\overline{p}$, a stratified map (see Definition 1.5) $f\colon X \to Y$, induces a chain map $f_*\colon C_*^{\overline{p}}(X) \to C_*^{\overline{p}}(Y)$.
- Proposition 3.4: In the case of a locally conelike space (see Definition 3.1) our complex gives the same intersection homology than the Goresky and MacPherson original one.

The first property is a key in the proof of the topological invariance stated at the end of this paper. In the proof of the second statement, we use the fact that a locally conelike space is a stratified space (Theorem B) and we rely on King's paper where a chain complex, similar to ours, is connected to the Goresky and MacPherson complex.

Section 4 contains a simplicial set presentation of this theory. We choose the context of face sets, also called simplicial sets without degeneracies, introduced by Rourke and Sanderson ([30]). We denote by $\mathbf{\Delta}_{\mathcal{F}}^{[n]}$ the category whose objects are the decomposed euclidian simplices, $\Delta^{j_0} * \cdots * \Delta^{j_n}$, and maps are joins of face operators, i.e., they are compositions of maps $f = *_{i=0}^{n}\partial_i$, with ∂_i a face operator. The filtered face sets are defined as functors, $\underline{K}$, from $\mathbf{\Delta}_{\mathcal{F}}^{[n]}$ to the category of sets. For instance, the filtered simplices of a filtered topological space constitute a filtered face set, see Example 4.5.

Sections 5, 6 and 7 are concerned with various cohomologies defined on a filtered face set $\underline{K}$. Adapting the topological situation described above, we associate, to $\underline{K}$ and to a loose perversity $\overline{p}$, a cochain complex $C^*_{\mathrm{GM},\overline{p}}(\underline{K},\mathbb{Z})$ corresponding to the original Goresky and MacPherson one. We look also for other cochain complexes built on $\underline{K}$ and quasi-isomorphic to the previous one. To proceed, we introduce a simplicial version of a blow-up, already defined in the thesis of the second author ([31]) and used for singular manifolds in [4] and for some particular simplicial complexes in [6]. To any filtered simplex, $\Delta = \Delta^{j_0} * \cdots * \Delta^{j_n}$, we associate $\widetilde{\Delta} = c\Delta^{j_0} \times \cdots \times c\Delta^{j_{n-1}} \times \Delta^{j_n}$, called the blow-up of Δ. If $A_{PL}(\Delta^i)$ is the CDGA of rational polynomial forms on Δ^i, we define $\widetilde{A}_{PL}(\Delta) = A_{PL}(c\Delta^{j_0}) \otimes \cdots \otimes A_{PL}(c\Delta^{j_{n-1}}) \otimes A_{PL}(\Delta^{j_n})$. We do not want to go into technical points in this introduction but the reader should be aware that this definition makes sense only if $\Delta^{j_n} \neq \emptyset$. Thus, the compatibility condition for the definition of the CDGA, $\widetilde{A}_{PL}(\underline{K})$, of global forms on $\underline{K}$ concerns only the face maps which satisfy this condition. We define a perverse degree for any global form $\omega \in \widetilde{A}_{PL}(\underline{K})$ which is a sort of degree along the fibre, see Example 5.9. Now, for any loose perversity $\overline{q}$, a global form $\omega \in \widetilde{A}_{PL}(\underline{K})$ is said $\overline{q}$-admissible if $\|\omega\|_i \leq \overline{q}(i)$ for all i. The complex $\widetilde{A}_{PL,\overline{q}}(\underline{K})$ is generated by the $\overline{q}$-admissible forms ω whose boundary $d\omega$ is $\overline{q}$-admissible also. We prove that the integration map

$$\int : \widetilde{A}_{PL,\overline{q}}(\underline{K}) \to C^*_{\mathrm{GM},\overline{p}}(\underline{K};\mathbb{Q})$$

induces a quasi-isomorphism if $\overline{p}$ and $\overline{q}$ are perversities verifying $\overline{q} \geq 0$ and $\overline{p}(j) + \overline{q}(j) = j - 2$.

We may consider again this construction replacing $A_{PL}(\Delta^i)$ by the cochain algebra $C^*(\Delta^i;\mathbb{Z})$. This gives a cochain complex $\widetilde{C}^*(\underline{K};\mathbb{Z})$ connected to the Goresky and MacPherson cochain complex by a quasi-isomorphism

$$\widetilde{C}^*_{\overline{q}}(\underline{K};\mathbb{Z}) \simeq C^*_{\mathrm{GM},\overline{p}}(\underline{K};\mathbb{Z}),$$

if $\overline{p}$ and $\overline{q}$ are perversities verifying $\overline{q} \geq 0$ and $\overline{p}(j) + \overline{q}(j) = j - 2$. This opens a treatment of intersection cohomology with coefficients in $\mathbb{Z}$ from local cochains. The existence of the previous quasi-isomorphisms are contained in Theorem C and Theorem D.

In *Section 8,* we study particular cases and examples. For instance, we prove that the $\overline{\infty}$-intersection cohomology is the cohomology of the *regular* part, $\underline{K}^{[0]}$, which consists of simplices $\Delta^{j_0} * \cdots * \Delta^{j_n}$ such that $\Delta^{j_i} = \emptyset$ if $i \leq n-1$. Also, the $\overline{0}$-intersection cohomology is the cohomology of $\underline{K}$ if $\underline{K}$ is a normal filtered face set (see Definition 8.2). As in [16], we prove that any filtered face set, $\underline{K}$, admits a unique normalization, $N(\underline{K})$, which has the same intersection cohomology than $\underline{K}$, see Proposition 8.7. Examples of cone and suspension of a filtered face set are detailed. We emphasize the fact that we find the intersection cohomologies with a filtration not coming from a geometrical dimension. This can be done with the original Goresky and MacPherson perversities if we work with the blow-up of differential forms but we must consider negative perversities if we use the original Goresky and MacPherson cochains. Example 8.9 is a concrete illustration of this situation.

In Section 9, we supply a cylinder object, define the product of a filtered face set with a face set and a notion of homotopy between maps of filtered face sets.

In *Section 10,* we come back to our rational homotopy project and recall the algebraic category, $\mathrm{CDGA}_{\mathcal{F}}$, of perverse CDGA's defined by Hovey in [22], see Definition 10.1 and Definition 10.3. Our previous construction of the blow-up of Sullivan's forms gives a functor $\mathcal{A}\colon \mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets \to \mathrm{CDGA}_{\mathcal{F}}$, $\underline{K} \mapsto \mathcal{A}(\underline{K})_\bullet = \widetilde{A}_{PL,\bullet}(\underline{K})$.

In Section 11, we build a contravariant adjoint functor $\langle - \rangle\colon \mathrm{CDGA}_{\mathcal{F}} \to \mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets$ to $\mathcal{A}$, i.e., we have a bijection

$$\hom_{\mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets}(\underline{K}, \langle A_\bullet \rangle) \cong \hom_{\mathrm{CDGA}_{\mathcal{F}}}(A_\bullet, \mathcal{A}(\underline{K})_\bullet).$$

This pair of contravariant adjoint functors extends to a bijection between homotopy classes (see Theorem E)

$$[\underline{K}, \langle A_\bullet \rangle]_{\mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets} \cong [A_\bullet, \mathcal{A}(\underline{K})_\bullet]_{\mathrm{CDGA}_{\mathcal{F}}}.$$

Finally, *in Section 12,* we prove that, for any filtered face set, $\underline{K}$, the perverse algebra $\mathcal{A}(\underline{K})_\bullet$ and its homology, $H(\mathcal{A}(\underline{K})_\bullet)$, admit positively graded minimal models (see Theorem F). We obtain thus a definition of formality: the filtered face set $\underline{K}$ is said *intersection-formal* if there is an isomorphism between the minimal models of $\mathcal{A}(\underline{K})_\bullet$ and $H(\mathcal{A}(\underline{K})_\bullet)$.

Some of our first results are proved in the general case of loose perversities but the construction of the minimal models relies strongly on the structure of the lattice of Goresky and MacPherson perversities. This was first observed by Stienne ([34]) who determined the properties of the predecessors of a given Goresky and MacPherson perversity and a part of the construction of our perverse minimal model is inspired from his work.

We end with the topological invariance of the minimal model. This uses Theorem A and the existence of an intrinsic filtration of a locally conelike space, due to Sullivan, and detailed in [23].

In all this text, for a graded perverse object, a, we denote by $|a|$ its degree and by $\|a\|$ its perverse degree.

Contents

1. Stratified spaces 5
2. Intersection homology of stratified spaces 7
3. Intersection homology of locally conelike spaces 15
4. Filtered face sets 17
5. Local system on filtered face sets 23
6. Proof of Theorem C 28
7. Cochains on filtered face sets 34
8. Particular cases: $\overline{q} = \overline{0}$ and $\overline{q} = \overline{\infty}$ 39

9. Homotopy of filtered face sets 45
10. Perverse differential graded algebras 49
11. Filtered face sets and perverse differential graded algebras 53
12. Minimal models 58
References 71

1. Stratified spaces

We begin with definitions and main properties of stratified spaces and maps. As these notions vary along the literature, the reader should be aware of the definitions. For instance, we quote that, in this work, the strata are the connected components of $X_i \backslash X_{i-1}$.

Among the results of this section, mention the fact that the relation $S \subset \overline{S'}$ defines an order on the set of strata and that a stratifed map respects this order.

Definition 1.1. A *filtered space* is a topological space, X, together with a filtration by closed subspaces

$$X_0 \subseteq X_1 \subseteq \ldots \subseteq X_n = X.$$

The *dimension* of X is n, denoted by $d(X) = n$. The subspace X_i is called the *i-skeleton* of X but the index i and the dimension, $d(X)$, are not related to a notion of geometrical dimension.

The connected components, S, of $X_i \backslash X_{i-1}$ are called *strata* of X and we set $d(S) = i$. We denote by $\mathcal{S}_X$ the family of non-empty strata of X.

The subspace X_{n-1} is the *singular set* and is also denoted by Σ. By convention, we set $X_j = \emptyset$ if $j < 0$ and $X_j = X$ if $j > n$. The dimension of the empty set is equal to -1. The strata in $X_n \backslash X_{n-1}$ are called the *regular strata* of X. *We always suppose that $X_n \backslash X_{n-1}$ is not empty.*

The filtration of a topological space X by $X_{n-1} = \emptyset$ and $X_n = X$ is called *trivial* of dimension n.

If $(X, (X_i)_{0 \leq i \leq n})$ and $(Y, (Y_i)_{0 \leq i \leq m})$ are filtered spaces, we define a canonical filtration on the product $X \times Y$ by

$$(X \times Y)_i = \bigcup_{k+j=i} X_k \times Y_j, \text{ for } i \in \{0, \ldots, n+m\}.$$

Definition 1.2. Let $(X, (X_i)_{0 \leq i \leq n})$ be a filtered space, the open cone on X is the quotient $\mathring{c}X = X \times [0,1[/X \times \{0\}$. We denote by ϑ the summit of the cone. The *conic filtration,* $(\mathring{c}X)_{0 \leq i \leq n+1}$, on the open cone is defined by $(\mathring{c}X)_i = \mathring{c}X_{i-1}$, with the convention $\mathring{c}\emptyset = \{\vartheta\}$. Observe that $\mathring{c}X \backslash \{\vartheta\}$ is the product $X \times]0,1[$, where $]0,1[$ owns the trivial filtration of dimension 1.

This conic filtration generates a canonical filtration on the suspension ΣX. The join $X * Y$ of two filtered spaces is also canonically filtered by observing that $X * Y$ is the union of three open sets, $X \times Y \times]0,1[$, $X \times \mathring{c}Y$, $\mathring{c}X \times Y$, and the filtrations on the respective intersections coincide.

Definition 1.3. A *stratified space* is a filtered space such that any pair of strata, S and S' with $S \cap \overline{S'} \neq \emptyset$, verify $S \subset \overline{S'}$.

For instance, the CS sets of King [23] are stratified spaces, see Theorem B and Definition 3.1. Also, the previous constructions of cone, suspension, product and join of stratified spaces are stratified.

Proposition 1.4. *If $(X, (X_i)_{0 \leq i \leq n})$ is a stratified space, the following properties are satisfied.*

(i) If we have $S \subset \overline{S'}$ and $S' \subset \overline{S}$, then $S = S'$.
(ii) The relation $S \preceq S'$ defined on the set of strata by $S \subset \overline{S'}$ is an order relation.
(iii) The dimension map, $d \colon \mathcal{S}_X \to \mathbb{N}$, is strictly increasing.
(iv) The closure of a stratum is a union of lower strata, i.e., $\overline{S} = \cup_{S' \preceq S} S'$.

Proof. (i) Suppose $d(S) \leq d(S')$. Since each stratum, S, is closed in $X_{d(S)} \backslash X_{d(S)-1}$ and $X_{d(S)}$ is closed in X, the stratum S is closed in $X \backslash X_{d(S)-1}$, which implies

$$S = \overline{S} \cap (X \backslash X_{d(S)-1}).$$

Then, the inclusion $S' \subset \overline{S}$ implies $S' \backslash X_{d(S)-1} \subset S$. The inclusion $S' \cap X_{d(S)-1} \subset S' \cap X_{d(S')-1}$ and the equality $S' \cap X_{d(S')-1} = \emptyset$, by definition of $d(S')$, imply $S' \subset S$ and the equality $S' = S$.

(ii) This is a direct consequence of the previous property, the reflexivity and transitivity being obvious.

(iii) Suppose $S \preceq S'$. As $X_{d(S')}$ is closed, we have $S \subset \overline{S'} \subset X_{d(S')}$ which implies $d(S) \leq d(S')$. Since $S \subset X \backslash X_{d(S)-1}$, we have

$$S = S \cap (X \backslash X_{d(S)-1}) \subset \overline{S'} \cap (X \backslash X_{d(S)-1}).$$

If $d(S) = d(S')$, we know that S' is closed in $X \backslash X_{d(S)-1}$, which implies

$$S' = \overline{S'} \cap (X \backslash X_{d(S)-1}).$$

We get $S \subset S'$ and $S = S'$.

(iv) The inclusion $\cup_{S' \preceq S} S' \subset \overline{S}$ is obvious. Consider $x \in \overline{S}$. Since the strata form a partition, there exists a unique stratum S' such that $x \in S'$. From $\overline{S} \cap S' \neq \emptyset$ and from Definition 1.3, we deduce $S' \subset \overline{S}$. We get $x \in S'$ and $S' \preceq S$, which proves the remaining part of the equality.

□

Definition 1.5. A *filtered map, $f \colon (X, (X_j)_{0 \leq j \leq n}) \to (Y, (Y_j)_{0 \leq j \leq m})$*, is a continuous map, $f \colon X \to Y$, such that $f^{-1} Y_{m-\ell} \subseteq X_{n-\ell}$, for any $\ell \geq 0$. A *stratified map* is a filtered map between stratified spaces, such that, for any stratum $S \in \mathcal{S}_X$, there exists a stratum $S^f \in \mathcal{S}_Y$ with $f(S) \subset S^f$. A stratified map is *stratum preserving* if $n = m$ and $f^{-1} Y_{n-\ell} = X_{n-\ell}$, for any $\ell \geq 0$.

The previous conditions on stratum-preserving map correspond to definitions of Friedman [11] and Quinn, [29], [28]. Filtered maps and stratified maps are stable by composition. Observe also that a filtered map is stratified if, and only if, for any stratum $S' \in \mathcal{S}_Y$, there exists a family of strata, $\{S_i \mid i \in I\} \subset \mathcal{S}_X$, such that

$$f^{-1}(S') = \bigcup_{i\in I} S_i.$$

For instance, the following maps are stratified:

- the canonical projections $X \times Y \to X$, $X \times Y \to Y$,
- the product of two stratified maps.
- If $f\colon Z \to X$ and $g\colon Z \to Y$ are stratified, then $(f,g)\colon Z \to X \times Y$ is stratified also.

As strata give a partition, the stratum S^f of Definition 1.5 is uniquely determined by f and S. Therefore we get a map $\mathcal{S}_f$ between the sets of strata.

Proposition 1.6. *If $f\colon X \to Y$ is a stratified map, the induced map, $\mathcal{S}_f\colon(\mathcal{S}_X, \preceq) \to (\mathcal{S}_Y, \preceq)$, defined by $\mathcal{S}_f(S) = S^f$, is increasing.*

Proof. Let $S_1 \preceq S_2$ in $\mathcal{S}_X$. Since f is continuous and $S_1 \subset \overline{S_2}$, we have $f(S_1) \subset \overline{f(S_2)}$. This implies $S_1^f \cap \overline{S_2^f} \neq \emptyset$, which gives $\mathcal{S}_f(S_1) \preceq \mathcal{S}_f(S_2)$. ☐

2. Intersection homology of stratified spaces

We define filtered simplices of a filtered space and the notion of perverse degree of a filtered simplex. The key point is that stratified maps decrease the perverse degree, cf. Theorem A. If $\overline{p}$ is a loose perversity, we introduce a chain complex, $C_*^{\overline{p}}(X)$, associated to a stratified space and well adapted to the simplicial point of view developed in Section 4. Its homology is called *intersection homology* and denoted $H_*^{\overline{p}}(X)$. We show that stratified maps induce homomorphisms in intersection homology. We define also a notion of s-homotopy between stratified maps and prove that two s-homotopic stratified maps induce the same homomorphism in intersection homology. Finally, we establish some properties of the intersection homology of stratified spaces, as a Mayer-Vietoris sequence and a particular case of a Künneth formula.

Definition 2.1. Let $(X, (X_i)_{0\leq i\leq n})$ be a filtered space and Δ be a simplex. A *filtered simplex* is a continous map, $\sigma\colon \Delta \to X$, such that one of the following equivalent properties is satisfied.

(a) For all $i \in \{0, \dots, n\}$, $\sigma^{-1}X_i$ is a face of Δ or $\sigma^{-1}X_i = \emptyset$.

(b) There exists a decomposition $\Delta = \Delta^{j_0} * \Delta^{j_1} * \dots * \Delta^{j_n}$ such that

$$\sigma^{-1}X_i = \Delta^{j_0} * \Delta^{j_1} * \dots * \Delta^{j_i},$$

for all $i \in \{0, \dots, n\}$. We call it *the σ-decomposition of* Δ.

We denote by $C_*(X)$ the usual singular chain complex of X and by $C_*^{\mathcal{F}}(X)$ the subcomplex generated by the filtered simplices.

In this definition, the simplices Δ^{j_i} can be empty. An empty set appears automatically if the filtration of X is stationary for an index i. We use the convention $\emptyset * Y = Y$, for any space Y.

Definition 2.2. Let $(X,(X_i)_{0\leq i\leq n})$ be a filtered space and let $\sigma\colon \Delta \to X$ be a filtered simplex. The *perverse degree* of σ is the $(n+1)$-uple

$$\|\sigma\| = (\|\sigma\|_0, \ldots, \|\sigma\|_n),$$

where $\|\sigma\|_\ell$ is the dimension of the smallest skeleton containing $\sigma^{-1}X_{n-\ell}$, with the convention $\|\sigma\|_\ell = -\infty$ if $\sigma^{-1}X_{n-\ell} = \emptyset$.
If $\sigma\colon \Delta = \Delta^{j_0} * \cdots * \Delta^{j_n} \to X$ is such that $\Delta^{j_0} * \cdots * \Delta^{j_{n-\ell}} \neq \emptyset$ then $\|\sigma\|_\ell = \dim \sigma^{-1}X_{n-\ell} = \dim(\Delta^{j_0} * \cdots * \Delta^{j_{n-\ell}})$.

Theorem A. *A stratified map, $f\colon X \to Y$, induces a chain map $f_*\colon C_*^{\mathcal{F}}(X) \to C_*^{\mathcal{F}}(Y)$, defined by $\sigma \mapsto f\circ\sigma$. Moreover, for all $\ell \geq 0$, we have*

$$\|f_*(\sigma)\|_\ell \leq \|\sigma\|_\ell.$$

Before proving this property, we need results on the behavior of the image of strata by a filtered simplex.

Lemma 2.3. *Let $(X,(X_i)_{0\leq i\leq n})$ be a stratified space and let $\sigma\colon \Delta \to X$ be a filtered simplex. Then the following properties are satisfied.*

- *(i) For any face ∇ of Δ, the restriction $\sigma_{|\nabla}$ is a filtered simplex.*
- *(ii) For any face ∇ of Δ, there is a stratum $S \in \mathcal{S}_X$ such that $\sigma(\mathring{\nabla}) \subset S$.*
- *(iii) The family of strata that have a non-empty intersection with the image of σ is totally ordered. If $S_1 \prec S_2 \prec \cdots \prec S_p$ is this family, then we have*

$$\sigma^{-1}X_i = \begin{cases} \emptyset & \text{if } i < d(S_1), \\ \sigma^{-1}(S_1 \sqcup \cdots \sqcup S_k) & \text{if } d(S_k) \leq i < d(S_{k+1}),\ 1 \leq k < p, \\ \sigma^{-1}(S_1 \sqcup \cdots \sqcup S_p) & \text{if } i \geq d(S_p). \end{cases} \tag{1}$$

Proof. The first property is obvious and it is sufficient to prove the second one for $\nabla = \Delta$. Let k be the integer such that $\sigma(\Delta) \subset X_k$ and $\sigma(\Delta) \not\subset X_{k-1}$. Thus there exists $t \in \Delta$ with $\sigma(t) \notin X_{k-1}$ and we consider the stratum, S, characterized by $\sigma(t) \in S \subset X_k \backslash X_{k-1}$. By hypothesis, the pullback $\Delta' = \sigma^{-1}(X_{k-1})$ is a face (perhaps empty) of Δ. We have $\Delta' \neq \Delta$ because $t \notin \Delta'$. From $\sigma(\Delta\backslash\Delta') \subset X_k\backslash X_{k-1}$ and $\sigma(t) \in \sigma(\Delta\backslash\Delta') \cap S$, we deduce $\sigma(\Delta\backslash\Delta') \subset S$ and

$$\sigma(\mathring{\Delta}) \subset \sigma(\Delta\backslash\Delta') \subset S.$$

(iii) Let $i \in \{0,\ldots,n\}$. From the definition of filtered simplex, we know that $\sigma^{-1}X_i\backslash\sigma^{-1}X_{i-1}$ is connected. Moreover, this complement can be decomposed in a disjoint union of closed subsets of $\sigma^{-1}X_i\backslash\sigma^{-1}X_{i-1}$,

$$\sigma^{-1}X_i\backslash\sigma^{-1}X_{i-1} = \bigsqcup_{d(S)=i} \sigma^{-1}S.$$

Suppose $\sigma^{-1}X_i\backslash\sigma^{-1}X_{i-1} \neq \emptyset$. As it is connected, it is the pullback of *one* stratum $\mathtt{S}_i$ of $X_i\backslash X_{i-1}$. Therefore, there exists a family of integers, $0 \leq d_1 < d_2 < \cdots < d_p \leq n$, such that

$$\sigma^{-1}X_i\backslash\sigma^{-1}X_{i-1} = \begin{cases} \emptyset & \text{if } \ i \notin \{d_1,\ldots,d_p\}, \\ \sigma^{-1}(\mathtt{S}_i) & \text{if } \ i = d_k \text{ for some } k \in \{1,\ldots,p\}. \end{cases}$$

Setting $\mathsf{S}_{d_k} = S_k$, this gives the equalities (1).

Let $k \in \{2, \dots, p\}$, we are reduced to prove $S_{k-1} \prec S_k$. Recall the equality $\sigma^{-1}X_{d_k} = \sigma^{-1}(S_1 \sqcup \dots \sqcup S_k)$. We have proved $\sigma^{-1}X_{d_k}\backslash\sigma^{-1}X_{d_{k-2}} = \sigma^{-1}(S_{k-1} \sqcup S_k)$ and $\sigma^{-1}\overline{S_k} = \bigcup\limits_{S\preceq S_k} \sigma^{-1}S$, see (iv) of Proposition 1.4. In particular, $\sigma^{-1}(\overline{S_k} \cap (X_{d_{k-1}}\backslash X_{d_{k-2}})) = \sigma^{-1}S_{k-1}$ or $\emptyset$ and we have

$$\begin{aligned}
\sigma^{-1}X_{d_k}\backslash\sigma^{-1}X_{d_{k-2}} &= \sigma^{-1}(S_{k-1} \sqcup (\overline{S_k} \cap (X_{d_k}\backslash X_{d_{k-1}})))\\
&= \sigma^{-1}(S_{k-1} \cup (\overline{S_k} \cap (X_{d_k}\backslash X_{d_{k-2}})))\\
&= \sigma^{-1}(S_{k-1}) \cup \sigma^{-1}(\overline{S_k} \cap (X_{d_k}\backslash X_{d_{k-2}})).
\end{aligned}$$

The subset $\sigma^{-1}S_{k-1}$(resp. $\sigma^{-1}(\overline{S_k} \cap (X_{d_k}\backslash X_{d_{k-2}}))$ is closed in the subset $\sigma^{-1}X_{d_{k-1}}\backslash\sigma^{-1}X_{d_{k-2}}$ (resp. $\sigma^{-1}X_{d_k}\backslash\sigma^{-1}X_{d_{k-2}}$), which is closed in the connected space $\sigma^{-1}X_{d_k}\backslash\sigma^{-1}X_{d_{k-2}}$.

From the formula above and the connectivity of $\sigma^{-1}X_{d_k}\backslash\sigma^{-1}X_{d_{k-2}}$, we deduce $\sigma^{-1}(S_{k-1}) \cap \sigma^{-1}(\overline{S_k} \cap (X_{d_k}\backslash X_{d_{k-2}})) \neq \emptyset$ which implies $S_{k-1} \cap \overline{S_k} \neq \emptyset$ and $S_{k-1} \preceq S_k$, by definition. Finally, as $d(S_{k-1}) = d_{k-1} \neq d_k = d(S_k)$, we have $S_{k-1} \prec S_k$. □

Proof of Theorem A. Let $f\colon (X, (X_i)_{0\leq i\leq n}) \to (Y, (Y_i)_{0\leq i\leq m})$ be a stratified map.

If $(f \circ \sigma)^{-1}(Y_{m-\ell}) = \emptyset$ then $\|f \circ \sigma\|_\ell = -\infty$.

If $\sigma^{-1}X_{n-\ell} = \emptyset$, then from $(f \circ \sigma)^{-1}(Y_{m-\ell}) \subset \sigma^{-1}X_{n-\ell} = \emptyset$, we deduce $\|f \circ \sigma\|_\ell = \|\sigma\|_\ell = -\infty$.

We may thus suppose $(f \circ \sigma)^{-1}(Y_{m-\ell}) \neq \emptyset$ and $\sigma^{-1}X_{n-\ell} \neq \emptyset$, which implies $0 \leq \ell \leq m$. We have to prove that $(f \circ \sigma)^{-1}(Y_{m-\ell})$ is a face of Δ and that

$$\dim(f \circ \sigma)^{-1}(Y_{m-\ell}) \leq \dim \sigma^{-1}(X_{n-\ell}).$$

Recall that the family of strata meeting the image of σ can be ordered as

$$S_1 \prec S_2 \prec \dots \prec S_p,$$

which implies, with Proposition 1.6,

$$S_1^f \preceq S_2^f \preceq \dots \preceq S_p^f.$$

Let T be a stratum of Y such that $T \cap (f \circ \sigma)(\Delta) \neq \emptyset$., i.e., $f^{-1}(T) \cap \sigma(\Delta) \neq \emptyset$. As the map f is stratified, there exists $k \in \{1, \dots, p\}$, such that $T = S_k^f$. Since f is filtered, we have $S_k \subset X_{n-(m-d(T))}$ and

$$m - d(T) \leq n - d(S_k).$$

We denote by $0 \leq k_0 < \dots < k_q = p$ the integers such that

$$\begin{cases}
T_1 = S_1^f = \dots = S_{k_1}^f, \text{ with } S_{k_1}^f \neq S_{k_1+1}^f,\\
T_2 = S_{k_1+1}^f = \dots = S_{k_2}^f, \text{ with } S_{k_2}^f \neq S_{k_2+1}^f,\\
\dots\\
T_q = S_{k_{q-1}+1}^f = \dots = S_{k_q}^f = S_p^f.
\end{cases}$$

From that, we deduce

(1) $\{T \in \mathcal{S}_Y \mid T \cap (f \circ \sigma)(\Delta) \neq \emptyset\} = \{T_1 \prec T_2 \prec \dots \prec T_q\}$,
(2) $m - d(T_j) \leq n - d(S_{k_j})$,

(3) $f^{-1}(T_j) = \displaystyle\bigcup_{k=k_{j-1}+1}^{k_j} S_k$, for any $j \in \{1,\dots,q\}$.

From (iii) of Proposition 1.4, we deduce

$$0 \le d(T_1) < d(T_2) < \cdots < d(T_q) \le m.$$

With the convention $d(T_{q+1}) = m$ and $d(T_0) = 0$, we denote by $a \in \{0,\dots,q\}$ the integer such that

$$d(T_a) \le m - \ell < d(T_{a+1}).$$

As $(f\circ\sigma)^{-1}Y_{m-\ell} \neq \emptyset$, the integer a is different of 0 and we have obtained

$$\begin{aligned}(f\circ\sigma)^{-1}(Y_{m-\ell}) &= \bigcup_{j=0}^{m-\ell} \sigma^{-1}f^{-1}(Y_j\backslash Y_{j-1}) = \bigcup_{i=0}^{a}\sigma^{-1}f^{-1}(T_i)\\ &= \bigcup_{k=1}^{k_a}\sigma^{-1}(S_k) = \sigma^{-1}(X_{d(S_{k_a})}),\end{aligned}$$

which is a face of the simplex Δ. The simplex $f\circ\sigma$ is thus filtered. From Property (2) above, we deduce

$$d(S_{k_a}) \le n - m + d(T_a) \le n - m + (m-\ell) \le n - \ell,$$

which implies $\sigma^{-1}(X_{d(S_{k_a})}) \subset \sigma^{-1}(X_{n-\ell})$ and the expected inequality between the dimensions,

$$\dim (f\circ\sigma)^{-1}(Y_{m-\ell}) = \dim\sigma^{-1}(X_{d(S_{k_a})}) \le \dim\sigma^{-1}(X_{n-\ell}).$$

□

Remark 2.4. In the previous proof, the simplex Δ owns two decompositions, one induced by the map $\sigma\colon \Delta \to X$ and the other one by $f\circ\sigma\colon\Delta\to Y$. In general, they do not coincide, the second one being an amalgamation of terms of the first one. More precisely, if $\Delta = \Delta_1 * \Delta_2 * \cdots * \Delta_p$ is the σ-decomposition of Δ, the $(f\circ\sigma)$-decomposition of Δ is of the shape $\Delta = \Delta'_1 * \Delta'_2 * \cdots * \Delta'_q$, with $\Delta'_a = \Delta_{k_{a-1}+1} * \cdots * \Delta_{k_a}$, for any $a \in \{1,\dots,q\}$. With the notation of the proof, this decomposition verifies

$$(f\circ\sigma)^{-1}(T_1 \sqcup \cdots \sqcup T_a) = \Delta'_1 * \cdots * \Delta'_a, \text{ for any } a \in \{1,\dots,q\}.$$

Consider the case of a stratum-preserving stratified map. With property (iii) of Proposition 1.4, the relation $S_1 \prec S_2 \prec \cdots \prec S_p$ and the stratum-preserving condition imply $S_1^f \prec S_2^f \prec \cdots \prec S_p^f$. Therefore the equation (3) above reduces to $f^{-1}(T_j) = S_{k_j}$ and we have $(f\circ\sigma)^{-1}(T_j) = \sigma^{-1}(S_{k_j})$. In summary, *in the case of a stratum-preserving stratified map, the two decompositions are identical.*

We introduce now the notion of perversity which is the fundamental tool of intersection theory. We follow the convention of [23].

Definition 2.5. A *loose perversity* is a map $\overline{p}\colon \mathbb{N} \to \mathbb{Z}$, $i \mapsto \overline{p}(i)$, such that $\overline{p}(0) = 0$. A loose perversity is *positive* if $\overline{p}(i) \ge 0$, for any $i \in \mathbb{N}$.

A *perversity* is a loose perversity such that $\overline{p}(i) \le \overline{p}(i+1) \le \overline{p}(i) + 1$, for all $i \in \mathbb{N}$. A *Goresky-MacPherson perversity* (or *GM-perversity*) is a perversity such that $\overline{p}(1) = \overline{p}(2) = 0$.

If $\overline{p}_1$ and $\overline{p}_2$ are two loose perversities, we set $\overline{p}_1 \leq \overline{p}_2$ if we have $\overline{p}_1(i) \leq \overline{p}_2(i)$, for all $i \in \mathbb{N}$. The lattice of GM-perversities, denoted $\mathcal{P}^n$, admits a maximal element, $\overline{t}$, called the *top perversity* and defined by $\overline{t}(i) = i - 2$, if $i \geq 2$.

To these perversities, we add an element, $\overline{\infty}$, which is the constant map on ∞. We call it the *infinite perversity* despite the fact that it is not a perversity in the sense of the previous definition.

Definition 2.6. ([16]) Let $(X, (X_i)_{0 \leq i \leq n})$ be a filtered space and $\overline{p}$ be a loose perversity. A *$\overline{p}$-admissible simplex* is a filtered simplex, $\sigma\colon \Delta \to X$, such that

$$\|\sigma\|_i = \dim \sigma^{-1} X_{n-i} \leq \dim \Delta - i + \overline{p}(i),$$

for all $i \in \{0, \ldots, n\}$. A *chain c is $\overline{p}$-admissible* if there exist $\overline{p}$-admissible simplices, σ_j, so that $c = \sum_j \lambda_j \sigma_j$, $\lambda_j \in \mathbb{Z}$.

Remark 2.7. A $\overline{p}$-admissible simplex, $\sigma\colon \Delta \to \underline{K}$, verifies

$$\dim \Delta = \|\sigma\|_0 \leq \dim \Delta - 0 + \overline{p}(0).$$

In the definition of loose perversity, we have imposed $\overline{p}(0) = 0$ and this restriction makes empty the condition on $\|\sigma\|_0$. On the other side, we may observe also that, with a negative value for $\overline{p}(0)$, we should not have any $\overline{p}$-admissible simplex.

As usual in this theory, the abelian free group generated by the admissible $\overline{p}$-chains is not a sub-complex of the singular chains and we need to set the following definition.

Definition 2.8. Let $(X, (X_i)_{0 \leq i \leq n})$ be a filtered space and $\overline{p}$ be a loose perversity. A chain c is *an intersection chain* for $\overline{p}$ if c and its boundary ∂c are $\overline{p}$-admissible chains. We denote by $C_*^{\overline{p}}(X)$ the complex of intersection chains. The associated homology and cohomology are denoted by $H_*^{\overline{p}}(X)$ and $H^*_{\overline{p}}(X)$ and called *intersection homology and cohomology for $\overline{p}$.*

In the sequel, the notation $H_*^{\overline{p}}(X)$ and $H^*_{\overline{p}}(X)$ mean with coefficients in $\mathbb{Z}$. In the other cases, we mention explicitly the ring of coefficients; for instance, rational intersection cohomology is denoted by $H^*_{\overline{p}}(X; \mathbb{Q})$. The next result is a direct consequence of Definition 2.6 and Theorem A.

Proposition 2.9. *A stratified map, $f\colon X \to Y$, induces a morphism between the complexes of intersection chains, $f_*\colon C_*^{\overline{p}}(X) \to C_*^{\overline{p}}(Y)$, for any loose perversity $\overline{p}$.*

In the next definition, $[0, 1]$ is trivially filtered of dimension 1 and $X \times [0, 1]$ endows the product filtration.

Definition 2.10. Two stratified maps, f_0, $f_1\colon X \to Y$, are *s-homotopic* if there exists a stratified map, $f\colon X \times [0, 1] \to Y$, such that $f(x, 0) = f_0(x)$ and $f(x, 1) = f_1(x)$.

Proposition 2.11. *If f_0, $f_1\colon X \to Y$ are two s-homotopic stratified maps, they induced the same homomorphism in intersection homology,*

$$H_*^{\overline{p}}(f_0) = H_*^{\overline{p}}(f_1)\colon H_*^{\overline{p}}(X) \to H_*^{\overline{p}}(Y),$$

for any loose perversity $\overline{p}$.

Proof. Let $f\colon X\times I\to Y$ be an s-homotopy from f_0 to f_1 and $\iota_k\colon X\to X\times I$, $x\mapsto(x,k)$, the canonical inclusions for $k=0,1$. In the non-stratified setting, the proof (see [10, Théorème 5.33] for instance) uses a chain homotopy between $C_*(\iota_0)$ and $C_*(\iota_1)$, defined as follows.

If $\Delta^n=\langle e_0,\dots,e_n\rangle$, we denote by $a_i=(e_i,0)$ and $b_i=(e_i,1)$ the points of $\Delta^n\times I$ corresponding to vertices. A singular $(n+1)$-chain of $\Delta^n\times I$ is defined by

$$P=\sum_{i=0}^{n}(-1)^i\langle a_0,\dots,a_i,b_i,\dots,b_{n+1}\rangle.$$

This chain generates a homotopy $h\colon C_*(X)\to C_*(X\times I)$ defined by $h_*(\sigma)=C_*(\sigma\times\mathrm{id})(P)$ and which satisfies the expected equality $(dh+hd)(\sigma)=C_*(\iota_1)(\sigma)-C_*(\iota_0)(\sigma)$.

Let σ be a $\overline{p}$-admissible simplex of X, we are reduced to prove that $h(\sigma)$ is a $\overline{p}$-admissible simplex of $X\times[0,1]$. Let $j\in\{0,\dots,n\}$. We denote by $\tau_j\colon\Delta^{n+1}=\langle v_0,\dots,v_{n+1}\rangle\to\Delta^{n+1}\times I$ the $(n+1)$-simplex defined by $(v_0,\dots,v_j,v_{j+1},\dots,v_{n+1})\mapsto(a_0,\dots,a_j,b_j,\dots,b_n)$. It is sufficient to prove that τ_j is such that $\dim\tau_j^{-1}(F)\le\dim F$, for any face, F, of the simplicial decomposition

$$\Delta^n\times I=\cup_{i=0}^{n}\tau_i(\Delta^{n+1}).$$

Therefore, $\tau_j^{-1}(F)$ is identified to $F\cap\tau_j(\Delta^{n+1})$ and the result follows. □

In the end of this section, we establish basic properties of the intersection homology.

Proposition 2.12. *Let X be a filtered space with $d(X)=n$ and $\overline{p}$ be a loose perversity. Then the following properties are satisfied.*

(i) *If $\mathfrak{U}=\{U,V\}$ is an open cover of X, there exists a long exact sequence*

$$\ldots\to H_*^{\overline{p}}(U\cap V)\to H_*^{\overline{p}}(U)\oplus H_*^{\overline{p}}(V)\to H_*^{\overline{p}}(X)\to H_{*-1}^{\overline{p}}(U\cap V)\to\ldots$$

(ii) *The canonical projection* $\mathrm{pr}\colon\mathbb{R}\times X\to X$ *induces an isomorphism*

$$H_*^{\overline{p}}(\mathrm{pr})\colon H_*^{\overline{p}}(\mathbb{R}\times X)\to H_*^{\overline{p}}(X),$$

whose inverse is induced by the inclusion $X\to\mathbb{R}\times X$, $x\mapsto(0,x)$.

(iii) *If $U_0\subset U_1\subset\cdots$ are open sets of X so that $X=\cup_iU_i$, then the natural map $\lim_{\to}H_*^{\overline{p}}(U_i)\to H_*^{\overline{p}}(X)$ is an isomorphism.*

(iv) *If M is a topological manifold, then the following sequence is split exact for each i,*

$$0\longrightarrow(H_*(M)\otimes H_*^{\overline{p}}(X))_i\longrightarrow H_i^{\overline{p}}(X)\longrightarrow(H_*(M)*H_*^{\overline{p}}(X))_{i-1}\longrightarrow 0.$$

(v) *Suppose X is compact and $\overline{p}$ is a perversity. If the cone $\mathring{c}X$ is endowed with the conic filtration, we have*

$$H_k^{\overline{p}}(\mathring{c}X)=\begin{cases}H_k^{\overline{p}}(X) & \text{if } k\le n-\overline{p}(n+1)-1,\\ 0 & \text{if } k\ge n-\overline{p}(n+1) \text{ and } k\ne0,\\ \mathbb{Z} & \text{if } k\ge n-\overline{p}(n+1) \text{ and } k=0.\end{cases}$$

and

$$H_k^{\overline{p}}(\mathring{c}X, X\times]0,1[) = \begin{cases} 0 & \text{if } k \leq n-\overline{p}(n+1), \\ \overline{H}^{\overline{p}}_{k-1}(X) & \text{if } k \geq n-\overline{p}(n+1)+1, \end{cases}$$

where $\overline{H}^{\overline{p}}_*$ *denotes the reduced homology.*

Proof. (i) This Mayer-Vietoris exact sequence is a direct consequence of Lemma 2.13, as in the classical case.

(ii) This is a consequence of Proposition 2.11.

(iii) This is obvious.

(iv) A careful reading of the proof of [23, Theorem 4] shows that this particular Künneth formula exists if properties (i), (ii) and (iii) are satisfied.

(v) Let X be a compact filtered space, $\mathring{c}X$ the associated open cone, endowed with the conic filtration. The next computation is already done in previous works.

Let $\sigma\colon \Delta^k \to \mathring{c}X$ be a $\overline{p}$-admissible filtered simplex. We consider two cases.

• *Let* $k = \dim\Delta \leq n-\overline{p}(n+1)$. As

$$\|\sigma\|_{n+1} \leq k-(n+1)+\overline{p}(n+1) < 0,$$

we have $\sigma^{-1}(\mathring{c}X)_0 = \emptyset$ and the simplex σ does not meet the summit ϑ of the cone $\mathring{c}X$. This implies $C^{\overline{p}}_{\leq n-\overline{p}(n+1)}(\mathring{c}X) = C^{\overline{p}}_{\leq n-\overline{p}(n+1)}(X\times]0,1[)$ and property (ii) implies

$$H_k^{\overline{p}}(\mathring{c}X) = H_k^{\overline{p}}(X\times]0,1[) = H_k^{\overline{p}}(X),$$

for all $k \leq n-\overline{p}(n+1)-1$.

• *Let* $k = \dim\Delta \geq n-\overline{p}(n+1)$. In this case, we show that the classical homotopy operator, used to prove the contractibility of the singular chain complex of a cone, is compatible with the intersection setting. If $s\in[0,1]$ and $\langle x,t\rangle \in \mathring{c}X$, we set $s\cdot\langle x,t\rangle = \langle x,st\rangle$. If $\sigma\colon\Delta\to\mathring{c}X$ is a simplex, we define $c\sigma\colon \Delta * \{\vartheta\} \to \mathring{c}X$ by

$$c\sigma(sa+(1-s)\vartheta) = s\cdot\sigma(a).$$

As X is compact, this map is continue and we extend it in a linear map $c\sigma\colon C_*(\mathring{c}X)\to C_*(\mathring{c}X)$. Let $i\in\{0,\dots,n+1\}$ and $\sigma\colon\Delta\to\mathring{c}X$ be a $\overline{p}$-admissible simplex. We have

$$\begin{aligned}(c\sigma)^{-1}((\mathring{c}X)_{n+1-i})) &= \{sa+(1-s)\vartheta \mid s\cdot\sigma(a)\in(\mathring{c}X)_{n+1-i} = \mathring{c}X_{n-i}\} \\ &= \sigma^{-1}(\mathring{c}X_{n-i}) * \{\vartheta\},\end{aligned}$$

which is a face of Δ. If $\sigma^{-1}(\mathring{c}X_{n-1}) \neq \emptyset$, we have

$$\begin{aligned}\dim(c\sigma)^{-1}((\mathring{c}X)_{n+1-i}) &= 1+\dim\sigma^{-1}(\mathring{c}X_{n-i}) \leq 1+\dim\Delta - i+\overline{p}(i) \\ &= \dim(\Delta * \{\vartheta\}) - i + \overline{p}(i).\end{aligned}$$

Therefore, the simplex $c\sigma$ is $\overline{p}$-admissible.

If $\sigma^{-1}(\mathring{c}X_{n-1}) = \emptyset$, then $\dim(c\sigma)^{-1}((\mathring{c}X)_{n+1-i}) = 0$ and, by hypothesis, we have

$$\begin{aligned}0 &\leq k-n+\overline{p}(n+1) = k+1-(n+1)+\overline{p}(n+1) \\ &\leq k+1-i+\overline{p}(i),\end{aligned}$$

because $\overline{p}$ is a perversity. The last expression is equal to $\dim(\Delta * \{\vartheta\}) - i + \overline{p}(i)$ and the simplex $c\sigma$ is $\overline{p}$-admissible.

Now we compute $H_k^{\overline{p}}(\mathring{c}X)$ by distinguishing two cases.

- — $k \neq 0$. The result is clear for $k < 0$ and we may suppose $k > 0$. For any $\eta \in C_k^{\overline{p}}(\mathring{c}X)$, we have $\partial c\eta + c\partial\eta = \eta$, which implies $H_k^{\overline{p}}(\mathring{c}X) = 0$.
- — $k = 0$. For any $\eta \in C_0^{\overline{p}}(\mathring{c}X)$, we have the formula $\partial c\eta = \eta - \vartheta$. This implies that all the generators of $C_0^{\overline{p}}(\mathring{c}X)$ are identified in homology and, therefore, $H_0^{\overline{p}}(\mathring{c}X) = \mathbb{Z}$.

Finally, the computation of the relative homology, $H_k^{\overline{p}}(\mathring{c}X, X \times]0,1[)$, is direct from the previous result and the long exact sequence associated to $(\mathring{c}X, X \times]0,1[)$. □

Lemma 2.13. *Let $\mathfrak{U}$ be an open cover of the filtered space X and $\overline{p}$ be a loose perversity. Let $I^{\mathfrak{U}}C_*^{\overline{p}}(X)$ be the subcomplex of $C_*^{\overline{p}}(X)$ whose elements have a support included in an element of $\mathfrak{U}$. We denote its homology by $I^{\mathfrak{U}}H_*^{\overline{p}}(X)$. Then the map $I^{\mathfrak{U}}H_*^{\overline{p}}(X) \to H_*^{\overline{p}}(X)$, generated by the inclusion, is an isomorphism.*

Proof. We adapt the classical proof to the intersection setting. For that, we take over the presentation of [33, Section 4 of Chapter 4]. The key is the construction of an augmentation-preserving chain map $sd\colon C_*(X) \to C_*(X)$ and a chain deformation $T\colon C_*(X) \to C_*(X)$ from sd to id, where $C_*(X)$ denotes the singular chain complex of X.

We define now sd and T in the intersection setting. Denote by $\Delta^{per}(\Delta)$ the subcomplex generated by the linear simplices, ξ, of Δ such that

$$\dim \xi^{-1}(F) \leq \dim F,$$

for any face F of Δ. Observe that if $\sigma\colon \Delta \to X$ is filtered and $\xi \in \Delta^{per}(\Delta)$, then

- $\sigma_*(\xi)^{-1}(X_i) = \xi^{-1}\sigma^{-1}(X_i)$ is a face of Δ because σ is filtered and ξ linear,
- $\|\sigma_*(\xi)\|_i = \dim \xi^{-1}\sigma^{-1}(X_{n-i}) \leq \dim \sigma^{-1}(X_{n-i}) = \|\sigma\|_i$.

Therefore, if σ is $\overline{p}$-admissible and $\xi \in \Delta^{per}(\Delta)$ then $\sigma_*(\xi)$ is $\overline{p}$-admissible. The proof of the statement is reduced to the verification of the inclusions,

$$sd(\Delta^{per}(\Delta)) \subset \Delta^{per}(\Delta) \text{ and } T(\Delta^{per}(\Delta)) \subset \Delta^{per}(\Delta).$$

- Let $\xi \in \Delta_\ell^{per}(\Delta)$, $\xi\colon \nabla \to \Delta$, and recall that sd is defined by

$$sd(\xi) = \left\{ \begin{array}{ll} \xi & \text{if } \ell = -1,\, 0, \\ b(\xi) \cdot sd(\partial\xi) & \text{if } \ell > 0, \end{array} \right.$$

where $b(\xi)$ is the barycenter of ξ and the definition of $b(\xi) \cdot sd(\partial\xi)$ is recalled below. The result is clear for $\ell = -1$, 0; suppose it is true for any simplex of dimension less than, or equal to, $\ell - 1$. Let $\nabla = \Delta^\ell$. By induction, we know that $sd(\partial(\xi)) = \sum_i n_i\tau_i$, with $\tau_i \in \Delta_{\ell-1}^{per}(\Delta)$. Therefore, we have $b(\xi) \cdot sd(\partial(\xi)) = \sum_i n_i\, b(\xi) \cdot \tau_i$, where $b(\xi) \cdot \tau_i$ is defined on $\nabla = \{\vartheta\} * \Delta^{l-1}$ by

$$(b(\xi) \cdot \tau_i)(t\vartheta + (1-t)a) = tb(\xi) + (1-t)\tau_i(a).$$

We have to prove that $b(\xi) \cdot \tau_i \in \Delta^{per}(\Delta)$. If F is a face of Δ, we claim that $\dim(b(\xi) \cdot \tau_i)^{-1}(F) \leq \dim F$. We consider two cases:

  - $b(\xi)\notin F$. Then $(b(\xi)\cdot\tau_i)^{-1}(F)=\tau_i^{-1}(F)$ and we already know that $\dim\tau_i^{-1}(F)\leq\dim F$.
  - $b(\xi)\in F$. In this case, $\xi(\nabla)\subset F$ which gives $\dim\nabla=\dim\xi^{-1}(F)\leq\dim F$. The inclusion $\xi(\nabla)\subset F$ implies also that $\partial\xi$ is a $(\ell-1)$-simplex of F. Thus $\tau_i(\nabla)\subset F$ and $(b(\xi)\cdot\tau_i)^{-1}(F)=\nabla$. Finally, we obtain $\dim(b(\xi)\cdot\tau_i)^{-1}(F)\leq\dim F$.

- Let $\xi\in\Delta_\ell^{per}(\Delta)$, $\xi\colon\nabla\to\Delta$, and recall that T is defined by

$$T(\xi)=\left\{\begin{array}{ll}\xi & \text{if } \ell=-1,\\ b(\xi)\cdot sd(\xi-T(\partial(\xi))) & \text{if } \ell\geq 0.\end{array}\right.$$

The result is obvious for $\ell=-1$ and the rest of the proof is completely similar at the proof done at the end of the first step.

Let $\sigma\in C_\ell^{\overline{p}}(X)$. The proof ends as in [33] by setting

$$sd(\sigma)=C_*(\sigma)(sd([\Delta^\ell]))\text{ and }T(\sigma)=C_*(\sigma)(T([\Delta^\ell])).$$

□

3. Intersection homology of locally conelike spaces

> This section is the bridge between the intersection homology defined in Section 2 and Goresky-MacPherson original one's. For doing that, we use the CS sets of King (called locally conelike spaces here) and a result of King ([23, Theorem 10]) which gives the isomorphism between King's homology and Goresky-MacPherson homology. We deduce also the independence of the stratification from another theorem of King, ([23, Theorem 9]).

Definition 3.1. A *locally conelike space* of dimension n is a filtered space,

$$\emptyset\subset X_0\subseteq X_1\subseteq\cdots\subseteq X_{n-2}=X_{n-1}\subsetneq X_n=X$$

such that, for all i, $X_i\backslash X_{i-1}$ is a metrizable topological i-dimensional manifold or the empty set, and for each point $x\in X_i\backslash X_{i-1}$, there exist

(a) an open neighborhood, V, of x in X, endowed with the induced filtration,
(b) an open neighborhood, U, of x in $X_i\backslash X_{i-1}$,
(c) a compact, locally conelike space, L, of dimension $n-i-1$, whose cone, $\mathring{c}L$ is endowed with the conic filtration,
(d) a homeomorphism, $\varphi\colon U\times\mathring{c}L\to V$, such that
    (i) $\varphi(u,\vartheta)=u$, for any $u\in U$, with ϑ the summit of the cone,
    (ii) $\varphi(U\times L_j\times]0,1[)=V\cap X_{i+j+1}$, for any $j\in\{0,\dots,n-i-1\}$.

The couple (V,φ) is called a *conic chart* of x and the locally conelike space L the *link* of x.

Observe that this definition makes sense with an induction on the dimension, starting from locally conelike spaces of dimension 0 which are discrete topological spaces, by definition. Also, property (d) (i) implies $V\cap X_{i-1}=\emptyset$ (which implies X_{i-1} closed in X) and both properties (d) (i) and (ii) imply $V\cap X_i=\varphi(U\times\{\vartheta\})=U$. As $X_i\backslash X_{i-1}$ is locally connected, we may choose U connected, i.e., $U=V\cap S$, where S is the stratum containing x.

If $x \in X_n \backslash X_{n-1}$ is a point of the regular part, we choose for L the empty set, $\mathring{c}\emptyset = \vartheta$ and $V = U$, therefore $X_n \backslash X_{n-1}$ is an open subset of X.

Example 3.2. The following spaces are locally conelike:

- any topological manifold, with the trivial filtration,
- any open set of a locally conelike space,
- the product and the join of two locally conelike spaces,
- the open cone of a compact locally conelike space.

Theorem B. *A locally conelike space is a stratified space.*

Proof. Let S and S' be strata such that $S \cap \overline{S'} \neq \emptyset$, we have to prove $S \subset \overline{S'}$. Set $i = d(S)$. The space $X_i \backslash X_{i-1}$ being locally connected, we may cover S by connected open sets U of S satisfying the local condition for some V. Denote by $\mathfrak{U}$ this cover of S and let $U \in \mathfrak{U}$. We first prove the

Claim: if $U \cap \overline{S'} \neq \emptyset$, then $U \subset \overline{S'}$.

Suppose that $U \cap \overline{S'} \neq \emptyset$ and recall $U = V \cap S = \varphi(U \times \{\vartheta\})$ and $V \cap X_{i-1} = \emptyset$. Set $j = d(S')$. The property $U \cap \overline{S'} \neq \emptyset$ implies $V \cap \overline{S'} \neq \emptyset$. As V is open, we deduce $V \cap S' \neq \emptyset$ and $j \geq i$, because $V \cap X_{i-1} = \emptyset$.

• If $i = j$, we have

$$\emptyset \neq S' \cap V \subset X_i \cap V = U = S \cap V,$$

which implies $S \cap S' \neq \emptyset$ and $S = S'$. This gives $U \subset S' \subset \overline{S'}$ as in the claim.

• If $i < j$, we have

$$\begin{aligned} \emptyset \neq S' \cap V \subset (X_j \backslash X_{j-1}) \cap V &= \varphi(U \times (L_{j-i-1} \backslash L_{j-i-2}) \times]0,1[) \\ &= \varphi(\bigsqcup_T U \times T \times]0,1[), \end{aligned}$$

where T runs over the connected components of $L_{j-i-1} \backslash L_{j-i-2}$. We may write

$$(X_j \backslash X_{j-1}) \cap V = \bigsqcup_{S''} \bigsqcup_{C''} C'',$$

where S'' runs over the strata of $X_j \backslash X_{j-1}$ and C'' over the connected components of $S'' \cap V$. By local connectivity of $X_j \backslash X_{j-1}$, the C'' are open in $(X_j \backslash X_{j-1}) \cap V$. As $S' \cap V \neq \emptyset$, there exists a non empty connected component C' of $S' \cap V$ and a non empty connected component, T, of $L_{j-i-1} \backslash L_{j-i-2}$ with

$$C' = \varphi(U \times T \times]0,1[).$$

Observe that $U \times \{\vartheta\}$ is in the closure of $U \times T \times]0,1[$ in $U \times \mathring{c}L$. This implies that U is included in the closure of C' in V and gives $U \subset \overline{S'}$.

As the claim is proved, we know that, for any $U \in \mathfrak{U}$, we have $U \subset \overline{S'}$ or $U \cap \overline{S'} = \emptyset$. The condition $S \cap \overline{S'} \neq \emptyset$ and the connectivity of S imply $U \subset \overline{S'}$, and therefore $S \subset \overline{S'}$. □

In [23], King defines a subcomplex of the singular chain complex of a filtered space as follows.

Definition 3.3. Let $(X, (X_i)_{0 \leq i \leq n})$ be a filtered space and let $\overline{p}$ be a loose perversity. A singular simplex, $\sigma \colon \Delta \to X$, is said *King $\overline{p}$-admissible* if $\sigma^{-1}(X_{n-j} \backslash X_{n-j-1})$ is contained in the $(\dim \Delta - j + \overline{p}(j))$-skeleton of Δ, for all $j \in \{0,1,\dots,n\}$. A chain c is King $\overline{p}$-admissible if there exist $\overline{p}$-admissible simplices, σ_j, so that $c=\sum_j \lambda_j\sigma_j$, $\lambda_j \in \mathbb{Z}$.

Let $K_*^{\overline{p}}(X)$ be the chain complex which consists of the singular chains c, such that c and its boundary ∂c are King $\overline{p}$-admissibles.

Proposition 3.4. *Let X be a locally conelike space and let $\overline{p}$ be a perversity. Then the canonical inclusion $C_*^{\overline{p}}(X) \to K_*^{\overline{p}}(X)$ induces an isomorphism in homology.*

Proof. Let $\sigma\colon \Delta \to X$ be a $\overline{p}$-admissible simplex. As the set $\sigma^{-1}X_{n-i}$ is a face of Δ of dimension less than, or equal to, $(\dim\Delta - i + \overline{p}(i))$, its subset $\sigma^{-1}(X_{n-i}\backslash X_{n-i-1})$ is also included in this skeleton. This shows that any σ in $C_*^{\overline{p}}(X)$ belongs also to $K_*^{\overline{p}}(X)$ and we have a canonical inclusion $C_*^{\overline{p}}(X) \subset K_*^{\overline{p}}(X)$

We have now only to check the five conditions of [23, Theorem 10]. Conditions (1), (3), (4) and (5) are direct consequences of Proposition 2.12 and Theorem B. The assertion (2) is a consequence of the Künneth formula established in Proposition 2.12, (iv). $\square$

Remark 3.5. If $\sigma\colon \Delta \to X$ is a simplex of a filtered space X of dimension n, we consider the two following conditions,

(a) $\sigma^{-1}X_{n-j}$ is included in the $(\dim\Delta - j + \overline{p}(j))$-skeleton of Δ, for all j,
(b) $\sigma^{-1}(X_{n-j}\backslash X_{n-j-1})$ is included in the $(\dim\Delta - j + \overline{p}(j))$-skeleton of Δ, for all j.

In the previous proof, we used that condition (a) implies condition (b). As already quoted by King ([23]), in the case of a perversity $\overline{p}$, condition (b) implies also condition (a). For proving that, we observe that $\sigma^{-1}X_{n-j}$ is a union of $\sigma^{-1}(X_{n-j-k}\backslash X_{n-j-k-1})$, $k \geq 0$. An induction based on

$$\begin{aligned}\dim\Delta-(i+1)+\overline{p}(i+1) &\leq \dim\Delta-(i+1)+\overline{p}(i)+1\\ &\leq \dim\Delta - i+\overline{p}(i),\end{aligned}$$

shows that

$$\dim\Delta-(j+k)+\overline{p}(j+k) \leq \dim\Delta - j+\overline{p}(j),$$

for any k, $k \geq 0$.

4. Filtered face sets

We begin with a brief recall on face sets introduced by Rourke and Sanderson and define our main objects, the filtered face sets. We present also the notions of filtered skeleton, link and expanded link which are crucial in the proofs of sections 5, 6 and 7.

For any integer n, we set $[n]=\{0,1,\dots,n\}$. Denote by $\boldsymbol{\Delta}$ the category whose morphisms are the *injective* order-preserving maps $f\colon [n] \to [m]$. A morphism of $\boldsymbol{\Delta}$ is a composition of face operators $\delta_i\colon [n] \to [n+1]$ defined by

$$\delta_i(j)=\left\{\begin{array}{cl} j & \text{if } j<i,\\ j+1 & \text{if } j\geq i,\end{array}\right.$$

which satisfy $\delta_j\delta_i=\delta_i\delta_{j-1}$, if $i<j$.

Definition 4.1. A *face set* is a contravariant functor, $\underline{K}$, from $\mathbf{\Delta}$ to the category of sets, $[n] \mapsto \underline{K}_n$. An element $\sigma \in \underline{K}_n$ is called *a simplex of dimension n,* that we denote $|\sigma| = n$. The *n-skeleton* of $\underline{K}$ is $K^{(n)} = \cup_{p\leq n}\underline{K}_p$. The images by $\underline{K}$ of face operators are maps $\partial_i \colon \underline{K}_{n+1} \to \underline{K}_n$ such that $\partial_i\partial_j = \partial_{j-1}\partial_i$, if $i < j$.

If $\underline{K}$ and $\underline{K}'$ are face sets, a *face map,* $\underline{f} \colon \underline{K} \to \underline{K}'$, is a natural transformation between the two functors $\underline{K}$ and $\underline{K}'$. We denote by $\mathbf{\Delta}$ – Sets the category of face maps.

For basic properties of face sets and their relation with the more classical simplicial setting, we refer to the foundational paper [30], observing that the homotopy theory of face sets is the same as the homotopy theory of simplicial sets.

Example 4.2. The face set $\underline{\Delta}^n$ is defined as follows: $\underline{\Delta}^n_p$ is the set of injective order-preserving maps from $[p]$ to $[n]$. It has only one simplex of dimension n, denoted by $[\Delta^n]$. If $\underline{K}$ is a face set, there is a bijection between the set of elements $\sigma \in \underline{K}_n$ and the set of face maps, $\underline{\sigma} \colon \underline{\Delta}^n \to \underline{K}$, defined by $\underline{\sigma}(f) = \underline{K}(f)(\sigma)$. In the sequel, we do not make any distinction between $\underline{\sigma} \colon \underline{\Delta}^n \to \underline{K}$ and $\sigma \in \underline{K}_n$.

Recall from [30], that the realization of a face set, $\underline{K}$, is the CW-complex, $\|\underline{K}\|$, defined by

$$\|\underline{K}\| = \bigcup_{n=0}^{\infty} (\underline{K}_n \times \Delta^n)/(\partial_i x, t) \sim (t, \partial_i t),$$

where

- Δ^n is the standard n-simplex of $\mathbb{R}^{n+1}$, whose vertices $v_0, \dots, v_n$ verify $v_i = (t_0, \dots, t_n)$, $t_j = 0$ if $i \neq j$ and $t_i = 1$,
- in barycentric coordinates, the map $\partial_i \colon \Delta^{n-1} \to \Delta^n$ is the linear application defined by $\partial_i(v_j) = v_{\delta_i(j)}$,

For instance, the realization of the face set $\underline{\Delta}$ described in Example 4.2 is the standard simplex Δ^n and, in the sequel, we identify $[n]$, $\underline{\Delta}^n$ and Δ^n. The CW-complex $\|\underline{K}\|$ has one n-cell for each $\sigma \in \underline{K}_n$, of characteristic map $\tilde{\sigma} \colon \Delta^n \to \|\underline{K}\|$ defined by $\tilde{\sigma}(t_0, \dots, t_n) = [\sigma, (t_0, \dots, t_n)]$, where $[-]$ denotes the equivalence classes defining the realization $\|\underline{K}\|$.

We adapt now these objects to the context of intersection theory. First, *we fix an integer n which corresponds to the dimension of filtered sets; that means, we fix the number of elements in the filtration to n,* see Remark 4.6 for the topological meaning of this process.

Let $\mathbf{\Delta}^{[n]}_{\mathcal{F}}$ be the category whose

- objects are the join $\Delta = \Delta^{j_0} * \Delta^{j_1} * \dots * \Delta^{j_n}$, where Δ^{j_i} is the simplex of dimension j_i, possibly empty, with the conventions $\Delta^{-1} = \emptyset$ and $\emptyset * X = X$,
- maps are the $\sigma \colon \Delta = \Delta^{j_0} * \Delta^{j_1} * \dots * \Delta^{j_n} \to \Delta' = \Delta^{k_0} * \Delta^{k_1} * \dots * \Delta^{k_n}$, of the shape $\sigma = *_{i=0}^{n} \sigma_i$, with $\sigma_i \colon \Delta^{j_i} \to \Delta^{k_i}$ an injective order-preserving map for each i, with $j_i \geq -1$.

The category $\mathbf{\Delta}^{[n],+}_{\mathcal{F}}$ is the full subcategory of $\mathbf{\Delta}^{[n]}_{\mathcal{F}}$ whose objects are the $\Delta^{j_0} * \Delta^{j_1} * \cdots * \Delta^{j_n}$ with $\Delta^{j_n} \neq \emptyset$, i.e., $j_n \geq 0$.

Definition 4.3. A *filtered face set* is a contravariant functor, $\underline{K}$, from the category $\mathbf{\Delta}^{[n]}_{\mathcal{F}}$ to the category of sets, i.e., $(j_0, \ldots, j_n) \mapsto \underline{K}_{(j_0,\ldots,j_n)}$. The restriction of the filtered face set, $\underline{K}$, to $\mathbf{\Delta}^{[n],+}_{\mathcal{F}}$ is denoted $\underline{K}_+$.

If $\underline{K}$ and $\underline{K}'$ are filtered face sets, a *filtered face map, $\underline{f}\colon \underline{K} \to \underline{K}'$*, is a natural transformation between the two functors $\underline{K}$ and $\underline{K}'$. We denote by $\mathbf{\Delta}^{[n]}_{\mathcal{F}}$*−Sets* the category of filtered face maps.

Face sets are filtered face sets whose simplices, $\sigma\colon \Delta^{j_0} * \cdots * \Delta^{j_n} \to \underline{K}$, are such that $j_i = -1$ for all $i < n$.

Definition 4.4. The *perverse degree* of a simplex, $\sigma\colon \Delta^{j_0} * \Delta^{j_1} * \ldots * \Delta^{j_n} \to \underline{K}$, of a filtered face set, $\underline{K}$, is the $(n+1)$-uple

$$\|\sigma\| = (\|\sigma\|_0, \ldots, \|\sigma\|_n),$$

where $\|\sigma\|_\ell = \dim(\Delta^{j_0} * \cdots * \Delta^{j_{n-\ell}})$ if $\Delta^{j_0} * \cdots * \Delta^{j_{n-\ell}} \neq \emptyset$ and $\|\sigma\|_\ell = -\infty$ otherwise.

Observe that *any filtered face map preserves the perverse degree.* The next example is the main motivation for the introduction of filtered face sets.

Example 4.5. If X is a filtered space, we define a filtered face set by

$$\mathrm{ISing}^{\mathcal{F}}_*(X)_{j_0,\ldots,j_n} = \{\sigma\colon \Delta^{j_0} * \cdots * \Delta^{j_n} \to X,\ \sigma \text{ filtered}\},$$

for any $(j_0, \ldots, j_n)$. Moreover, any stratum-preserving stratified map, $X \to Y$, induces a filtered face map, $\mathrm{ISing}^{\mathcal{F}}_*(X) \to \mathrm{ISing}^{\mathcal{F}}_*(Y)$, see Theorem A and Remark 2.4.

Remark 4.6. Imposing a common dimension, n, to filtered face sets, is not a restriction for the study of intersection homology. Observe, from the definition, that the admissibility of a simplex of a filtered space, X, depends only on the codimension in X. Therefore, as noticed by Friedman in [11], one can add a fixed integer to each index of the filtration of a space without modifying its intersection homology. A practical consequence is that we can always suppose that the spaces we are considering have the same dimension without loosing generality.

The restriction to the subcategory $\mathbf{\Delta}^{[n],+}_{\mathcal{F}}$ is due to the fact that, in the topological setting, we do not consider simplices entirely included in the singular part. One may observe also (see Examples 5.6 and 5.9) that the blow-up's of differential forms are not defined if $\Delta^{j_n} = \emptyset$.

We define now the filtered notions of boundary and skeleta which coincide with usual notions of boundary and skeleta in the case $n = 0$.

Definition 4.7. Let $\Delta = \Delta^{j_0} * \cdots * \Delta^{j_n} \in \mathbf{\Delta}^{[n]}_{\mathcal{F}}$ be a filtered simplex, not reduced to a point. The *filtered boundary* of Δ is the filtered face set $\partial^{\mathcal{F}}\Delta$ defined by

$$\partial^{\mathcal{F}}\Delta = \Delta^{j_0} * \cdots * \Delta^{j_{i-1}} * \partial\Delta^{j_i} * \Delta^{j_{i+1}} * \cdots \Delta^{j_{n-1}} * \Delta^{j_n},$$

if $j_k = -1$ for $k < i$ and $j_i \geq 0$. If $\sigma\colon \Delta \to \underline{K}$ is a simplex of a filtered face set $\underline{K}$, we denote by $\partial^{\mathcal{F}}\sigma$ its restriction to the filtered boundary,

$$\partial^{\mathcal{F}}\sigma\colon \partial^{\mathcal{F}}\Delta \longrightarrow \Delta \xrightarrow{\sigma} \underline{K}$$

Definition 4.8. The *depth of the simplex* $\sigma\colon \Delta^{j_0} * \cdots * \Delta^{j_n} \to \underline{K}$ is the integer $v(\sigma) \in \{0, \ldots, n\}$ defined by

$$v(\sigma) = \max\{n - a \mid \Delta^{j_a} \neq \emptyset\}.$$

The depth $v(\underline{K})$ of a filtered face set, $\underline{K}$, is the maximum of the depth of its simplices.

Definition 4.9. Let $r \geq 0$ and $k \geq 0$. The *filtered skeleta* of a filtered face set, $\underline{K}$, are the filtered face sets, $\underline{K}^{[r],k}$, defined by

$$\underline{K}^{[r],k} = \{\sigma \in \underline{K} \mid v(\sigma) < r \text{ or } (v(\sigma) = r \text{ and } j_{v(\sigma)} \leq k)\}.$$

We set $\underline{K}^{[r]} = \cup_{k \geq 0} \underline{K}^{[r],k}$ and, by convention, $\underline{K}^{[r],-1} = \underline{K}^{[r-1]}$, if $r > 0$. The *regular part* of $\underline{K}$ is the face set $\underline{K}^{[0]}$.

Let $\underline{K}$ be a filtered face set. For any $\sigma\colon \Delta = \Delta^{j_0} * \cdots * \Delta^{j_n} \to \underline{K}^{[r],k}$, we denote by

- $R_1(\sigma)$ the restriction of σ to $\Delta^{j_{n-r}}$,
- $R_2(\sigma)$ the restriction of σ to $\Delta^{j_{n-r+1}} * \cdots * \Delta^{j_n}$.

We denote by $\mathcal{J}(\underline{K}^{[r],k})$ the set of $R_1(\sigma)$, when $\sigma \in \underline{K}^{[r],k} \backslash \underline{K}^{[r],k-1}$ and observe that the domain of $R_2(\sigma)$ can be empty. We set $\xi \lhd \sigma$ if ξ is a face (of any codimension) of a simplex σ.

Definition 4.10. Let $\underline{K}$ be a filtered face set and $\tau \in \mathcal{J}(\underline{K}^{[r],k})$.

(a) *The star of τ in $\underline{K}$* is the filtered face subset, $\underline{K}(\tau)$, of $\underline{K}$ defined by

$$\underline{K}(\tau) = \{\xi \in \underline{K} \mid \exists \sigma \in \underline{K}^{[r],k} \text{ with } R_1(\sigma) = \tau \text{ and } \xi \lhd \sigma\}.$$

(b) *The link of τ in $\underline{K}$* is the filtered face subset, $\mathcal{L}(\underline{K}, \tau)$, of $\underline{K}$ defined by

$$\mathcal{L}(\underline{K}, \tau) = \{\xi \in \underline{K} \mid \exists \sigma \in \underline{K}(\tau) \text{ with } \xi \lhd R_2(\sigma)\}.$$

Remark 4.11. Directly from the definition, we note that the filtered face set $\underline{K}(\tau)$ is never empty and that the link $\mathcal{L}(\underline{K}, \tau)$ can be empty. Observe also that R_2 can be written as a composite of face maps. More precisely, denote by $\partial_0\colon \Delta^{j_{n-r}-1} * \cdots * \Delta^{j_n} \to \Delta^{j_{n-r}} * \cdots * \Delta^{j_n}$ the first face operator and by $\partial_0^{1+j_{n-r}}\colon \Delta^{j_{n-r+1}} * \cdots * \Delta^{j_n} \to \Delta^{j_{n-r}} * \cdots * \Delta^{j_n}$ the iteration of ∂_0. By definition, we have $R_2 = \partial_0^{1+j_{n-r}}$.

Any face operator ∂_i defined on $\Delta^{j_{n-r+1}} * \cdots * \Delta^{j_n}$ induces a face operator on $\Delta^{j_{n-r}} * \Delta^{j_{n-r+1}} * \cdots * \Delta^{j_n}$ denoted $\partial_{i+1+j_{n-r}}$. With the usual rule of commutations of face operators, we have $\partial_i R_2 = R_2 \partial_{i+1+j_{n-r}}$.

We observe that

$$\underline{K}^{[r],k} = \underline{K}^{[r],k-1} \cup_{\tau \in \mathcal{J}(\underline{K}^{[r],k})} \underline{K}(\tau). \tag{2}$$

As $\underline{K}(\tau)$ is a filtered face set, the filtered face set $\underline{K}(\tau)(\partial_i\tau)$ is already defined for any face operator $\partial_i\colon \Delta^{k-1} \to \Delta^k$, i.e.,

$$\underline{K}(\tau)(\partial_i\tau) = \{\xi \in \underline{K}(\tau) \mid \exists \sigma \in \underline{K}(\tau) \text{ with } \xi \lhd \sigma \text{ and } R_1(\sigma) = \partial_i\tau\}.$$

A simplex σ as above is the face of a simplex $\gamma \in \underline{K}$ such that $R_1(\gamma) = \tau$ and we may also write

$$\underline{K}(\tau)(\partial_i\tau) = \{\xi \in \underline{K} \mid \exists\gamma \in \underline{K} \text{ with } \xi \lhd \partial_i\gamma \text{ and } R_1(\gamma) = \tau\}.$$

Definition 4.12. Let $\underline{K}$ be a filtered face set, $r \geq 0$, $k \geq 1$ and $\tau \in \mathfrak{J}(\underline{K}^{[r],k})$. We define a filtered face set $\underline{K}(\tau, \partial\tau)$ by

$$\underline{K}(\tau, \partial\tau) = \cup_i \underline{K}(\tau)(\partial_i\tau).$$

Observe that

$$\underline{K}(\tau) = \underline{K}(\tau)^{[r],k} \text{ and } \underline{K}(\tau, \partial\tau) = \underline{K}(\tau, \partial\tau)^{[r],k-1}.$$

With the determination of $\underline{K}(\tau)(\partial_i\tau)$ done above, we can characterise the elements of $\underline{K}(\tau, \partial\tau)$ as the set of $\xi \in \underline{K}(\tau)$ such that ξ is a face of $\sigma \in \underline{K}(\tau)$ with $R_1(\sigma) = \partial_i\tau$ for some face operator ∂_i.

If $j_{n-r} = 0$, we denote $\underline{K}(\tau)$ by $\underline{K}(\{\vartheta\})$ and $\mathcal{L}(\underline{K}, \tau)$ by $\mathcal{L}(\underline{K}, \{\vartheta\})$. We determine now the intersection of the two filtered face sets in the right hand side of equation (2).

Proposition 4.13. *If $\underline{K}$ is a filtered face set, $r \geq 1$, $k \geq 0$ and $\tau \in \mathfrak{J}(\underline{K}^{[r],k})$, we have*

$$\underline{K}^{[r],k-1} \cap \underline{K}(\tau) = \begin{cases} \underline{K}(\tau, \partial\tau), & \textit{if } k \geq 1, \\ \mathcal{L}(\underline{K}, \tau), & \textit{if } k = 0. \end{cases}$$

Proof. Suppose first $k \geq 1$. An element $\xi \in \underline{K}^{[r],k-1} \cap \underline{K}(\tau)$ is a face of a $\sigma \in \underline{K}(\tau)$ with $R_1(\sigma) = \tau$ and we have two possibilities:

- $v(\xi) = r$ and $j_{n-r} \leq k - 1$. In this case, $R_1(\xi)$ is a proper face of τ. Let $\tau' = \partial_i\tau$ such that $R_1(\xi) \subset \partial_i\tau$ and set $\sigma' = \partial_i\sigma$. Then we have $R_1(\sigma') = \tau'$, $\sigma' \in \underline{K}(\tau)$ and ξ is a face of σ'. This implies $\xi \in \underline{K}(\tau, \partial\tau)$.
- or $v(\xi) \leq r - 1$. In this case, ξ is a face of $R_2(\sigma)$. Set $\tau' = \partial_i\tau$ and $\sigma' = \partial_i\sigma$ for a face operator ∂_i. Then, we have $\sigma' \in \underline{K}(\tau)$, $R_1(\sigma') = \tau'$ and $R_2(\sigma)$ is a face of σ'. Thus ξ is also a face of σ' and we conclude $\xi \in \underline{K}(\tau, \partial\tau)$.

The reverse inclusion comes directly from $\underline{K}(\tau, \partial\tau) = \underline{K}(\tau, \partial\tau)^{[r],k-1}$. The case $k = 0$ is proved in a similar way, by using $\mathcal{L}(\underline{K}, \tau) = \mathcal{L}(\underline{K}, \tau)^{[r-1]}$. □

Definition 4.14. Let $\underline{K}$ be a filtered face set, $r \geq 1$, $k \geq 0$ and $\tau \in \mathfrak{J}(\underline{K}^{[r],k})$. The *expanded link* of τ in $\underline{K}$ is the filtered face set $\mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)$ defined by

$$\mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau) = \{(R_2\sigma, \sigma) \mid \sigma \in \underline{K}, R_1(\sigma) = \tau \text{ and } \sigma \neq \tau\},$$

with face maps,

$$\partial_i(R_2(\sigma), \sigma) = (\partial_i R_2(\sigma), \partial_{i+1+k}\sigma) = (R_2(\partial_{i+1+k}\sigma), \partial_{i+1+k}\sigma),$$

for all $i \in \{0, \ldots, \dim R_2(\sigma)\}$, and perverse degrees,

$$\|(R_2(\sigma), \sigma)\|_\ell = \|R_2(\sigma)\|_\ell,$$

for all $\ell \in \{0, \ldots, n\}$.

By definition, a simplex $(\alpha,\sigma)\in\mathcal{L}^{\tt exp}(\underline{K},\tau)$ if, and only if, $R_1(\sigma)=\tau$ and $R_2(\sigma)=\alpha$. The difference between the link and the expanded link lies in the fact that two simplices σ and σ' of $\underline{K}(\tau)$, verifying $R_2(\sigma)=R_2(\sigma')$, give the same element in $\mathcal{L}(\underline{K},\tau)$ and two distinct elements in $\mathcal{L}^{\tt exp}(\underline{K},\tau)$. In the expanded link, we keep trace of σ and σ' as parameters. Also, the condition $\sigma\neq\tau$ means $R_2(\sigma)\neq\emptyset$.

Definition 4.15. A *morphism of pairs of filtered face sets,* $f\colon(\underline{K}_1,\underline{L}_1)\to(\underline{K}_2,\underline{L}_2)$, is a filtered face map, $f\colon\underline{K}_1\to\underline{K}_2$, such that $f(\underline{L}_1)\subset\underline{L}_2$. A *relative isomorphism,* $f\colon(\underline{K}_1,\underline{L}_1)\to(\underline{K}_2,\underline{L}_2)$, is a morphism of pairs of filtered face sets, such that $f\colon\underline{K}_1\to\underline{K}_2$ is surjective and its restriction to the complementary subsets is a bijection respecting the perverse degree, $f\colon\underline{K}_1\backslash\underline{L}_1\cong\underline{K}_2\backslash\underline{L}_2$.

Example 4.16. From Proposition 4.13, we deduce that the map

$$(\cup_{\tau_i\in\mathfrak{J}(\underline{K}^{[r],k})}\underline{K}(\tau_i),\cup_{\tau_i\in\mathfrak{J}(\underline{K}^{[r],k})}\underline{K}(\tau_i,\partial\tau_i))\to(\underline{K}^{[r],k},\underline{K}^{[r],k-1})$$

is a relative isomorphism.

Example 4.17. Let $\underline{K}$ be a filtered face set of depth $v(\underline{K})=\ell-1$ with $\ell\in\{1,\dots,n\}$. Let $k\in\mathbb{N}$. We denote by $\Delta^k*\underline{K}$ the filtered face set, of depth ℓ, generated by the simplices

$$\Delta^k*\Delta^j\xrightarrow{\mathrm{id}*\sigma}\Delta^k*\underline{K},$$

with $\sigma\in\underline{K}$ and $k=j_{n-\ell}$. The simplices of this filtered face set $\Delta^k*\underline{K}$ are of three types, $\partial_I\Delta^k$ with $I\subsetneq\{0,\dots,k\}$, $\sigma\in\underline{K}$ or $\partial_I\Delta^k*\sigma$. Their perverse degrees are defined by

- $\|\partial_I\Delta^k\|_{\ell'}=\left\{\begin{array}{cl}-\infty & \text{if } \ell<\ell',\\ k-|I| & \text{if } \ell\geq\ell',\end{array}\right.$
- $\|\sigma\|_{\ell'}=\left\{\begin{array}{cl}-\infty & \text{if } \ell\leq\ell',\\ \|\sigma\|_{\ell'} & \text{if } \ell>\ell',\end{array}\right.$
- $\|\partial_I\Delta^k{*}\sigma\|_{\ell'}=\left\{\begin{array}{cl}-\infty & \text{if } \ell<\ell',\\ k-|I| & \text{if } \ell'=\ell,\\ & \text{or } (\ell>\ell' \text{ and } \|\sigma\|_{\ell'}=-\infty),\\ k-|I|+\|\sigma\|_{\ell'}+1 & \text{if } \ell>\ell' \text{ and } \|\sigma\|_{\ell'}\neq-\infty.\end{array}\right.$

If $\underline{L}$ is a subcomplex of Δ^k, we define similarly a filtered face set $\underline{L}*\underline{K}$, called *the join of* $\underline{L}$ *and* $\underline{K}$*,* as, for instance, $\partial\Delta^k*\underline{K}$. There exists certainly a notion of join of two filtered face sets (see [13]) but we do not need it.

Proposition 4.18. *Let* $\underline{K}$ *be a filtered face set,* $r\geq 1$*,* $k\geq 0$ *and* $\tau\in\mathfrak{J}(\underline{K}^{[r],k})$*. There exists a relative isomorphism*

$$f\colon(\Delta^k*\mathcal{L}^{\tt exp}(\underline{K},\tau),\partial\Delta^k*\mathcal{L}^{\tt exp}(\underline{K},\tau))\to(\underline{K}(\tau),\underline{K}(\tau,\partial\tau)),$$

preserving the perverse degree.

Proof. Let $(\alpha,\sigma)\in\mathcal{L}^{\tt exp}(\underline{K},\tau)$ and $I\subset\{0,\dots,k\}$. The application f is defined by

$$f(\partial_I\Delta^k)=\partial_I\tau,\ f(\alpha,\sigma)=\sigma \text{ and } f(\partial_I\Delta^k*(\alpha,\sigma))=\partial_I\sigma.$$

By construction, we have $f(\partial\Delta^k * \mathcal{L}^{\mathtt{exp}}(\underline{K},\tau)) = \underline{K}(\tau,\partial\tau)$. Denote $\ell = n-r$. We check first that f decreases the perverse degree,

$$\begin{aligned}
\|f(\partial_I\Delta^k * (\alpha,\sigma))\|_{\ell'} &= \|\partial_I\sigma\|_{\ell'} = -\infty = \|\partial_I\Delta^k * (\alpha,\sigma)\|_{\ell'}, \text{ if } \ell < \ell',\\
\|f(\partial_I\Delta^k * (\alpha,\sigma))\|_{\ell'} &= \|\partial_I\sigma\|_{\ell'} = \|\partial_I\Delta^k * (\alpha,\sigma)\|_{\ell'} = k - |I|,\\
&\qquad \text{if } \ell' = \ell \text{ or } (\ell > \ell' \text{ and } \|\sigma\|_{\ell'} = -\infty),\\
\|f(\partial_I\Delta^k * (\alpha,\sigma))\|_{\ell'} &= \|\partial_I\sigma\|_{\ell'} = \|\partial_I\Delta^k * (\alpha,\sigma)\|_{\ell'} = k - |I| + \|\sigma\|_{\ell'} + 1,\\
&\qquad \text{if } \ell > \ell' \text{ and } \|\sigma\|_{\ell'} \neq -\infty.
\end{aligned}$$

A similar verification is done for the other cases. We leave also to the reader the verifications of the compatibility with face maps, i.e., $f \circ \partial_i = \partial_i \circ f$, for all i.

For proving the surjectivity, we consider $\beta \in \underline{K}(\tau)$. By definition, the simplex β is a face of a simplex $\sigma \in \underline{K}(\tau)$ such that $R_1(\sigma) = \tau$. That means, there exist $I \subset \{0,\dots,k\}$ and $J \subset \{k+1,\dots\}$ with $\partial_I\partial_J\sigma = \beta$. They verify $R_1(\partial_J\sigma) = \tau$ and $R_2(\beta) = R_2(\partial_I\partial_J\sigma) = R_2(\partial_J\sigma)$. We distinguish two cases:

- if $R_2(\partial_J\sigma) = \emptyset$, then $\partial_J\sigma = \tau$ and $\beta = \partial_I\tau = f(\partial_I\Delta^k)$,
- if $R_2(\partial_J\sigma) \neq \emptyset$, then $(R_2(\partial_J\sigma),\partial_J\sigma) \in \mathcal{L}^{\mathtt{exp}}(\underline{K},\tau)$ and $\beta = f(\partial_I\Delta^k * (R_2(\partial_J\sigma),\partial_J\sigma))$.

We study now the restriction to the complementary subsets. Let $\gamma \in \underline{K}(\tau)\backslash\underline{K}(\tau,\partial\tau)$. By definition, we have $R_1(\gamma) = \tau$. If $R_2(\gamma) = \emptyset$, then $\gamma = \tau = f(\Delta^k)$. If $R_2(\gamma) \neq \emptyset$, then $\gamma = f(\Delta^k * (R_2(\gamma),\gamma))$. This gives the surjectivity and the injectivity is obvious from inspection of the definition of f in this case. ☐

Remark 4.19. The join $\Delta^k * \mathcal{L}^{\mathtt{exp}}(\underline{K},\tau)$ on the expanded link (see Example 4.17) can also be defined functorially.

Let $\mathbf{\Delta}_{\mathcal{F}}^{[n,\ell],k}$ be the full subcategory of $\mathbf{\Delta}_{\mathcal{F}}^{[n]}$ whose objects are the joins $\Delta^{j_0} * \cdots * \Delta^{j_n}$ with $j_i = -1$ if $i < \ell$ and $j_\ell \geq k$. We denote by $\mathbf{\Delta}_{\mathcal{F}}^{[n,\ell],k}$*–Sets* the category of functors from $\mathbf{\Delta}_{\mathcal{F}}^{[n,\ell],k}$ to the category of sets. As this last one is complete and cocomplete, the forgetful functor $\mathcal{O}\colon \mathbf{\Delta}_{\mathcal{F}}^{[n]}\text{–}Sets \to \mathbf{\Delta}_{\mathcal{F}}^{[n,\ell],k}\text{–}Sets$ has a left adjoint $\mathcal{F}$.

Let $\tau \in \mathcal{J}(\underline{K}^{[r],k})$ and $\sigma\colon \Delta^{j_\ell} * \cdots * \Delta^{j_n} \to \underline{K}(\tau)$ be a generic simplex of $\underline{K}(\tau)$. In $\mathcal{O}(\underline{K}(\tau))$, only the simplices such that $j_\ell = k$ are kept and the face operators $(\partial_0)^i$ with $i \leq k+1$ do not appear. In $\mathcal{F}(\mathcal{O}(\underline{K}(\tau)))$, we must add the operators missing and a simplex of $\mathcal{F}(\mathcal{O}(\underline{K}(\tau)))$ is of the shape $\beta * R_2(\sigma)$ with $\beta = \partial_0^j(\tau)$, for some $j \leq k+1$. Theses faces being added freely, they are not submitted to any identification and $\mathcal{F}(\mathcal{O}(\underline{K}(\tau)))$ is of the shape $\Delta^k * \underline{L}$ for some filtered face set, $\underline{L}$. An inspection of the definition of the expanded link gives

$$\mathcal{F}(\mathcal{O}(\underline{K}(\tau))) = \Delta^k * \mathcal{L}^{\mathtt{exp}}(\underline{K},\tau).$$

5. LOCAL SYSTEM ON FILTERED FACE SETS

We define perverse local systems of coefficients, $\mathcal{M}$, over a filtered face set $\underline{K}$ and the intersection cohomology of $\underline{K}$ with coefficients in $\mathcal{M}$, for a loose perversity $\overline{p}$. The blow-up $\widetilde{F}$ of certain local systems over face sets, F, are examples of this

> situation. We detail the particular cases of cochains (over $\mathbb{Z}$) and Sullivan forms (over $\mathbb{Q}$). The main result (Theorem C) gives sufficient conditions on the local systems, F and G, for having an isomorphism between the intersection cohomologies of a filtered face set, $\underline{K}$, with coefficients in the blow-up's $\widetilde{F}$ and $\widetilde{G}$.

In this section, all cochain complexes are over a commutative ring R. The particular case $R = \mathbb{Q}$ is explicitly quoted if necessary.

Homology and cohomology theories on a face set, $\underline{K}$, can be expressed with local systems, see [21] for instance in the simplicial case. We adapt these local systems to the filtered situation as follows.

Definition 5.1. A *strict perverse cochain complex* is a cochain complex in which each element, c, has a perverse degree,

$$\|c\| = (\|c\|_1, \ldots, \|c\|_n),$$

with $\|c\|_i \in \mathbb{N} \cup \{-\infty\}$, and such that $\|c + c'\| \leq \max(\|c\|, \|c'\|)$. A *strict perverse* DGA is a perverse cochain complex and a DGA such that $\|c \cdot c'\| \leq \|c\| + \|c'\|$.

Definition 5.2. A *perverse local system of cochains over a filtered face set*, $\underline{K}$, is a family of strict perverse cochains, $\mathcal{M}_\sigma$, indexed by the simplices σ of $\underline{K}$, and a family of cochain maps, $\partial_i \colon \mathcal{M}_\sigma \to \mathcal{M}_{\partial_i \sigma}$, decreasing the perverse degree and such that $\partial_i \partial_j = \partial_{j-1} \partial_i$, if $i < j$ and $\partial_i \partial_j \in \mathbf{\Delta}_{\mathcal{F}}^{[n],+}$.

The *space of global sections* of $\mathcal{M}$ over $\underline{K}$ is the cochain complex $\mathcal{M}(\underline{K})$ defined as follows: an element $\omega \in \mathcal{M}^j(\underline{K})$ is a function which assigns to each simplex $\sigma \in \underline{K}$ an element $\omega_\sigma \in \mathcal{M}_\sigma^j$ such that $\omega_{\partial_i \sigma} = \partial_i(\omega_\sigma)$ for all σ and all face operators $\partial_i \in \mathbf{\Delta}_{\mathcal{F}}^{[n],+}$. The *perverse degree of* $\omega \in \mathcal{M}(\underline{K})$ is the supremum of the perverse degrees of the ω_σ, $\sigma \in \underline{K}$.

The adaptation of Definition 5.2 to strict perverse DGA's is straightforward and if $\mathcal{M}$ is a local system of strict perverse DGA's, the space of global sections is a DGA. The laws and differential on $\mathcal{M}(\underline{K})$ are defined by $(\lambda\omega + \mu\omega')_\sigma = \lambda\omega_\sigma + \mu\omega'_\sigma$, $(d\omega)_\sigma = d\omega_\sigma$ and $(\omega\omega')_\sigma = \omega_\sigma\omega'_\sigma$ in the case of DGA's.

Let $\mathcal{M}$ be a perverse local system on a filtered face set $\underline{K}$. Observe that any filtered face map, $f \colon \underline{L} \to \underline{K}$, induces a perverse local system on $\underline{L}$, defined by

$$(f^*\mathcal{M})_\sigma = \mathcal{M}_{f \circ \sigma}.$$

Therefore, a perverse local system, $\mathcal{M}$, on $\underline{K}$ induces a perverse local system on any filtered face subset of $\underline{K}$; we still denote by $\mathcal{M}$ these induced systems.

Definition 5.3. Let $\overline{q}$ be a loose perversity and $\mathcal{M}$ be a perverse local system over a filtered face set $\underline{K}$. A global section $\omega \in \mathcal{M}(\underline{K})$ is $\overline{q}$-*admissible* if $\|\omega\|_i \leq \overline{q}(i)$, for any $i \in \{1, \ldots, n\}$. A global section ω is of *intersection for* $\overline{q}$ if ω and $d\omega$ are $\overline{q}$-admissible.

We denote by $\mathcal{M}_{\overline{q}}(\underline{K})$ the complex of global sections of intersection for $\overline{q}$ and by $H^*_{\overline{q}}(\underline{K}; \mathcal{M})$ its homology, called the *intersection cohomology of* $\underline{K}$ *with coefficients in* $\mathcal{M}$, *for the loose perversity* $\overline{q}$.

In the case $n = 0$, a perverse local system is called a local system. The PL-forms of Sullivan and the face cochains are examples of local systems. We recall their construction.

Example 5.4 (Universal system of cochains)**.** A *universal system of cochains* is a functor F from the category $\mathbf{\Delta}$ to the category of cochains. Such functor defines a local system of cochains on any face set, $\underline{K}$, by setting $F_\sigma = F(\Delta^{|\sigma|})$ and $\partial_i = F(\delta_i)\colon F_\sigma \to F_{\partial_i\sigma}$. The same process works for the definition of universal systems of DGA's.

Example 5.5 (Face cochain algebra)**.** By setting $C^r(n) = C^r(\underline{\Delta}^n; R)$, we define a universal system of DGA's. If $\underline{K}$ is a face set, the associated DGA of global sections, $C^*(\underline{K}; R)$, is defined by

- $C^j(\underline{K}; R)$, is the free R-module of all set maps $f\colon \underline{K}_j \to R$,
- the differential δ is given by
$$(\delta f)(\sigma) = \sum_{i=0}^{j+1}(-1)^i f(\partial_i\sigma),\ f \in C^j(\underline{K}; R),\ \sigma \in \underline{K}_{j+1},$$
- the product of $f \in C^j(\underline{K}; R)$ and $g \in C^k(\underline{K}; R)$ is defined by
$$(f.g)(\sigma) = f(\partial_j^F\sigma)g(\partial_k^B\sigma),\ \sigma \in \underline{K}_{j+k},$$
where $\partial_j^F\colon \underline{K}_{j+k} \to \underline{K}_j$ and $\partial_k^B\colon \underline{K}_{j+k} \to \underline{K}_k$ are given by $\partial_j^F\sigma = (\partial_{j+1} \circ \cdots \partial_{j+k})\sigma$ and $\partial_k^B\sigma = (\partial_0 \circ \cdots \circ \partial_0)\sigma$.

Example 5.6 (Sullivan's polynomial forms)**.** Let $R = \mathbb{Q}$. We define a universal system of CDGA's by setting

$$A_{PL}(n) = \wedge(t_0, \dots, t_n, dt_0, \dots, dt_n)/(t_0 + \cdots t_n = 1,\ dt_0 + \cdots dt_n = 0),$$

with $|t_i| = 0$ and $|dt_i| = 1$. Geometrically, the elements of $A_{PL}(n)$ are the polynomial differential forms, with rational coefficients, on the simplex Δ^n. The associated CDGA of global sections is denoted $A_{PL}(\underline{K})$, if $\underline{K}$ is a face set.

The systems we are using in the sequel of this work come from a universal one but there exist important systems which are not of this particular type. For instance, if $\underline{f}\colon \underline{E} \to \underline{K}$ is a Kan fibration between face sets, for any $\sigma\colon \Delta \to \underline{K}$, we denote by $\underline{E}_\sigma$ the pullback of $\underline{f}$ and σ. The association $\sigma \mapsto A_{PL}(\underline{E}_\sigma)$ is a local system over $\underline{K}$ which is the key tool in [21] and [15].

We introduce now the fundamental notion of *blow-up of a universal system*, based on the blow-up of a simplex which first appears in [31], see also [32].

Definition 5.7. The *blow-up of a filtered simplex,* $\Delta = \Delta^{j_0} * \cdots * \Delta^{j_n}$, is the map,

$$\mu\colon \widetilde{\Delta} = c\Delta^{j_0} \times \cdots \times c\Delta^{j_{n-1}} \times \Delta^{j_n} \to \Delta = \Delta^{j_0} * \cdots * \Delta^{j_n},$$

defined by

$$\begin{aligned}\mu([y_0, s_0], \dots, [y_{n-1}, s_{n-1}], y_n) \quad = \quad & s_0y_0 + (1 - s_0)s_1y_1 + \cdots \\ & +(1 - s_0) \cdots (1 - s_{n-2})s_{n-1}y_{n-1} \\ & +(1 - s_0) \cdots (1 - s_{n-2})(1 - s_{n-1})y_n,\end{aligned}$$

where $y_i \in \Delta^{j_i}$ and $[y_i, s_i] \in c\Delta^{j_i}$.

The prism $\widetilde{\Delta}$ is sometimes also called the blow-up of $\Delta^{j_0} * \cdots * \Delta^{j_n}$. The map μ *induces a diffeomorphism* between the interior of the blow-up $\widetilde{\Delta}$ and the interior of the simplex Δ. In the previous representation of the elements of $\widetilde{\Delta}$, the face $\Delta^{j_i} \times \{1\}$ of the simplex $c\Delta^{j_i}$ corresponds to $s_i = 1$.

Remark 5.8. The blow-up μ can also be expressed in terms of barycentric coordinates by

$$\mu((x_0, t_0), \ldots, (x_{n-1}, t_{n-1}), x_n) = (x_0, t_0 x_1, t_0 t_1 x_2 \ldots, t_0 \cdots t_{n-1} x_n),$$

where $x_i \in \mathbb{R}^{j_i+1}$ and $t_i \in \mathbb{R}$. In this context, the face $\Delta^{j_i} \times \{1\}$ of the simplex $c\Delta^{j_i}$ corresponds to $t_i = 0$.

Example 5.9. As already observed in [31] in the case of differential forms, any universal system of cochains, F, from the category $\mathbf{\Delta}$ to the category of cochains, gives rise to a perverse local system of cochains, $\widetilde{F}$, on any filtered face set, $\underline{K}$, defined on $\sigma\colon \Delta^{j_0} * \cdots * \Delta^{j_n} \to \underline{K}$ by

$$\widetilde{F}_\sigma = F(c\Delta^{j_0}) \otimes \cdots \otimes F(c\Delta^{j_{n-1}}) \otimes F(\Delta^{j_n}),$$

with the obvious face maps. The *perverse degree* of a global section $\omega \in \widetilde{F}(\underline{K})$ is defined with a method similar to a construction of Brylinski ([5]).

Let $\ell \in \{1, \ldots, n\}$ such that $\Delta^{j_{n-\ell}} \neq \emptyset$. We first define the *$\ell$-perverse degree,* $\|\omega_\sigma\|_\ell$, of an element $\omega_\sigma \in F(c\Delta^{j_0}) \otimes \cdots \otimes F(c\Delta^{j_{n-1}}) \otimes F(\Delta^{j_n})$, as follows. The restriction

$$\omega_{\sigma,n-\ell} \in F(c\Delta^{j_0}) \otimes \cdots \otimes F(\Delta^{j_{n-\ell}} \times \{1\}) \otimes \cdots \otimes F(c\Delta^{j_{n-1}}) \otimes F(\Delta^{j_n})$$

of ω_σ can be written as $\omega_{\sigma,n-\ell} = \sum_k \omega'_{\sigma,n-\ell}(k) \otimes \omega''_{\sigma,n-\ell}(k)$, with

- $\omega'_{\sigma,n-\ell}(k) \in F(c\Delta^{j_0}) \otimes \cdots \otimes F(c\Delta^{j_{n-\ell-1}}) \otimes F(\Delta^{j_{n-\ell}} \times \{1\})$ and
- $\omega''_{\sigma,n-\ell}(k) \in F(c\Delta^{j_{n-\ell+1}}) \otimes \cdots \otimes F(\Delta^{j_n})$.

If $\omega_{\sigma,n-\ell} \neq 0$, we set $\|\omega_\sigma\|_\ell = \sup_k \left\{ |\omega''_{\sigma,n-\ell}(k)| \text{ such that } \omega'_{\sigma,n-\ell}(k) \neq 0 \right\}$. If $\omega_{\sigma,n-\ell} = 0$ or $\Delta^{j_{n-\ell}} = \emptyset$, we set $\|\omega_\sigma\|_\ell = -\infty$.

The *perverse degree* of a global section $\omega \in \widetilde{F}(\underline{K})$ is the n-uple

$$\|\omega\| = (\|\omega\|_1, \ldots, \|\omega\|_n),$$

where $\|\omega\|_\ell$ is the supremum of the $\|\omega_\sigma\|_\ell$ for all $\sigma \in \underline{K}$.

Finally, observe that, in the case of differential forms, the ℓ-perverse degree of ω_σ is the degree along the fibers of $\omega_{\sigma,n-\ell}$ relatively to the projection (see [20, Page 122])

$$\mathrm{pr}\colon c\Delta^{j_0} \times \cdots \times \left(\Delta^{j_{n-\ell}} \times \{1\}\right) \times \cdots \times c\Delta^{j_{n-1}} \times \Delta^{j_n} \to c\Delta^{j_0} \times \cdots \times \left(\Delta^{j_{n-\ell}} \times \{1\}\right).$$

Definition 5.10. Let F be a universal system defined on $\mathbf{\Delta}$ and $\underline{K}$ be a filtered face set. The perverse local system, $\widetilde{F}$, described above, is called the *blow-up* of F over $\underline{K}$.

Let $\underline{K}$ be a filtered face set. To any such $\widetilde{F}$ and any loose perversity $\overline{q}$, we associate the complex $\widetilde{F}_{\overline{q}}(\underline{K})$ and its cohomology $H^*_{\overline{q}}(\underline{K}, \widetilde{F})$ as in Definition 5.3. Observe that the value $\overline{q}(0)$ does not matter because there is no perverse condition on the regular part.

Example 5.9 and Definition 5.10 are easily adaptated to universal systems of DGA's. The algebra of Sullivan's polynomial forms and the face cochain algebra of Examples 5.6 and 5.5 give the main examples of blow-up's used in this work, denoted respectively by $\widetilde{A}_{PL}$ and $\widetilde{C}^*$, the cochain complexes of intersection being denoted by $\widetilde{A}_{PL,\overline{q}}$ and $\widetilde{C}^*_{\overline{q}}$. The differential forms corresponding to $\overline{q} = \overline{0}$ are the forms introduced by Verona in [37].

Proposition 5.11. *Let F be a universal system defined on $\mathbf{\Delta}$. Any filtered face map $f\colon \underline{K} \to \underline{K}'$ induces a cochain map (or* DGA*'s map), $f^*\colon \widetilde{F}(\underline{K}') \to \widetilde{F}(\underline{K})$, defined by $(f^*\omega)_\sigma = \omega_{f(\sigma)}$.*

Moreover, the map f^* decreases the perverse degree and induces a cochain map $f^*\colon \widetilde{F}_{\overline{q}}(\underline{K}') \to \widetilde{F}_{\overline{q}}(\underline{K})$, for any loose perversity $\overline{q}$.

Proof. Let $\omega \in \widetilde{F}(\underline{K}')$ and $\sigma \in \underline{K}$. Any map f which is compatible with face operators satisfies

$$(f^*\omega)_{\partial_i\sigma} = \omega_{f(\partial_i\sigma)} = \omega_{\partial_i f(\sigma)} = \partial_i^*\omega_{f(\sigma)} = \partial_i^*(f^*(\omega_\sigma)).$$

Thus $f^*\omega$ is an element of $\widetilde{F}(\underline{K})$. The compatibility with the perverse degree is direct. □

The next result is the key point in the existence of quasi-isomorphisms. It is based on the following definitions introduced in [21].

Definition 5.12. A universal local system, F, is *extendable* if the restriction map $F(\Delta^n) \to F(\partial\Delta^n)$ admits a section, for any n.

Definition 5.13. A universal local system, F, is of *differential coefficients* if, for any n and any i, the face operator $\partial_i\colon \Delta^{n-1} \to \Delta^n$ induces an isomorphism $\partial_i^*\colon H(F([n])) \to H(F([n-1]))$.

Theorem C. *Let R be a commutative ring. Let $\overline{q}$ be a loose perversity and F, G be two extendable universal local systems of differential coefficients, such that there exists a natural transformation, $\psi_n\colon F([n]) \to G([n])$, inducing an isomorphism in homology in the case $n = 0$ and such that $H^0(F([0]) = H^0(G([0]) = R$. Then a morphism $\Psi\colon \widetilde{F}_{\overline{q}}(\underline{K}) \to \widetilde{G}_{\overline{q}}(\underline{K})$ is defined by $(\Psi\omega)_\sigma = \psi_{|\sigma|}(\omega_\sigma)$ and the induced map*

$$\Psi^*\colon H^*_{\overline{q}}(\underline{K}; \widetilde{F}) \to H^*_{\overline{q}}(\underline{K}; \widetilde{G})$$

is an isomorphism.

The classical integration map, $\int\colon A_{PL}(\Delta^n) \to C^*(\Delta^n)$, defines an integration map between the blow-up's of the local systems of cochains and of Sullivan's forms,

$$\int\colon \widetilde{A}_{PL}(\underline{K}) \to \widetilde{C}^*(\underline{K}),$$

as follows. Let $\sigma\colon \Delta^{j_0} * \cdots * \Delta^{j_n} \to \underline{K}$ and $\omega \in \widetilde{A}_{PL}(\underline{K})$. If $\omega_\sigma = \omega_0 \otimes \cdots \otimes \omega_n \in A_{PL}(c\Delta^{j_0}) \otimes \cdots \otimes A_{PL}(\Delta^{j_n})$, we set

$$\left(\int \omega\right)_\sigma = \int_{c\Delta^{j_0}} \omega_0 \times \cdots \times \int_{\Delta^{j_n}} \omega_n = \int_{c\Delta^{j_0} \times \cdots \times \Delta^{j_n}} \omega_0 \wedge \cdots \wedge \omega_n.$$

The compatibility with the differentials is the classical Stokes' theorem. In the case of face sets, Sullivan proves that this integration induces a quasi-isomorphism in homology, see [35, Theorem 7.1]. This Sullivan's theorem and Theorem C imply the next result.

Corollary 5.14. *Let* $R=\mathbb{Q}$, $\underline{K}$ *be a filtered face set and* $\overline{q}$ *be a loose perversity. The integration map* $\int\colon \widetilde{A}_{PL,\overline{q}}(\underline{K})\to \widetilde{C}^*_{\overline{q}}(\underline{K})$ *induces an isomorphism in homology.*

6. Proof of Theorem C

> This section is devoted to the proof of Theorem C. For that, we introduce the notion of filtered theory of cochains that will be used also for Goresky-MacPherson cochains.

All cochain complexes are over a commutative ring R.

Definition 6.1. A functor C^* from the category of filtered face maps to the category of cochain complexes is a *filtered theory of cochains* if the following properties are satisfied.

(a) Extension axiom. For any filtered face set, $\underline{K}$, any $r\geq 0$ and any $k\geq 0$, the restriction map, $C^*(\underline{K}^{[r],k})\to C^*(\underline{K}^{[r],k-1})$, is surjective. The kernel is denoted by $C^*(\underline{K}^{[r],k},\underline{K}^{[r],k-1})$.

(b) Relative isomorphism axiom. Each relative isomorphism (see Definition 4.15), of the shape $f\colon (\underline{K}^{[r],k},\underline{K}^{[r],k-1})\to(\underline{L}^{[r],k},\underline{L}^{[r],k-1})$, induces an isomorphism $C^*(\underline{K}^{[r],k},\underline{K}^{[r],k-1})\cong C^*(\underline{L}^{[r],k},\underline{L}^{[r],k-1})$.

(c) Wedge axiom. For any union of pairs of filtered face sets, $\cup_i(\underline{K}_i^{[r],k},\underline{K}_i^{[r],k-1})$, such that the complementary subsets $\underline{K}_i^{[r],k}\backslash\underline{K}_i^{[r],k-1}$ are disjoint, we have an isomorphism

$$C^*(\cup_i\underline{K}_i^{[r],k},\cup_i\underline{K}_i^{[r],k-1}))\cong\prod_i C^*(\underline{K}_i^{[r],k},\underline{K}_i^{[r],k-1}).$$

(d) Filtered Dimension axiom. For any filtered face set, $\underline{K}$, and any r, we have $H^m(C^*(\underline{K}^{[r]},\underline{K}^{[r],k}))=0$ if $m<k$.

Definition 6.2. Let $\Psi\colon C^*\to D^*$ be a natural transformation between two filtered theories of cochains, such that $\Psi(\underline{K})\colon C^*(\underline{K})\to D^*(\underline{K})$ is a quasi-isomorphism for a filtered face set $\underline{K}$, of depth $v(\underline{K})<n$. The natural transformation Ψ is *cone-compatible for* $\underline{K}$ if the map

$$\Psi(\Delta^k*\underline{K})\colon C^*(\Delta^k*\underline{K})\to D^*(\Delta^k*\underline{K})$$

is a quasi-isomorphism, for any $k\geq 0$.

Proposition 6.3. *Let R be a commutative ring. Let $\Psi\colon C^*\to D^*$ be a natural transformation between two filtered theories of cochains, cone-compatible for any filtered face set of depth strictly less than n, and inducing an isomorphism in homology, $H^*(\Psi)\colon C^*(\{\vartheta\})\to D^*(\{\vartheta\})$, for any singleton $\{\vartheta\}$. Then $\Psi(\underline{K})\colon C^*(\underline{K})\to D^*(\underline{K})$ is a quasi-isomorphism for any filtered face set $\underline{K}$.*

Proof. Suppose the result is true for any filtered face set $\underline{L}$ such that $\underline{L} = \underline{L}^{[r],k-1}$. (Observe that the induction is satisfied if $r = 0$, $k = 1$.) The extension axiom gives a morphism of exact sequences

$$\begin{tikzcd}
0 \arrow[r] & C^*(\underline{K}^{[r],k}, \underline{K}^{[r],k-1}) \arrow[r] \arrow[d, "f_1"'] & C^*(\underline{K}^{[r],k}) \arrow[r] \arrow[d, "f_2"] & C^*(\underline{K}^{[r],k-1}) \arrow[r] \arrow[d, "f_3"] & 0 \\
0 \arrow[r] & D^*(\underline{K}^{[r],k}, \underline{K}^{[r],k-1}) \arrow[r] & D^*(\underline{K}^{[r],k}) \arrow[r] & D^*(\underline{K}^{[r],k-1}) \arrow[r] & 0.
\end{tikzcd}$$

The map f_3 is a quasi-isomorphism by induction. To prove that f_2 is one also, we are reduced to study the map f_1 induced between the kernels. Recall that $\mathfrak{J}(\underline{K}^{[r],k})$ is the set of $\tau = R_1(\sigma)$, when $\sigma \in \underline{K}^{[r],k} \backslash \underline{K}^{[r],k-1}$. From Example 4.16, the wedge and the relative isomorphism axioms, we get

$$C^*(\underline{K}^{[r],k}, \underline{K}^{[r],k-1}) = \prod_{\tau \in \mathfrak{J}(r,k)} C^*(\underline{K}(\tau), \underline{K}(\tau, \partial\tau)) \tag{3}$$

and a similar result for D^*. (In the case $k = 0$, we have $\underline{K}(\tau, \partial\tau) = \mathcal{L}(\underline{K}, \tau)$ and $\underline{K}^{[r],-1} = \underline{K}^{[r-1]}$.) From Proposition 4.18 and the relative isomorphism axiom, there exists an isomorphism between $C^*(\underline{K}(\tau), \underline{K}(\tau, \partial\tau))$ and $C^*(\Delta^k * \mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau), \partial\Delta^k * \mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau))$. With a similar argument on D^*, the induction hypothesis on $H^*(\Psi)$ and the cone-compatibility, the proof is done in the case $\underline{K}^{[r]} = \underline{K}^{[r],k}$ for some k.

The axiom of filtered dimension implies the nullity of the relative cohomologies $H^m(C(\underline{K}^{[r]}, \underline{K}^{[r],k}))$ and $H^m(D(\underline{K}^{[r]}, \underline{K}^{[r],k}))$, in any degree m for k great enough. Therefore, we get the isomorphism $H^m(C(\underline{K}^{[r]})) \cong H^m(D(\underline{K}^{[r]}))$, for any m. As $\underline{K} = \underline{K}^{[n]}$, the proof is done. $\square$

Proposition 6.3 admits a variation (used in Proposition 8.4) whose proof follows from an inspection of the previous arguments.

Proposition 6.4. *Let R be a commutative ring. Let $\Psi \colon C^* \to D^*$ be a natural transformation between two filtered theories of cochains, cone-compatible for any* connected *filtered face set of depth strictly less than n, and inducing an isomorphism in homology, $H^*(\Psi) \colon C^*(\{\vartheta\}) \to D^*(\{\vartheta\})$, for any singleton $\{\vartheta\}$. Then $\Psi(\underline{K}) \colon C^*(\underline{K}) \to D^*(\underline{K})$ is a quasi-isomorphism for any filtered face set $\underline{K}$ whose expanded links, $\mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)$, are connected.*

We determine a class of universal local system generating filtered theory of cochains by blow-up's.

Proposition 6.5. *Let F be an extendable universal local system of differential coefficients, such that $H^0(\{\vartheta\}) = R$, and $\overline{q}$ be a loose perversity. Then, the cochain complex $\widetilde{F}_{\overline{q}}$ is a filtered theory of cochains.*

Proof. We follow the properties of Definition 6.1.

- Extension axiom. Let $\underline{K}$ be a filtered face set and $\tau \in \mathfrak{J}(\underline{K}^{[r],k})$ with $k > 0$. We first prove that $\widetilde{F}_{\overline{q}}(\underline{K}(\tau)) \to \widetilde{F}_{\overline{q}}(\underline{K}(\tau, \partial\tau))$ is surjective.

Let $\sigma \in \underline{K}(\tau) \backslash \underline{K}(\tau, \partial\tau)$ and $\omega \in \widetilde{F}(\underline{K}(\tau, \partial\tau))$. The global section $\omega_{\partial^{\mathcal{F}}\sigma}$ defined on the filtered boundary $\partial^{\mathcal{F}}\sigma$ of σ can de decomposed as $\omega_{\partial^{\mathcal{F}}\sigma} = \sum_m \omega'_m \otimes \omega''_m$ with $\omega'_m \in F(c\partial\Delta^k)$ and $\omega''_m \in F(c\Delta^{j_{n-r+1}}) \otimes \cdots \otimes F(\Delta^{j_n})$. By hypothesis and [21, Proposition 12.21], the map $F(c\Delta^k) \to F(c\partial\Delta^k)$ admits a section, ρ, and we may define $\omega_\sigma = \sum_m \rho(\omega'_m) \otimes \omega''_m$. By construction, the correspondence $\sigma \mapsto \omega_\sigma$ is compatible with face operators and we get a global section $\omega \in \widetilde{F}(\underline{K}(\tau))$. By definition of the perverse degree on $\widetilde{F}$, we have $\|\rho(\omega)\|_i \leq \|\omega\|_i$ and $\|d\rho(\omega)\|_i \leq \|d\omega\|_i$ for all i. Therefore, we get an induced surjective map, $\widetilde{F}_{\overline{q}}(\underline{K}(\tau)) \to \widetilde{F}_{\overline{q}}(\underline{K}(\tau, \partial\tau))$.

We consider now the case $k = 0$. The map $F(\Delta^1) = F(c\{\vartheta\}) \to F(\partial\Delta^1) = F((\{\vartheta\} \times \{1\}) \sqcup (\{\vartheta\} \times \{0\}))$ being surjective, there exists $\hat{\alpha} \in F^0(c\{\vartheta\})$ such that

$$\hat{\alpha}_{|\vartheta\}\times\{1\}} = 0 \text{ and } \hat{\alpha}_{|\vartheta\}\times\{0\}} = 1.$$

Let $\sigma\colon \{\vartheta\} * \Delta^{j_{n-r+1}} * \cdots * \Delta^{j_n} \to \underline{K}$ be a simplex of $\underline{K}(\{\vartheta\})$. We identify the product $c\Delta^{j_{n-r+1}} \times \cdots \times \Delta^{j_n}$ to the subset $(\{\vartheta\} \times \{0\}) \times c\Delta^{j_{n-r+1}} \times \cdots \times \Delta^{j_n}$ of $c\{\vartheta\} \times c\Delta^{j_{n-r+1}} \times \cdots \times \Delta^{j_n}$ and denote by σ' the restriction of σ to $c\Delta^{j_{n-r+1}} \times \cdots \times \Delta^{j_n}$. Let $\omega \in \widetilde{F}(\mathcal{L}(\underline{K}, \{\vartheta\}))$. We set $\hat{\omega}_\sigma = \hat{\alpha} \otimes \omega_{\sigma'}$. This construction is compatible with the face operators and define an element $\hat{\omega} \in \widetilde{F}(\underline{K}(\{\vartheta\}))$ whose restriction to $\widetilde{F}(\mathcal{L}(\underline{K}, \{\vartheta\}))$ coincides with ω. As $H^0(\{\vartheta\}) = R$, the restriction of $d\hat{\alpha}$ to the boundary $\partial\Delta^1$ is equal to 0 and, for any $\omega' \in \widetilde{F}(\mathcal{L}(\underline{K}, \{\vartheta\}))$, we have

$$\|d\hat{\alpha} \otimes \omega'\|_\ell \leq \|\hat{\alpha} \otimes \omega')\|_\ell = \begin{cases} -\infty & \text{if} \quad \ell \geq r, \\ \|\omega'\|_\ell & \text{if} \quad \ell < r. \end{cases}$$

The surjectivity of $\widetilde{F}_{\overline{q}}(\underline{K}(\{\vartheta\})) \to \widetilde{F}_{\overline{q}}(\mathcal{L}(\underline{K}, \{\vartheta\}))$ follows.

Let $k \geq 0$. We detail only the case $k > 0$, the case $k = 0$ being similar here. Observe that $\underline{K}^{[r],k}$ can be obtained as a pushout, built on disjoint unions,

$$\begin{tikzcd} \sqcup_{\tau\in\mathfrak{J}(r,k)}\underline{K}(\tau, \partial\tau) \arrow[r] \arrow[d] & \sqcup_{\tau\in\mathfrak{J}(r,k)}\underline{K}(\tau) \arrow[d] \\ \underline{K}^{[r],k-1} \arrow[r] & \underline{K}^{[r],k}. \end{tikzcd} \tag{4}$$

By definition, the blow-up $\widetilde{F}_{\overline{q}}$ is compatible with colimits and we get a pullback

$$\begin{tikzcd} \widetilde{F}_{\overline{q}}\left(\sqcup_{\tau\in\mathfrak{J}(r,k)}\underline{K}(\tau, \partial\tau)\right) & \widetilde{F}_{\overline{q}}\left(\sqcup_{\tau\in\mathfrak{J}(r,k)}\underline{K}(\tau)\right) \arrow[l] \\ \widetilde{F}_{\overline{q}}\left(\underline{K}^{[r],k-1}\right) \arrow[u] & \widetilde{F}_{\overline{q}}\left(\underline{K}^{[r],k}\right). \arrow[l] \arrow[u] \end{tikzcd}$$

By definition also, we have $\widetilde{F}_{\overline{q}}\left(\sqcup_{\tau\in\mathfrak{J}(r,k)}\underline{K}(\tau, \partial\tau)\right) = \prod_{\tau\in\mathfrak{J}(r,k)} \widetilde{F}_{\overline{q}}(\underline{K}(\tau, \partial\tau)$ and $\widetilde{F}_{\overline{q}}\left(\sqcup_{\tau\in\mathfrak{J}(r,k)}\underline{K}(\tau)\right) = \prod_{\tau\in\mathfrak{J}(r,k)} \widetilde{F}_{\overline{q}}(\underline{K}(\tau))$. The surjectivity of $\widetilde{F}_{\overline{q}}(\underline{K}(\tau)) \to \widetilde{F}_{\overline{q}}(\underline{K}(\tau, \partial\tau))$ implies the surjectivity of $\widetilde{F}_{\overline{q}}(\underline{K}^{[r],k}) \to \widetilde{F}_{\overline{q}}(\underline{K}^{[r],k-1})$.

If $\underline{L}$ and $\underline{L}'$ are filtered face sets, connected by a pushout,

$$\begin{tikzcd} \sqcup_{\tau\in I}\underline{K}(\tau, \partial\tau) \arrow[r] \arrow[d] & \sqcup_{\tau\in I}\underline{K}(\tau) \arrow[d] \\ \underline{L}' \arrow[r] & \underline{L}, \end{tikzcd}$$

as in (4), the previous argument implies the surjectivity of $\widetilde{F}_{\overline{q}}(\underline{L}) \to \widetilde{F}_{\overline{q}}(\underline{L}')$. We will use that property several times in the proof of the last axiom.

• Relative isomorphism axiom.
The relative isomorphism $f\colon (\underline{K}^{[r],k}, \underline{K}^{[r],k-1}) \to (\underline{L}^{[r],k}, \underline{L}^{[r],k-1})$ induces a morphism of complexes

$$f^*\colon \widetilde{F}_{\overline{q}}(\underline{L}^{[r],k}, \underline{L}^{[r],k-1}) \to \widetilde{F}_{\overline{q}}(\underline{K}^{[r],k}, \underline{K}^{[r],k-1}).$$

If $\eta \in \widetilde{F}_{\overline{q}}(\underline{K}^{[r],k}, \underline{K}^{[r],k-1})$ and $\sigma \in \underline{L}^{[r],k}$, we set

$$\omega_\sigma = \left\{ \begin{array}{ccl} \eta_{f^{-1}(\sigma)} & \text{if} & \sigma \in \underline{L}^{[r],k} \backslash \underline{L}^{[r],k-1}, \\ 0 & \text{if} & \sigma \in \underline{L}^{[r],k-1}. \end{array} \right.$$

As $f^{-1}(\sigma)$ exists and is uniquely defined, the element ω is well defined. We have to check the compatibility with face operators. Let ∂_j be a face map and $\sigma \in \underline{L}^{[r],k}$.

- if $\partial_j\sigma \in \underline{L}^{[r],k} \backslash \underline{L}^{[r],k-1}$, then we have
$$\omega_{\partial_j\sigma} = \eta_{f^{-1}(\partial_j\sigma)} = \eta_{\partial_j f^{-1}(\sigma)} = \partial_j^*\eta_{f^{-1}(\sigma)} = \partial_j^*\omega_\sigma.$$
(We used $\partial_j f^{-1}(\sigma) = f^{-1}(\partial_j\sigma)$ which can be verified by composing with f.)
- if $\partial_j\sigma \in \underline{L}^{[r],k-1}$, then we have $\partial_j^*\omega_\sigma = \partial_j^*\eta_{f^{-1}(\sigma)} = \eta_{\partial_j f^{-1}(\sigma)} = 0$, because $\partial_j f^{-1}(\sigma) \in \underline{K}^{[r],k-1}$. (If $\partial_j f^{-1}(\sigma) \in \underline{K}^{[r],k} \backslash \underline{K}^{[r],k-1}$ then we should have $\partial_j\sigma \in \underline{L}^{[r],k} \backslash \underline{L}^{[r],k-1}$.) By construction, we have also $\omega_{\partial_j\sigma} = 0$.

• Wedge axiom. The restriction maps give a morphism of complexes

$$\widetilde{F}_{\overline{q}}(\cup_i \underline{K}_i^{[r],k}, \cup_i \underline{K}_i^{[r],k-1}) \to \prod_i \widetilde{F}_{\overline{q}}(\underline{K}_i^{[r],k}, \underline{K}_i^{[r],k-1}).$$

This morphism admits an inverse: for any family $(\omega_i)_i \in \prod_i \widetilde{F}_{\overline{q}}(\underline{K}_i^{[r],k}, \underline{K}_i^{[r],k-1})$, we define a global section $\omega \in \widetilde{F}_{\overline{q}}(\cup_i \underline{K}_i^{[r],k}, \cup_i \underline{K}_i^{[r],k-1})$ by

$$\omega_\sigma = \left\{ \begin{array}{cl} (\omega_i)_\sigma & \text{if } \sigma \in \underline{K}_i^{[r],k} \backslash \underline{K}_i^{[r],k-1}, \\ 0 & \text{otherwise.} \end{array} \right.$$

The indice i such that $\sigma \in \underline{K}_i^{[r],k} \backslash \underline{K}_i^{[r],k-1}$ being unique, the element ω_σ is well defined. We have to check its compatibility with face operators. Let $\sigma \in \underline{K}_i^{[r],k}$.

- If $\partial_j\sigma \in \underline{K}_i^{[r],k} \backslash \underline{K}_i^{[r],k-1}$, we have $\omega_{\partial_j\sigma} = (\omega_i)_{\partial_j\sigma} = \partial_j^*(\omega_i)_\sigma = \partial_j^*\omega_\sigma$.
- If $\partial_j\sigma \in \underline{K}_i^{[r],k-1}$, we have $\omega_{\partial_j\sigma} = 0$ and $\partial_j^*\omega_\sigma = \partial_j^*(\omega_i)_\sigma = (\omega_i)_{\partial_j\sigma} = 0$, because $\omega_i \in \widetilde{F}_{\overline{q}}(\underline{K}_i^{[r],k}, \underline{K}_i^{[r],k-1})$.

• Filtered dimension axiom. We have to prove the nullity of the relative cohomology with coefficients in $\widetilde{F}_{\overline{q}}$, i.e., $H_{\overline{q}}^m(\underline{K}^{[r]}, \underline{K}^{[r],k}; \widetilde{F}) = 0$ for $m < k$. We establish it in three steps.

(i) If $\tau \in \mathcal{I}(\underline{K}^{[r],k})$ and $m < k$, we prove

$$H^m_{\overline{q}}(\underline{K}(\tau), \underline{K}(\tau, \partial\tau); \widetilde{F}) = 0.$$

We denote by $\Lambda^{m,k}$ the subcomplex of Δ^k generated by the simplices containing the vertex v_0 and of dimension less than, or equal, to m and by $\Lambda^{m,\tau}$ the restriction of τ to $\Lambda^{m,k}$. Let $f_i \colon \Delta^m \to \Lambda^{m,k}$, $i \in I$, the m-dimensional simplices of $\Lambda^{m,k}$. We set $\tau_i = \Lambda^{m,\tau} \circ f_i$ and define $\underline{K}(\Lambda^{m,\tau}) = \cup_{i\in I}\underline{K}(\tau_i)$. We first prove that

$$H^*_{\overline{q}}(\underline{K}(\tau), \underline{K}(\Lambda^{m,\tau}); \widetilde{F}) = 0, \text{ for any } m,\ 0 \leq m \leq k. \tag{5}$$

This equality is true for $\tau \in \mathcal{I}(\underline{K}^{[r],0})$. We use an induction on the dimension $|\tau|$ of τ, by supposing that (5) is true for any τ' of dimension less than $|\tau|$. The result for τ is true for $m = k$. We do a new induction, supposing the result for τ is true for m, $m \leq k$, and proving it for $m-1$. As $\widetilde{F_{\overline{q}}}$ verifies the extension axiom, we are reduced to prove that

$$H^*_{\overline{q}}(\underline{K}(\Lambda^{m,\tau}), \underline{K}(\Lambda^{m-1,\tau}); \widetilde{F}) = 0, \tag{6}$$

which is a consequence of the inductive step by using the isomorphism,

$$\widetilde{F}_{\overline{q}}(\underline{K}(\Lambda^{m,\tau}), \underline{K}(\Lambda^{m-1,\tau})) \cong \prod_{i\in I} \widetilde{F}_{\overline{q}}(\underline{K}(\tau_i), \underline{K}(\Lambda^{m-1,\tau_i})).$$

The equality (5) is proved.

Consider now $\iota \colon \Delta^{k-1} \to \Delta^k$, the face opposite to v_0 and set $\tau' = \tau \circ \iota$. With the extension axiom, we have an exact sequence,

$$\begin{tikzcd} 0 \arrow[r] & \widetilde{F}_{\overline{q}}(\underline{K}(\tau), \underline{K}(\tau, \partial\tau)) \arrow[r] & \widetilde{F}_{\overline{q}}(\underline{K}(\tau), \underline{K}(\Lambda^{k-1,\tau})) \arrow[r] & \widetilde{F}_{\overline{q}}(\underline{K}(\tau'), \underline{K}(\tau', \partial\tau')) \arrow[r] & 0. \end{tikzcd}$$

The equality (5) implies the existence of an isomorphism of degree $+1$ between the relative cohomologies involving τ and τ'. Starting from $\tau' = \{\vartheta\}$ and $\widetilde{F}_{\overline{p}}(\underline{K}(\tau'), \underline{K}(\tau', \partial\tau')) = \widetilde{F}_p(\underline{K}(\vartheta), \mathcal{L}(\underline{K}, \vartheta))$, whose cohomology is zero in negative degree, we deduce

$$H^m_{\overline{q}}(\underline{K}(\tau), \underline{K}(\tau, \partial\tau); \widetilde{F}) = 0, \text{ if } m < |\tau|,$$

and the result follows.

(ii) As the map $(\cup_{\tau\in\mathcal{I}(\underline{K}^{[r],k})}\underline{K}(\tau), \cup_{\tau\in\mathcal{I}(\underline{K}^{[r],k})}\underline{K}(\tau, \partial\tau)) \to (\underline{K}^{[r],k}, \underline{K}^{[r],k-1})$ is a relative isomorphism, the previous step, the wedge and the relative isomorphism axioms imply

$$H^m_{\overline{q}}(\underline{K}^{[r],k}, \underline{K}^{[r],k-1}; \widetilde{F})) = \prod_{\tau\in\mathcal{I}(\underline{K}^{[r],k})} H^m_{\overline{q}}(\underline{K}(\tau), \underline{K}(\tau, \partial\tau), \widetilde{F}) = 0,$$

for any m, $m < k$.

(iii) Finally, we prove $H^m_{\overline{q}}(\underline{K}^{[r]}, \underline{K}^{[r],k}; \widetilde{F}) = 0$, if $m < k$.

Let $\omega \in \widetilde{F}_{\overline{q}}(\underline{K}^{[r]}, \underline{K}^{[r],k})$, of degree m, with $d\omega = 0$. The restriction $\omega_{|\underline{K}^{[r],k+1}}$ is a coboundary (with step (ii)) and there exists $\psi_0 \in \widetilde{F}_{\overline{q}}(\underline{K}^{[r],k+1}, \underline{K}^{[r],k})$ such that $\omega_{|\underline{K}^{[r],k+1}} = d\psi_0$. By iterating the extension axiom, the global section ψ_0 can be extended in a global section defined on $\underline{K}^{[r]}$ that we still denote ψ_0. By induction, we find a family of global sections $\psi_i \in \widetilde{F}_{\overline{q}}(\underline{K}^{[r]}, \underline{K}^{[r],k+i})$ so that $\omega - d(\sum_0^j \psi_i) \in$

$\widetilde{F}_{\overline{q}}(\underline{K}^{[r]},\underline{K}^{[r],k+j+1})$. We set $\psi=\sum_{i=0}^{\infty}\psi_i$. When we apply ψ to a simplex $\sigma\in\underline{K}(\tau)$, the sum ψ_σ is finite and we have $\psi\in\widetilde{F}_{\overline{q}}(\underline{K}^{[r]},\underline{K}^{[r],k})$ with $\omega=d\psi$.

□

Proposition 6.6. *Let $\underline{K}$ be a filtered face set of depth $v(\underline{K})=\ell-1<n$ and $\Delta^k*\underline{K}$ be a join with $k\geq 0$. Let F be a universal local system of coefficients and $\overline{q}$ be a loose perversity. Then we have*

$$H^i_{\overline{q}}(\Delta^k*\underline{K};\widetilde{F})=\begin{cases}\oplus_{i_1+i_2=i}H^{i_1}(F(\{\vartheta\})\otimes H^{i_2}_{\overline{q}}(\underline{K};\widetilde{F}) & \textit{if } i\leq\overline{q}(\ell),\\ 0 & \textit{if } i>\overline{q}(\ell).\end{cases}$$

Proof. First, we claim the existence of an isomorphism of cochain complexes

$$\widetilde{F}(\Delta^k*\underline{K})\cong F(c\Delta^k)\otimes\widetilde{F}(\underline{K}).$$

Observe that an element of $F(c\Delta^k)$ is entirely determined by its value on the maximal simplex $[c\Delta^k]$ and that any element $\eta\in F(c\Delta^k)\otimes\widetilde{F}(\underline{K})$ can be decomposed as a sum of $\eta'\otimes\eta''$ with $\eta'\in F(c\Delta^k)$ and $\eta''\in\widetilde{F}(\underline{K})$.

If $\omega\in\widetilde{F}(\Delta^k*\underline{K})$ and $\alpha\colon\Delta^{j_{n-\ell+1}}*\cdots*\Delta^{j_n}\to\underline{K}$, we set $\eta'_{[c\Delta^k]}\otimes\eta''_\alpha=\omega_{\mathrm{id}*\alpha}$. Reciprocally, if $\eta\in F(c\Delta^k)\otimes\widetilde{F}(\underline{K})$ and

$$\sigma=(\partial_0)^{k-k'}*\alpha\colon\Delta^{k'}*\Delta^{j_{n-\ell+1}}*\cdots*\Delta^{j_n}\to\Delta^k*\underline{K},$$

we define $\omega\in\widetilde{F}(\Delta^k*\underline{K})$ by $\omega_\sigma=((\partial_0)^{k-k'})^*\eta'_{[c\Delta^k]}\otimes\eta''_\alpha$. This proves the claim.

We introduce now the perversity $\overline{q}$. Let $\alpha\colon\Delta^{j_{n-\ell+1}}*\cdots*\Delta^{j_n}\to\underline{K}$ and $\eta\in F(c\Delta^k)\otimes\widetilde{F}(\underline{K})$. We decompose

$$\eta_{\Delta^k*\alpha}=\sum_i\eta'_{i,\alpha}\otimes\eta''_{i,\alpha}\in F(c\Delta^k)\otimes\widetilde{F}(\underline{K}),$$

By definition, the perverse degree of $\eta_{\Delta^k*\alpha}$ is determined as follows:

- if $\ell<\ell'$, then $\|\eta_{\Delta^k*\alpha}\|_{\ell'}=-\infty$,
- if $\ell'=\ell$, then,
  - if $\eta'_{i,\alpha|\Delta^k\times\{1\}}\otimes\eta''_{i,\alpha}=0$ for all i, we set $\|\eta_{\Delta^k*\alpha}\|_\ell=-\infty$,
  - if not, we set
    $$\|\eta_{\Delta^k*\alpha}\|_\ell=\max_i\{|\eta''_{i,\alpha}|\text{ such that }\eta'_{i,\alpha|\Delta^k\times\{1\}}\neq 0\},$$
- if $\ell>\ell'$, then,
  - if $\eta'_{i,\alpha}=0$ for all i, we set $\|\eta_{\Delta^k*\alpha}\|_{\ell'}=-\infty$,
  - if not, we set
    $$\|\eta_{\Delta^k*\alpha}\|_{\ell'}=\max_i\left\{\|\eta''_{i,\alpha}\|_{\ell'}\text{ such that }\eta'_{i,\alpha}\neq 0\right\},$$

  and the perverse degree of η coincide with the perverse degree of its component in $\widetilde{F}(\underline{K})$.

This implies the decomposition of $\widetilde{F}_{\overline{q}}(\Delta^k*\underline{K})$ as

$$\begin{aligned}\widetilde{F}_{\overline{q}}(\Delta^k*\underline{K}) \;=\;& \left(F(c\Delta^k)\otimes\widetilde{F}^{<\overline{q}(\ell)}_{\overline{q}}(\underline{K})\right)\oplus\left(F(c\Delta^k)\otimes\mathcal{Z}\widetilde{F}^{\overline{q}(\ell)}_{\overline{q}}(\underline{K})\right)\\ &\oplus\left(F(c\Delta^k,\Delta^k\times\{1\})\otimes(\widetilde{F}^{>\overline{q}(\ell)}_{\overline{q}}(\underline{K})\oplus\mathcal{T}\widetilde{F}^{\overline{q}(\ell)}_{\overline{q}}(\underline{K}))\right),\end{aligned}$$

where $\mathcal{Z}V$ denotes the vector space of cocycles of a cochain complex V and $\mathcal{T}V$ is a supplementary of $\mathcal{Z}V$ in V. As the universal local system F is of differentiable coefficients, we have $H^*F(c\Delta^k, \Delta^k \times \{1\}) = 0$. Therefore, the complex $\widetilde{F}^i_{\overline{q}}(\Delta^k * \underline{K})$ is quasi-isomorphic to

$$\left(F(c\Delta^k) \otimes \widetilde{F}^{<\overline{q}(\ell)}_{\overline{q}}(\underline{K})\right) \oplus \left(F(c\Delta^k) \otimes \mathcal{Z}\widetilde{F}^{\overline{q}(\ell)}_{\overline{q}}(\underline{K})\right).$$

The result follows now from the isomorphism, $H^*(F(c\Delta^k)) \cong H^*(F(\{\vartheta\}))$, coming from the hypothesis of differential coefficients. □

Proof of Theorem C. Let $\sigma\colon \Delta^{j_0} * \cdots * \Delta^{j_n} \to \underline{K}$. We define

$$\psi_\sigma\colon \widetilde{F}_\sigma = F(c\Delta^{j_0}) \otimes \cdots \otimes F(\Delta^{j_n}) \to \widetilde{G}_\sigma = G(c\Delta^{j_0}) \otimes \cdots \otimes G(\Delta^{j_n})$$

by $\psi_\sigma = \psi_{j_0+1} \otimes \cdots \otimes \psi_{j_n}$. By construction, the map ψ decreases the perverse degree and induces a cochain map $\Psi(\underline{K})\colon \widetilde{F}_{\overline{q}}(\underline{K}) \to \widetilde{G}_{\overline{q}}(\underline{K})$.

The hypothesis of Proposition 6.3 are satisfied: the blow-up's $\widetilde{F}_{\overline{q}}$ and $\widetilde{G}_{\overline{q}}$ are filtered theories of cochains thanks to Proposition 6.5 and the natural transformation $\widetilde{F}_{\overline{q}}(-) \to \widetilde{G}_{\overline{q}}(-)$ is cone-compatible for any filtered face set, thanks to Proposition 6.6. □

7. COCHAINS ON FILTERED FACE SETS

We compare two cochain complexes defined over a commutative ring R. The first one is the blow-up, $\widetilde{C}^*$, of the classical cochains on a face set, previously studied. The second one is the dual of a complex, $C^{\mathrm{GM},\overline{p}}_*$, whose homology is the Goresky and MacPherson homology in the case of a filtered face set associated to a locally conelike space. The main result (Theorem D) is the existence of a quasi-isomorphism, over R, between these two complexes.

Let $\underline{K}$ be a filtered face set. The chain complex $C^{\mathrm{GM}}_*(\underline{K})$ is defined as follows,

- as module, $C^{\mathrm{GM}}_*(\underline{K})$ is the free R-module generated by the simplices $\sigma \in \underline{K}$,
- any simplex $\sigma\colon \Delta^{j_0} * \cdots * \Delta^{j_n} \to \underline{K}$ has a perverse degree defined by $\|\sigma\|_\ell = \dim(\Delta^{j_0} * \cdots * \Delta^{j_{n-\ell}})$, if $\Delta^{j_0} * \cdots * \Delta^{j_{n-\ell}} \neq \emptyset$ and $\|\sigma\|_\ell = -\infty$ otherwise,
- the differential of a simplex $\sigma\colon \Delta^k = \Delta^{j_0} * \cdots * \Delta^{j_n} \to \underline{K}$ is given as usual by $\partial\sigma = \sum_{i=0}^k (-1)^i \sigma \circ \partial_i$, with $\partial_i \in \mathbf{\Delta}^{[n]}_{\mathcal{F}}$.

Let $\overline{p}$ be a loose perversity. Recall that a simplex $\sigma\colon \Delta = \Delta^{j_0} * \cdots * \Delta^{j_n} \to \underline{K}$ is $\overline{p}$-admissible if

$$\dim(\Delta^{j_0} * \cdots * \Delta^{j_{n-\ell}}) = \|\sigma\|_\ell \leq \dim \Delta - \ell + \overline{p}(\ell),$$

for any ℓ, $0 \leq \ell \leq n$. A chain c is $\overline{p}$-admissible if c can be written as a linear combination of $\overline{p}$-admissible simplices. A chain c is of $\overline{p}$-intersection if c and ∂c are $\overline{p}$-admissible. We denote by $C^{\mathrm{GM},\overline{p}}_*(\underline{K})$ the complex of intersection chains for $\overline{p}$. The *GM-cochain complexes* are the duals of these complexes, i.e.,

$$C^*_{\mathrm{GM}}(\underline{K}) = \hom(C^{\mathrm{GM}}_*(\underline{K}), R) \text{ and } C^*_{\mathrm{GM},\overline{p}}(\underline{K}) = \hom(C^{\mathrm{GM},\overline{p}}_*(\underline{K}), R).$$

We denote by $H_*^{\mathrm{GM},\overline{p}}(\underline{K};R)$ the homology of the complex $C_*^{\mathrm{GM},\overline{p}}(\underline{K})$, by $H^*_{\mathrm{GM},\overline{p}}(\underline{K};R)$ the homology of the complex $C^*_{\mathrm{GM},\overline{p}}(\underline{K})$ and called them, respectively, Goresky-MacPherson homology and cohomology of $\underline{K}$.

If X is a filtered space and $\underline{L} = \mathrm{ISing}_*^{\mathcal{F}}(X)$, as in Example 4.5, we have $C^*_{\mathrm{GM},\overline{p}}(\underline{L}) = \hom(C_*^{\overline{p}}(X),R)$.

Proposition 7.1. *Let R be a commutative ring and $\overline{p}$ be a loose perversity. The* GM-cochain complex $C^*_{\mathrm{GM},\overline{p}}(-)$ *is a filtered theory of cochains.*

Proof. We establish the properties of Definition 6.1.

- Extension axiom Let $\tau \in \mathfrak{J}(\underline{K}^{[r],k})$. We denote by $\mathcal{R}$ the map induced by the canonical inclusion,

$$\mathcal{R}\colon \hom(C_*^{\mathrm{GM},\overline{p}}(\underline{K}^{[r],k}),R) \to \hom(C_*^{\mathrm{GM},\overline{p}}(\underline{K}^{[r],k-1}),R).$$

A chain $c \in C_*(\underline{K}^{[r],k})$ can be decomposed as $c = c_1 + c_2$, with $c_1 \in C_*(\underline{K}^{[r],k-1})$ and $c_2 \in C_*(\underline{K}^{[r],k}\backslash\underline{K}^{[r],k-1})$. If c is of $\overline{p}$-intersection, then c_1, c_2 are $\overline{p}$-admissible and the boundary, ∂c, can be written as a linear combination of $\overline{p}$-admissible simplices. We decompose ∂c_2 in $\partial c_2 = \partial' c_2 + \partial'' c_2$, with $\partial' c_2 \in C_*(\underline{K}^{[r],k-1})$ and $\partial'' c_2 \in C_*(\underline{K}^{[r],k}\backslash\underline{K}^{[r],k-1})$. The elements of the linear combination $\partial'' c_2$ cannot be canceled with elements coming from $\partial' c_2$ or ∂c_1, therefore $\partial'' c_2$ is $\overline{p}$-admissible.

Let ∂_i be any face operator and σ be a $\overline{p}$-admissible simplex. If we have strict inequalities, $\|\partial_i\sigma\|_\ell < \|\sigma\|_\ell$, for any ℓ, then, directly from the definition, we observe that $\partial_i\sigma$ is $\overline{p}$-admissible. These strict inequalities occuring when ∂_i is a face operator acting on the first factor $\Delta^{j_{n-r}}$ of $\Delta^{j_{n-r}} * \cdots * \Delta^{j_n}$, we get the $\overline{p}$-admissibility of $\partial' c_2$. We have proved that c_2 is of $\overline{p}$-intersection; therefore $c_1 = c - c_2$ is of $\overline{p}$-intersection also.

This property allows the construction of a section, s, to $\mathcal{R}$, as follows: Let $\Phi \in \hom(C_*^{\mathrm{GM},\overline{p}}(\underline{K}^{[r],k-1}),R)$, we define $s(\Phi) \in \hom(C_*^{\mathrm{GM},\overline{p}}(\underline{K}^{[r],k}),R)$ by $s(\Phi)(c_1+c_2) = \Phi(c_1)$.

- Relative isomorphism axiom. Let

$$f\colon (\underline{K}^{[r],k},\underline{K}^{[r],k-1}) \to (\underline{L}^{[r],k},\underline{L}^{[r],k-1})$$

be a relative isomorphism preserving the perverse degree. The relative isomorphism axiom is equivalent to the fact that f induces an isomorphism between the quotients,

$$\overline{f}_*\colon \frac{C_*^{\mathrm{GM},\overline{p}}(\underline{K}^{[r],k})}{C_*^{\mathrm{GM},\overline{p}}(\underline{K}^{[r],k-1})} \xrightarrow{\cong} \frac{C_*^{\mathrm{GM},\overline{p}}(\underline{L}^{[r],k})}{C_*^{\mathrm{GM},\overline{p}}(\underline{L}^{[r],k-1})}.$$

Let $c \in C_*(\underline{K}^{[r],k})$. As in the previous item, we have a unique decomposition of c as $c = c' + c''$, with $c' \in C_*(\underline{K}^{[r],k-1})$ and c'' a linear combination of simplices of $\underline{K}^{[r],k}\backslash\underline{K}^{[r],k-1}$. As previously, we can also prove that if $c \in C_*^{\mathrm{GM},\overline{p}}(\underline{K}^{[r],k})$ then $c' \in C_*^{\mathrm{GM},\overline{p}}(\underline{K}^{[r],k-1})$ and c'' is a linear combination of simplices of $\underline{K}^{[r],k}\backslash\underline{K}^{[r],k-1}$ that is of $\overline{p}$-intersection. A similar decomposition exists for the elements of $C_*^{\mathrm{GM},\overline{p}}(\underline{L}^{[r],k})$.

We denote by $[c]$ the class of $c \in C_*^{\mathrm{GM},\overline{p}}(\underline{K}^{[r],k})$ in the quotient above. As f is a bijection between $\underline{K}^{[r],k}\backslash\underline{K}^{[r],k-1}$ and $\underline{L}^{[r],k}\backslash\underline{L}^{[r],k-1}$, we have $\overline{f}_*([c]) =$

$\overline{f}_*([c''])$, which is a linear combination of simplices in $\underline{L}^{[r],k}\backslash\underline{L}^{[r],k-1}$, that is of $\overline{p}$-intersection. This implies that $\overline{f}_*$ is an isomorphism.

• Wedge axiom. The dual of a direct sum being a product, we have to prove that the canonical injections induce an isomorphism,

$$\psi\colon \bigoplus_i \frac{C_*^{\mathrm{GM},\overline{p}}\left(\underline{K}_i^{[r],k}\right)}{C_*^{\mathrm{GM},\overline{p}}\left(\underline{K}_i^{[r],k-1}\right)} \xrightarrow{\cong} \frac{C_*^{\mathrm{GM},\overline{p}}\left(\cup_i\underline{K}_i^{[r],k}\right)}{C_*^{\mathrm{GM},\overline{p}}\left(\cup_i\underline{K}_i^{[r],k-1}\right)}.$$

The map ψ is defined by $\psi(([c_i])_i)=\sum_i [c_i]$, where $[-]$ denotes the equivalence classes of the quotients.

As we already did, we decompose a chain $c\in C_*^{\mathrm{GM},\overline{p}}\left(\cup_i\underline{K}_i^{[r],k}\right)$ in $c=c'+c''$, with $c'\in C_*^{\mathrm{GM},\overline{p}}\left(\cup_i\underline{K}_i^{[r],k-1}\right)$ and $c''\in C_*^{\mathrm{GM},\overline{p}}\left(\cup_i\underline{K}_i^{[r],k}\backslash \cup_i \underline{K}_i^{[r],k-1}\right)$. As the intersection of the sets, $\underline{K}_i^{[r],k}\backslash\underline{K}_i^{[r],k-1}$, is empty, the element c'' can be written $c''=\sum_i c_i''$, in a unique way, with $c_i''\in C_*^{\mathrm{GM},\overline{p}}\left(\underline{K}_i^{[r],k}\backslash\underline{K}_i^{[r],k-1}\right)$. The decompositions $c=c'+c''$ and $c''=\sum_i c_i''$ being unique, we have built an inverse to ψ, defined by $\psi^{-1}([c])=([c_i''])_i$.

• Filtered dimension axiom. Let $c\in C_*^{\mathrm{GM},\overline{p}}(\underline{K}^{[r]})$ of homological degree $|c|$ such that $|c|\leq k$. Denote by $[c]$ the class of c in the quotient

$$\frac{C_*^{\mathrm{GM},\overline{p}}(\underline{K}^{[r]})}{C_*^{\mathrm{GM},\overline{p}}(\underline{K}^{[r],k})}.$$

For degree reason, we have $[c]=0$ and the filtered dimension axiom is proved. □

Proposition 7.2. *Let R be a commutative ring. Let $\underline{K}$ be a filtered face set of depth $v(\underline{K})=\ell-1$ with $\ell\in\{1,\dots,n\}$. For any perversity $\overline{p}$ verifying $\overline{p}(\ell)\leq \ell-2$ and any $k\in\mathbb{N}$, we have*

$$H^i_{\mathrm{GM},\overline{p}}(\Delta^k*\underline{K};R)=\begin{cases} H^i_{\mathrm{GM},\overline{p}}(\underline{K};R) & \text{if } i\leq \ell-1-\overline{p}(\ell),\\ 0 & \text{if } i>\ell-1-\overline{p}(\ell).\end{cases}$$

Proof. Let $\sigma\colon\Delta^N=\Delta^{j_{n-\ell+1}}*\dots*\Delta^{j_n}\to\underline{K}$ and $\gamma\colon\Delta^{k'}\to\Delta^k$. We define

$$\psi\colon C_*(\Delta^k)\otimes C_*(\underline{K})\to C_{*+1}(\Delta^k*\underline{K})$$

by $\psi(\gamma\otimes\sigma)=\gamma*\sigma$. (Observe that this map is not compatible with face operators when γ or σ is a point.) The perverse degree of $\gamma*\sigma$ is determined by (see Example 4.17)

$$\|\gamma*\sigma\|_{\ell'}=\begin{cases} -\infty & \text{if } \ell<\ell',\\ k' & \text{if } \ell'=\ell,\\ & \text{or } (\ell>\ell' \text{ and } \|\sigma\|_{\ell'}=-\infty),\\ k'+\|\sigma\|_{\ell'}+1 & \text{if } \ell>\ell' \text{ and } \|\sigma\|_{\ell'}\neq-\infty.\end{cases}$$

By definition, the $\overline{p}$-admissibility condition of $\gamma*\sigma$ is

$$\|\gamma*\sigma\|_{\ell'}\leq k'+N+1-\ell'+\overline{p}(\ell'),\ \text{for any } \ell'. \tag{7}$$

If $\ell>\ell'$ and $\|\sigma\|_{\ell'}\neq-\infty$, the condition (7) is satisfied if, and only if, $\|\sigma\|_{\ell'}\leq N-\ell'+\overline{p}(\ell')$. This last condition is exactly the $\overline{p}$-admissibility of σ at ℓ'.

If $\ell' = \ell$ or ($\ell > \ell'$ and $\|\sigma\|_{\ell'} = -\infty$), the condition (7) is equivalent to $N \geq \ell' - 1 - \overline{p}(\ell')$. As $\overline{p}$ is a perversity, these conditions on ℓ' are equivalent to $N \geq \ell - 1 - \overline{p}(\ell)$.

As in [17, 1.14], we define a truncation of $C_*(\underline{K})$ by setting,

$$\tau_\ell^{\overline{p}} C_*(\underline{K}) = \left\{ \begin{array}{cl} 0 & \text{if } * < \ell - 1 - \overline{p}(\ell), \\ \mathfrak{Z}C_*^{\mathrm{GM},\overline{p}}(\underline{K}) & \text{if } * = \ell - 1 - \overline{p}(\ell), \\ C_*^{\mathrm{GM},\overline{p}}(\underline{K}) & \text{if } * > \ell - 1 - \overline{p}(\ell). \end{array} \right.$$

The previous calculations of perverse degree imply that the map

$$\psi \colon C_*(\Delta^k) \otimes \tau_\ell^{\overline{p}} C_*(\underline{K}) \to C_{*+1}^{\mathrm{GM},\overline{p}}(\Delta^k * \underline{K})$$

is well defined and induces

$$\overline{\psi} \colon C_*(\Delta^k) \otimes \tau_\ell^{\overline{p}} C_*(\underline{K}) \to \frac{C_{*+1}^{\mathrm{GM},\overline{p}}(\Delta^k * \underline{K})}{C_{*+1}^{\mathrm{GM},\overline{p}}(\underline{K})}.$$

Observe now that $\gamma * \{\vartheta\} \in C_*^{\mathrm{GM},\overline{p}}(\Delta^k * \underline{K})$ implies $0 \geq \ell - 1 - \overline{p}(\ell)$ which is in contradiction with the hypothesis $\ell - 2 - \overline{p}(\ell) \geq 0$. From $\gamma * \{\vartheta\} \notin C_*^{\mathrm{GM},\overline{p}}(\Delta^k * \underline{K})$, we deduce the compatibility of $\overline{\psi}$ with the differentials. This chain map $\overline{\psi}$ is clearly injective. It is also surjective because $C_*(\Delta^k) \cap C_*^{\mathrm{GM},\overline{p}}(\Delta^k * \underline{K}) = \{0\}$. This is a consequence of the fact that any γ in this intersection verifies $|\gamma| = \|\gamma\|_\ell \leq |\gamma| - \ell + \overline{p}(\ell) \leq |\gamma| - 2$, which is impossible.

Therefore, the map $\overline{\psi}$ is an isomorphism of complexes. The complex $C_*(\Delta^k)$ being contractible, we obtain an isomorphism

$$H_{*+1}(C^*_{\mathrm{GM},\overline{p}}(\Delta^k * \underline{K}, \underline{K})) \cong H_*(\hom(\tau_\ell^{\overline{p}} C_*(\underline{K}), R)).$$

For any cochain complex, (A, δ), $\delta(A^k) \subset A^{k+1}$, if we replace A by a truncation $\tau_j A$ defined by

$$(\tau_j A)^k = \left\{ \begin{array}{cl} 0 & \text{if } k < j, \\ A^j / \delta(A^{j-1}) & \text{if } k = j, \\ A^k & \text{if } k > j, \end{array} \right.$$

it is easy to see that $H^k(\tau_j A) = 0$ if $k < j$ and $H_k(\tau_j A) = H^k(A)$ if $k \geq j$. If the cochain complex A is the dual of a chain complex, C, the quotient by the image of δ corresponds to the dual of the cocycles in C. Thus, we have,

$$H_i(\hom(\tau_\ell^{\overline{p}} C_*(\underline{K}), R)) = \left\{ \begin{array}{cl} 0 & \text{if } i \leq \ell - 1 - \overline{p}(\ell), \\ H^i_{\mathrm{GM},\overline{p}}(\underline{K}; R) & \text{otherwise,} \end{array} \right.$$

which implies

$$H^i_{\mathrm{GM},\overline{p}}(\Delta^k * \underline{K}, \underline{K}; R) = \left\{ \begin{array}{cl} 0 & \text{if } i \leq \ell - 2 - \overline{p}(\ell), \\ H^{i-1}_{\mathrm{GM},\overline{p}}(\underline{K}; R)) & \text{otherwise.} \end{array} \right.$$

The result follows now from the fact that the map $\overline{\psi}$ induces the connecting map of the long exact sequence of the pair $(\Delta^k * \underline{K}, \underline{K})$. □

Theorem D. *Let $\underline{K}$ be a filtered face set, $\overline{p}$ and $\overline{q}$ be two perversities such that $\overline{q} \geq 0$ and $\overline{p}(i) + \overline{q}(i) = i - 2$ for any $i \in \{1, \ldots, n\}$. Let C^* be the universal local system of cochains with coefficients in a commutative ring R. The* GM-cochain complex *$C^*_{\mathrm{GM},\overline{p}}(\underline{K})$ and the blow-up $\widetilde{C}^*_{\overline{q}}(\underline{K})$ are related by a quasi-isomorphism, i.e., $H^*_{\mathrm{GM},\overline{p}}(\underline{K}; R) \cong H^*_{\overline{q}}(\underline{K}; \widetilde{C})$.*

Proof. First, we define $\chi\colon \widetilde{C}^*(\underline{K}) \to \hom(C_*^{\rm GM}(\underline{K}), R)$. If $\Phi \in \widetilde{C}^*(\underline{K})$, for any $\sigma\colon \Delta = \Delta^{j_0} * \cdots * \Delta^{j_n} \to \underline{K}$, we have $\Phi_\sigma = \Phi_{0,\sigma} \otimes \cdots \otimes \Phi_{n,\sigma} \in C^*(c\Delta^{j_0}) \otimes \cdots \otimes C^*(\Delta^{j_n})$. These cochains can be evaluated on the maximal simplex in each factor. We set

$$\chi(\Phi)(\sigma) = \Phi_{0,\sigma}([c\Delta^{j_0}]) \cdot \ldots \cdot \Phi_{n,\sigma}([\Delta^{j_n}])$$

and extend it by linearity. The compositions of χ with the differentials verify

$$\begin{aligned}
\chi(d\Phi)(\sigma) &= \sum_{i=0}^{n} \pm \Phi_{0,\sigma}([c\Delta^{j_0}]) \cdot \ldots \cdot d\Phi_{i,\sigma}([c\Delta^{j_i}]) \cdot \ldots \cdot \Phi_{n,\sigma}([\Delta^{j_n}]),\\
d\chi(\Phi)(\sigma) &= \chi(\Phi)(\partial\sigma)\\
&= \sum_{i=0}^{n} \pm \Phi_{0,\sigma}([c\Delta^{j_0}]) \cdot \ldots \cdot \Phi_{i,\sigma}([c\partial\Delta^{j_i}]) \cdot \ldots \cdot \Phi_{n,\sigma}([\Delta^{j_n}]).
\end{aligned}$$

From the definition of the differential of a cochain, we have $d\Phi_{i,\sigma}([c\Delta^{j_i}]) = \Phi_{i,\sigma}([\partial c\Delta^{j_i}])$. We observe also $\partial[c\Delta^{j_i}] = c\partial[\Delta^{j_i}] + [\Delta^{j_i} \times \{1\}]$. Therefore, to get the equality $d\chi(\Phi) = \chi(d\Phi)$, we have to prove the nullity of the products

$$\Pi_i = \Phi_{0,\sigma}([c\Delta^{j_0}]) \cdot \ldots \cdot \Phi_{i,\sigma}([\Delta^{j_i} \times \{1\}]) \cdot \ldots \cdot \Phi_{n,\sigma}([\Delta^{j_n}]),$$

for any $i \in \{0, \ldots, n-1\}$. For having $\Pi_i \neq 0$, if the restriction to $\Delta^{j_i} \times \{1\}$ is not 0, the perverse degrees of Φ must verify

$$\|\Phi\|_{n-i} = \dim(c\Delta^{j_{i+1}} \times \cdots \times c\Delta^{j_{n-1}} \times \Delta^{j_n}),$$

which is equivalent to

$$\|\Phi\|_{n-i} + \|\sigma\|_{n-i} = \dim\Delta - 1. \tag{8}$$

If $\Phi \in \widetilde{C}^*_{\overline{q}}(\underline{K})$ and $\sigma \in C_{*,\overline{p}}^{\rm GM}(\underline{K})$, their perverse degrees verify

$$\|\Phi\|_{n-i} \le \overline{q}(n-i) \text{ and } \|\sigma\|_{n-i} \le \dim\Delta - (n-i) + \overline{p}(n-i)$$

which imply

$$\begin{aligned}
\|\Phi\|_{n-i} + \|\sigma\|_{n-i} &\le \dim\Delta - (n-i) + \overline{p}(n-i) + \overline{q}(n-i)\\
&\le \dim\Delta - 2.
\end{aligned}$$

Thus the condition (8) cannot be satisfied and the products Π_i are equal to 0 for all i. We have proved the compatibility of χ with the differentials.

The fact that χ is a quasi-isomorphism follows from Proposition 6.3 Proposition 6.5, Proposition 6.6, Proposition 7.1 and Proposition 7.2. ☐

The first part of the next corollary is a direct consequence of Theorem D and Corollary 5.14. The second part follows from Proposition 3.4 and [23].

Corollary 7.3. *Let $R = \mathbb{Q}$. Let $\underline{K}$ be a filtered face set, $\overline{p}$ and $\overline{q}$ be perversities such that $\overline{q} \ge 0$ and $\overline{p}(i) + \overline{q}(i) = i - 2$ for any $i \in \{1, \ldots, n\}$. Then the complexes $\widetilde{A}_{PL,\overline{q}}(\underline{K})$, $\widetilde{C}^*_{\overline{q}}(\underline{K})$ and $C^*_{{\rm GM},\overline{p}}(\underline{K})$ are related by quasi-isomorphisms.*

In the particular case that $\overline{p}$ and $\overline{q}$ are perversities and $\underline{K}$ is the filtered face set associated to a locally conelike space, X, the homology of these complexes coincides with the original Goresky-MacPherson intersection cohomology of X.

Remark 7.4. If $\overline{p}\colon \{1,\dots,n\}\to\mathbb{Z}$ is a perversity such that $\overline{p}(1)<0$, we get,

$$\overline{p}(i)\leq\overline{p}(i-1)+1\leq\ldots\leq\overline{p}(1)+i-1<i-1,$$

for any $i\geq 1$. Thus the hypothesis $\overline{p}(\ell)\leq\ell-2$ of Proposition 7.2 and the hypothesis $\overline{q}\geq 0$ of Theorem D are satisfied.

In the case of a GM-perversity, with a similar argument, we get $\overline{p}(i)\leq\overline{p}(2)+i-2\leq i-2$, for any $i\geq 2$. But GM-perversities were performed for conelike spaces which are filtered spaces without strata of codimension 1 (see [16]) and the value $\overline{p}(1)$ is not considered in the notion of $\overline{p}$-admissibility. Thus, in this case, the previous hypotheses of Proposition 7.2 and of Theorem D are satisfied also.

Remark 7.5. The simplices in $\underline{K}\backslash\underline{K}_+$ are not considered in $\widetilde{C}^*_{\overline{q}}(\underline{K})$, by definition, and they do not appear in $C^*_{\mathrm{GM},\overline{p}}(\underline{K})$ if $\overline{q}(1)\geq 0$, or equivalently if $\overline{p}(1)\leq -1$. Indeed, any $\overline{p}$-admissible simplex $\sigma\colon\Delta=\Delta^{j_0}*\cdots*\Delta^{j_n}\to\underline{K}$ verifies

$$\|\sigma\|_1=\dim(\Delta^{j_0}*\cdots*\Delta^{j_{n-1}})\leq\dim\Delta-1+\overline{p}(1)\leq\dim\Delta-2$$

which implies $2\leq\dim\Delta-\dim(\Delta^{j_0}*\cdots*\Delta^{j_{n-1}})=j_n+1$ and $j_n\geq 1$.

8. Particular cases: $\overline{q}=\overline{0}$ and $\overline{q}=\overline{\infty}$

> We characterize the intersection cohomology in the cases $\overline{q}=\overline{0}$ and $\overline{q}=\overline{\infty}$. For $\overline{q}=\overline{0}$, we need to introduce the normalization of a filtered face set, as Goresky and MacPherson do in [16]. Here, the connectivity of the link is not sufficient, we need to use the expanded link.

Recall that the face set $\underline{K}^{[0]}$ is the regular part of a filtered face set, $\underline{K}$.

Proposition 8.1. *Let $\widetilde{F}$ be the blow-up of an extendable universal local system of differential coefficients F, over a commutative ring R and such that the homology of a point is reduced to R in degree 0. For any filtered face set, $\underline{K}$, the restriction map*

$$\widetilde{F}_{\overline{\infty}}(\underline{K})\to F(\underline{K}^{[0]})$$

is a quasi-isomorphism.

Proof. If $\omega\in\widetilde{F}_{\overline{\infty}}(\underline{K})$, we have an element $\omega_\sigma\in F(c\Delta^{j_0})\otimes\cdots\otimes F(c\Delta^{j_{n-1}})\otimes F(\Delta^{j_n})$, for each $\sigma\colon\Delta^{j_0}*\cdots*\Delta^{j_n}\to\underline{K}_+$. The simplices σ of $\underline{K}^{[0]}$ correspond to the particular case $j_i=-1$ for all $i\leq n-1$. A global section, still denoted $\omega\in F(\underline{K}^{[0]})$, is thus defined by restriction from $\underline{K}_+$ to $\underline{K}^{[0]}$.

We check easily that $\underline{K}\mapsto\widetilde{F}_{\overline{\infty}}(\underline{K})$ and $\underline{K}\mapsto F(\underline{K}^{[0]})$ are filtered theory of cochains, see Proposition 6.5. Let $\underline{K}$ be a filtered face set of depth $v(\underline{K})<n$ and such that $\widetilde{F}_{\overline{\infty}}(\underline{K})\to F(\underline{K}^{[0]})$ is a quasi-isomorphism. Using Proposition 6.3, we are reduced to prove that $\widetilde{F}_{\overline{\infty}}(\Delta^k*\underline{K})\to F(\Delta*\underline{K}^{[0]})$ is a quasi-isomorphism. On the right hand side, we have $(\Delta^k*\underline{K})^{[0]}=\underline{K}^{[0]}$. On the left hand side, with Proposition 6.6, we know that $H^*_{\overline{\infty}}(\Delta^k*\underline{K};\widetilde{F})\cong H^*_{\overline{\infty}}(\underline{K};\widetilde{F})$ and the statement is proved. □

For the study of the $\overline{0}$-cohomology, we introduce a concept similar to the normalization of Goresky and MacPherson in [16, Page 151]. We denote by $\sigma \lhd \phi$ the relation "σ is a face of ϕ".

Definition 8.2. A filtered face set, $\underline{K}$, is called *normal* if the two following conditions are satisfied.

(a) Any simplex $\sigma \in \underline{K}\backslash\underline{K}_+$ is a face of a simplex ϕ of $\underline{K}_+$, i.e., $\sigma \lhd \phi$.
(b) Moreover, the simplex $\phi \in \underline{K}_+$ is unique in the following sense.
Let $\sigma \in \underline{K}\backslash\underline{K}_+$. For any pair (ϕ, ϕ') of simplices of $\underline{K}_+$ such that $\sigma \lhd \phi$ and $\sigma \lhd \phi'$, there exists a family $\phi_1, \ldots, \phi_m$ of simplices of $\underline{K}_+$ such that $\sigma \lhd \phi_i$ for $i \in \{1, \ldots, m\}$ and $\phi \lhd \phi_1 \rhd \cdots \lhd \phi_m \rhd \phi'$.

Proposition 8.3. *Let $\underline{K}$ be a normal filtered face set. For any $r > 0$, $k \geq 0$ and $\tau \in \mathfrak{J}(\underline{K}^{[r],k})$, the expanded link, $\mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)$, is a connected, non empty and normal filtered face set.*

Proof. As $r > 0$, we have $\tau \in \underline{K}\backslash\underline{K}_+$ and there exists $\sigma \in \underline{K}_+$ such that $\tau \lhd \sigma$. We may suppose $R_1(\sigma) = \tau$. As $\sigma \neq \tau$, we have $(R_2(\sigma), \sigma) \in \mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)$ and $\mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau) \neq \emptyset$.

We prove now the connectivity of $\mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)$ by connecting any element $(R_2(\sigma'), \sigma') \in \mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)$ to $(R_2(\sigma), \sigma)$. We consider two cases.

- If $\sigma' \in \underline{K}_+$, as $R_1(\sigma') = R_1(\sigma) = \tau$, we may apply the unicity axiom of normal space. There exists a family $\phi_1, \ldots, \phi_m$ of simplices of $\underline{K}_+$ such that $\tau \lhd \phi_i$ for $i \in \{1, \ldots, m\}$ and $\sigma \lhd \phi_1 \rhd \cdots \lhd \phi_m \rhd \sigma'$. We may suppose $R_1(\phi_i) = \tau$ and, as $\phi_i \neq \tau$, we have $(R_2(\phi_i), \phi_i) \in \mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)$ for $i \in \{1, \ldots, m\}$. This implies the next relations in $\mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)$,
$$(R_2(\sigma), \sigma) \lhd (R_2(\phi_1), \phi_1) \rhd \cdots \lhd (R_2(\phi_m), \phi_m) \rhd (R_2(\sigma'), \sigma').$$
- If $\sigma' \notin \underline{K}_+$, there exists $\phi \in \underline{K}_+$ such that $\sigma' \lhd \phi$. As $R_1(\sigma') = \tau$, we may suppose $R_1(\phi) = \tau$ and we get $(R_2(\sigma'), \sigma') \lhd (R_2(\phi), \phi)$. By applying the first case to $(R_2(\phi), \phi) \in \mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)$, we can rely $(R_2(\sigma'), \sigma')$ to $(R_2(\sigma), \sigma)$.

We are reduced to the proof of the normality of $\mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)$. Let $(R_2(\sigma), \sigma) \in \mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)\backslash\mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)_+$. Then $\sigma \in \underline{K}\backslash\underline{K}_+$ and there exists $\phi \in \underline{K}_+$ with $\sigma \lhd \phi$. We may suppose $R_1(\phi) = \tau$ and we get $(R_2(\sigma), \sigma) \lhd (R_2(\phi), \phi)$ with $(R_2(\phi), \phi) \in \mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)$. This gives us the first property of a normal filtered face set. For the second one, consider $(R_2(\sigma), \sigma) \in \mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)\backslash\mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)_+$, $(R_2(\phi), \phi) \in \mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)_+$ and $(R_2(\phi'), \phi') \in \mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)_+$ such that
$$(R_2(\phi), \phi) \rhd (R_2(\sigma), \sigma) \lhd (R_2(\phi'), \phi').$$
We deduce ϕ and ϕ' in $\underline{K}_+$, $\sigma \in \underline{K}\backslash\underline{K}_+$, $R_1(\phi) = R_1(\phi') = R_1(\sigma) = \tau$ and $\phi \rhd \sigma \lhd \phi'$. From the unicity condition in a normal space, we deduce the existence of a family $\phi_1, \ldots, \phi_m$ of simplices of $\underline{K}_+$ such that $\sigma \lhd \phi_i$ for $i \in \{1, \ldots, m\}$ and $\phi \lhd \phi_1 \rhd \cdots \lhd \phi_m \rhd \phi'$. We may suppose $R_1(\phi_i) = \tau$ which gives $(R_2(\sigma), \sigma) \lhd (R_2(\phi_i), \phi_i)$ and $(R_2(\phi_i), \phi_i) \in \mathcal{L}^{\mathtt{exp}}(\underline{K}, \tau)$ for all $i \in \{1, \ldots, m\}$, and
$$(R_2(\phi), \phi) \lhd (R_2(\phi_1), \phi_1) \rhd \cdots \lhd (R_2(\phi_m), \phi_m) \rhd (R_2(\phi'), \phi').$$
□

Proposition 8.4. *Let $\widetilde{F}$ be the blow-up of an extendable universal local system of differential coefficients F, over a commutative ring R, such that the homology of a contractible face set is equal to R, concentrated in degree 0. For any normal filtered face set, $\underline{K}$, there exists a quasi-isomorphism*

$$F(\underline{K}) \to \widetilde{F}_{\overline{0}}(\underline{K}).$$

Proof. Let $\underline{K}$ be a normal filtered face set and $\omega \in F(\underline{K})$. For any simplex, $\sigma\colon \Delta = \Delta^{j_0} * \cdots * \Delta^{j_n} \to \underline{K}$, there exists $\omega_\sigma \in F(\Delta^{j_0} * \cdots * \Delta^{j_n})$. By pulling back ω_σ along the blow-up μ (see Definition 5.7), we get a global section on the blow-up of Δ and an element $\tilde{\omega}_\sigma \in F(c\Delta^{j_0}) \otimes \cdots \otimes F(c\Delta^{j_{n-1}}) \otimes F(\Delta^{j_n})$. These operations on forms are compatible with face operators and we obtain a cochain map $\rho\colon F(\underline{K}) \to \widetilde{F}_{\overline{0}}(\underline{K})$.

We have two filtered theories of cochains, $\underline{K} \mapsto \widetilde{F}_{\overline{0}}(\underline{K})$ and $\underline{K} \mapsto F(\underline{K})$. As $\underline{K}$ is normal, its expanded links are connected and we will use Proposition 6.4:
Let $\underline{L}$ be a connected filtered face set with $v(\underline{L}) < n$ and such that $F(\underline{L}) \to \widetilde{F}_{\overline{0}}(\underline{L})$ is a quasi-isomorphism. We are reduced to prove that $F(\Delta^k * \underline{L}) \to \widetilde{F}_{\overline{0}}(\Delta^k * \underline{L})$ is a quasi-isomorphism. On the left hand side, as the face set $\Delta^k * \underline{L}$ has the homotopy type of a point, the homology is reduced to R in degree 0. On the right hand side, with Proposition 6.6, we know that $H^*_{\overline{0}}(\Delta^k * \underline{L}; \widetilde{F}) \cong H^0_{\overline{0}}(\underline{L}; \widetilde{F})$ and the statement is proved grants to the hypothesis of connectivity of $\underline{L}$. □

Let $\overline{t}'$ be the loose perversity defined by $\overline{t}'(i) = i-2$, for any $i \in \{1, \ldots, n\}$. This perversity coincides with the top perversity $\overline{t}$ (Definition 2.5) for any $i \in \{2, \ldots, n\}$ and the difference between $\overline{t}$ and $\overline{t}'$ does not matter if we work with filtered spaces without strata of codimension 1, as it is the case in [16]. Proposition 8.4 and Theorem D give the characterization of the $\overline{t}'$-cohomology.

Corollary 8.5. *Let $\widetilde{C}$ be the blow-up of the universal local system of cochains, C, with coefficients in a commutative ring R. For any normal filtered face set, $\underline{K}$, we have an isomorphism*

$$H^*_{\overline{t}'}(\underline{K}; \widetilde{C}) \cong H^*_{\mathrm{GM},\overline{0}}(\underline{K}; R) \text{ and } H^*_{\mathrm{GM},\overline{t}'}(\underline{K}; R) \cong H^*(\underline{K}; R).$$

Definition 8.6. The *normalization of a filtered face set, $\underline{K}$,* is a map of filtered face sets,

$$\mathcal{N}\colon N(\underline{K}) \to \underline{K},$$

such that $N(\underline{K})$ is normal and the restriction map, $\mathcal{N}\colon N(\underline{K})_+ \to \underline{K}_+$, is an isomorphism.

Proposition 8.7. *Any filtered face set, $\underline{K}$, admits a unique normalization defined by*

$$N(\underline{K}) = \underline{K}_+ \cup \{\langle \sigma, \phi \rangle \mid \sigma \in \underline{K} \backslash \underline{K}_+,\ \phi \in \underline{K}_+ \text{ and } \sigma \lhd \phi\},$$

where

- *$\langle -, - \rangle$ denotes the equivalence class for the equivalence relation generated by*

$$(\sigma, \phi) \sim (\sigma, \phi') \iff \phi \lhd \phi',$$

- *the perverse degree of elements of* $\underline{K}_+$ *is keeped and* $\|\langle\sigma,\phi\rangle\|_i = \|\sigma\|_i$.

Moreover, if F *is an extendable universal local system of differential coefficients, over a commutative ring* R*, of blow-up* $\widetilde{F}$*, the map* $\mathcal{N}$ *induces*

- *an isomorphism* $H^*_{\overline{q}}(N(\underline{K});\widetilde{F}) \cong H^*_{\overline{q}}(\underline{K};\widetilde{F})$*, for any loose perversity* $\overline{q}$ *or* $\overline{q} = \overline{\infty}$,
- *an isomorphism* $H^{\mathrm{GM},\overline{p}}_*(N(\underline{K});R) \cong H^{\mathrm{GM},\overline{p}}_*(\underline{K};R)$*, for the Goresky-MacPherson homology if* $\overline{p}(1) < 0$.

Proof. Observe that, by construction, we have $N(\underline{K})_+ = \underline{K}_+$. Begin with the definition of a face operator, ∂_i. We consider three cases:

- if $\alpha \in \underline{K}_+$ is a simplex whose face in $\underline{K}$ owns to $\underline{K}_+$, we keep the same face operator than in $\underline{K}$,
- if $\alpha \in \underline{K}_+$ is a simplex whose face in $\underline{K}$ owns to $\underline{K}\backslash\underline{K}_+$, we set $\partial_i\alpha = \langle\partial_i\alpha,\alpha\rangle$, (in this case, ∂_i is the last face operator of α),
- we set $\partial_i\langle\sigma,\phi\rangle = \langle\partial_i\sigma,\phi\rangle$ otherwise.

Let $\alpha\colon \Delta = \Delta^{j_0} * \cdots * \Delta^{j_n} \to \underline{K}$ with $j = \dim\Delta$. For the commutation rule of face operators, we can reduce the verification to the next two cases:

- if $j_n = 0$ and $i < j$, we have
$$\partial_i\partial_j\alpha = \partial_i\langle\partial_j\alpha,\alpha\rangle = \langle\partial_i\partial_j\alpha,\alpha\rangle = \langle\partial_{j-1}\partial_i\alpha,\alpha\rangle = \partial_{j-1}\partial_i\alpha,$$
- if $j_n = 1$ and $i = j-1$, we have
$$\partial_i\partial_j\alpha = \langle\partial_i\partial_j\alpha,\partial_j\alpha\rangle = \langle\partial_{j-1}\partial_i\alpha,\alpha\rangle = \langle\partial_{j-1}\partial_i\alpha,\partial_i\alpha\rangle = \partial_{j-1}\partial_i\alpha.$$

We continue with the verification of the properties of a normal filtered face set. Let $\langle\sigma,\phi\rangle \in N(\underline{K})\backslash N(\underline{K})_+$ with $\phi\colon \Delta^{j_0} * \cdots * \Delta^{j_n} \to \underline{K}$. Let J be the smallest subset of indices such that $\partial_J\phi \notin \underline{K}_+$ and choose I such that $\partial_I\partial_J\phi = \sigma$. By definition of face operators, we have
$$\partial_I\partial_J\phi = \partial_I\langle\partial_J\phi,\phi\rangle = \langle\partial_I\partial_J\phi,\phi\rangle = \langle\sigma,\phi\rangle$$
and $\langle\sigma,\phi\rangle \lhd \phi$ which means that $\langle\sigma,\phi\rangle$ is a face of an element of $N(\underline{K})_+$. We study now the unicity property of the definition of normal filtered face set. Let $\langle\sigma,\phi\rangle \in N(\underline{K})\backslash N(\underline{K})_+$ and β, β' in $N(\underline{K})_+$ such that $\langle\sigma,\phi\rangle \lhd \beta$ and $\langle\sigma,\phi\rangle \lhd \beta'$. Let J be the smallest subset such that $\partial_J\beta \notin \underline{K}_+$ and I such that $\langle\sigma,\phi\rangle = \partial_I\partial_J\beta$. By definition of the face operator, we have
$$\langle\sigma,\phi\rangle = \partial_I\partial_J\beta = \partial_I\langle\partial_J\beta,\beta\rangle = \langle\partial_I\partial_J\beta,\beta\rangle,$$
which implies $\sigma = \partial_I\partial_J\beta$ (in $\underline{K}$!) and $\langle\sigma,\phi\rangle = \langle\sigma,\beta\rangle$. Similarly, we prove $\langle\sigma,\beta'\rangle = \langle\sigma,\phi\rangle$ and deduce
$$\langle\sigma,\phi\rangle = \langle\sigma,\beta\rangle = \langle\sigma,\beta'\rangle.$$
By definition of the equivalence relation, there exists a family of elements of $\underline{K}_+ = N(\underline{K})_+$, $(\beta_1,\ldots,\beta_p,\ldots,\beta_m)$, with
$$\beta \lhd \beta_1 \rhd \cdots \lhd \beta_p \rhd \phi \lhd \beta_{p+1} \rhd \cdots \lhd \beta_m \rhd \beta' \text{ and } \sigma \lhd \beta_i, \tag{9}$$
for all $i \in \{1,\ldots,m\}$. We are reduced to prove that $\langle\sigma,\phi\rangle \lhd \beta_i$, for all $i \in \{1,\ldots,m\}$. Observe that the relations (9) imply $\langle\sigma,\phi\rangle = \langle\sigma,\beta_i\rangle$, for all $i \in \{1,\ldots,m\}$. Let J be the smallest set of indices such that $\partial_J\beta_i \notin \underline{K}_+$ and I such that $\partial_I\partial_J\beta_i = \sigma$. By definition of the face operators, we get
$$\partial_I\partial_J\beta_i = \partial_I\langle\partial_J\beta_i,\beta_i\rangle = \langle\partial_I\partial_J\beta_i,\beta_i\rangle = \langle\sigma,\beta_i\rangle,$$

which implies $\langle\sigma,\beta_i\rangle \lhd \beta_i$. We have obtained $\langle\sigma,\phi\rangle = \langle\sigma,\beta_i\rangle \lhd \beta_i$.

A map of filtered face sets, $\mathcal{N}\colon N(\underline{K}) \to \underline{K}$, is defined by $\mathcal{N}(\alpha) = \alpha$, if $\alpha \in \underline{K}_+$ and $\mathcal{N}(\langle\sigma,\phi\rangle) = \sigma$. The verification of the compatibility with face operators is direct from the definitions and the restriction of $\mathcal{N}$ is the identity map from $N(\underline{K})_+$ to $\underline{K}_+$.

Suppose now that $\mathcal{N}\colon N(\underline{K}) \to \underline{K}$ and $\mathcal{N}'\colon N'(\underline{K}) \to \underline{K}$ are two normalizations of $\underline{K}$. We construct a map of filtered face sets, $\mathfrak{N}\colon N(\underline{K}) \to N'(\underline{K})$, such that $\mathcal{N}' \circ \mathfrak{N} = \mathcal{N}$, by:

$$\begin{array}{ll} \mathfrak{N}(\alpha) = \mathcal{N}'^{-1}(\mathcal{N}(\alpha)), & \text{if } \alpha \in N(\underline{K})_+ \\ \mathfrak{N}(\sigma) = \partial_I(\mathcal{N}'^{-1}(\mathcal{N}(\sigma))), & \text{if } \sigma = \partial_I\phi \in N(\underline{K})\backslash N(\underline{K})_+ \text{ and } \phi \in N(\underline{K})_+. \end{array}$$

We first have to prove that $\mathfrak{N}$ is well defined in the case $\sigma \notin N(\underline{K})_+$. Let $\phi' \in N(\underline{K})_+$ such that $\sigma = \partial_{I'}\phi'$. We may suppose that $\sigma \lhd \phi \lhd \phi'$ and set $\phi = \partial_J\phi'$. We have $\sigma = \partial_I\partial_J\phi'$ and

$$\partial_I(\mathcal{N}'^{-1}(\mathcal{N}(\phi)) = \partial_I(\mathcal{N}'^{-1}(\mathcal{N}(\partial_J\phi')) = \partial_I\partial_J(\mathcal{N}'^{-1}(\mathcal{N}(\phi'))),$$

which proves that $\mathfrak{N}$ is well defined.

We establish now the bijectivity of the map $\mathfrak{N}$. A map $\mathfrak{N}'\colon N'(\underline{K}) \to N(\underline{K})$ is defined in a similar manner and, on $N(\underline{K})_+$, they are obviously inverse. If $\sigma = \partial_I\phi$, we have

$$\mathfrak{N}'(\mathfrak{N}(\sigma)) = \mathfrak{N}'(\partial_I(\mathcal{N}'^{-1}(\mathcal{N}(\phi)))) = \partial_I(\mathcal{N}^{-1}(\mathcal{N}'(\mathcal{N}'^{-1}(\mathcal{N}(\phi))))) = \partial_I\phi = \sigma.$$

The map $\mathfrak{N}$ preserves the perverse degree. That is obvious on $N(\underline{K})_+$. Let $\sigma \in N(\underline{K})\backslash N(\underline{K})_+$. Then there exists $\phi \in N(\underline{K})_+$ and we may suppose $\sigma = \partial_m\phi$, with $m = |\phi|$. This implies

$$\|\mathfrak{N}(\sigma)\|_\ell = \|\partial_m\mathcal{N}'^{-1}(\mathcal{N}(\phi))\|_\ell = \left\{\begin{array}{ll} |\phi| - 1 = |\sigma| & \text{if } \ell = 0 \\ \|\phi\|_\ell = \|\sigma\|_\ell & \text{if } \ell > 0 \end{array}\right. = \|\sigma\|_\ell.$$

For proving the compatibility of $\mathfrak{N}$ with face operators, we consider three cases.

- It is direct if α and $\partial_i\alpha$ are in $N(\underline{K})_+$.
- Let $\alpha \in N(\underline{K})_+$ with $\partial_i\alpha \notin N(\underline{K})_+$. Then, we have
$$\partial_i\mathfrak{N}(\alpha) = \partial_i(\mathcal{N}'^{-1}(\mathcal{N}(\alpha))) = \mathfrak{N}(\partial_i\alpha).$$
- Let $\alpha \notin N(\underline{K})_+$. There exists $\phi \in N(\underline{K})_+$ with $\sigma = \partial_I\phi$ and we have
$$\partial_i\mathfrak{N}(\sigma) = \partial_i\partial_I(\mathcal{N}'^{-1}(\mathcal{N}(\phi))) = \mathfrak{N}(\partial_i\sigma).$$

The equality $\mathcal{N} \circ \mathfrak{N} = \mathcal{N}'$ is obvious on $N(\underline{K})_+$. If $\sigma = \partial_I\phi$, $\sigma \notin N(\underline{K})_+$ and $\phi \in N(\underline{K})_+$, we have

$$\mathcal{N}'(\mathfrak{N}(\sigma)) = \mathcal{N}'(\partial_I(\mathcal{N}'^{-1}(\mathcal{N}(\phi)))) = \partial_I\mathcal{N}(\phi) = \mathcal{N}(\partial_I\phi) = \mathcal{N}(\sigma).$$

We have established the existence and unicity of the normalization. We show now the existence of isomorphisms in cohomology and homology.

The restriction map of $\mathcal{N}$ to $N(\underline{K})_+$ being an isomorphism, $N(\underline{K})_+ \cong \underline{K}_+$, the map $\mathcal{N}$ induces an isomorphism for the cohomology with coefficients in $\widetilde{F}_{\overline{q}}$, for any loose perversity $\overline{q}$ or $\overline{q} = \overline{\infty}$.

Let $\sigma\colon \Delta = \Delta^{j_0} * \cdots * \Delta^{j_n} \to N(\underline{K})$ be a $\overline{p}$-admissible simplex for a perversity $\overline{p}$ such that $\overline{p}(1) < 0$. From $\|\sigma\|_1 = \dim(\Delta^{j_0} * \cdots * \Delta^{j_{n-1}})$ and

$$\|\sigma\|_1 \leq \dim\Delta - 1 + \overline{p}(1) \leq \dim\Delta - 2,$$

we deduce $j_n \geq 1$ and $\sigma \in \underline{K}_+ \cong N(\underline{K})_+$. This gives an isomorphism of chain complexes $C_*^{\mathrm{GM},\overline{p}}(N(\underline{K})) \cong C_*^{\mathrm{GM},\overline{p}}(\underline{K})$. □

Example 8.8. This example shows that the connectivity of $\mathcal{L}(\underline{K},\tau)$ is not sufficient for having a quasi-isomorphism between $F(\underline{K})$ and $\widetilde{F}_{\overline{0}}(\underline{K})$. Consider a face set, $\underline{K}$, which looks like a pinch ribbon. It is formed of two triangles, $\Delta^2_{(1)}$ and $\Delta^2_{(2)}$, with a common vertex and the opposite edge in common also. We can represent it as



with $[ad] = [be]$. We decompose the two triangles as $\{c\} * [ad]$ and $\{c\} * [be]$ creating a filtered face set still denoted $\underline{K}$. Its blow up is a rectangle and has no cohomology of degree 1 in perverse degree $\overline{0}$, i.e., $H^1_{\overline{0}}(\underline{K};\widetilde{C}) = 0$. The cohomology of the face set $\underline{K}$ has a generator in degree 1. It is easy to check that the link of $\{c\}$ is connected but its expanded link is not connected.

The normalization $N(\underline{K})$ of $\underline{K}$ is formed of $\underline{K}_+$ and two 0-simplices $\langle\{c\},\Delta^2_{(1)}\rangle$ and $\langle\{c\},\Delta^2_{(2)}\rangle$. For instance, we have $\partial_0[cd] = \langle\{c\},\Delta^2_{(1)}\rangle$. The determination of the other boundaries show that in $N(\underline{K})$ the two triangles still have a common edge but the two summits are now different.

Example 8.9. The cone of a face set $\underline{S}$ is the filtered face set $c\underline{S} = \{\vartheta\} * \underline{S} \in \mathbf{\Delta}^{[1]}_{\mathcal{F}}$. By definition, the strict perverse CDGA of the blow-up of Sullivan's forms is $A_{PL}(\Delta^1) \otimes A_{PL}(\underline{S}) = \wedge(t,dt) \otimes A_{PL}(\underline{S})$, with perverse degrees determined by $\|t\| = 0$, $\|dt\| = -\infty$ and $\|1 \otimes \omega\| = |\omega|$ if $\omega \in A_{PL}(\underline{S})$, $\omega \neq 0$. The perverse degree is extended to products by $\|\alpha_1\, \alpha_2\| = \|\alpha_1\| + \|\alpha_2\|$.

As there is only one stratum, a perversity is given by a non-negative number, $\overline{q}(1)$, and we have a quasi-isomorphism

$$\big(A_{PL}(\Delta^1) \otimes A_{PL}(\underline{S})\big)_{\overline{q}} \simeq A_{PL}^{<\overline{q}(1)}(\underline{S}) \oplus \mathcal{Z}A_{PL}^{\overline{q}(1)}(\underline{S}).$$

This means that *the strict perverse* CDGA, $\widetilde{A}_{PL}(c\underline{S})$, *is the* CDGA $A_{PL}(\underline{S})$ *with a perverse degree equal to the usual degree of forms.*

In the case of the filtered face set, $\underline{S}$, associated to $\mathbb{C}P(2)$, only three values suffice for the description of the various intersection cohomologies; they are $\overline{q}(1) = 0$, 2 and 4. More precisely, we have

$$\begin{aligned}
H^i_{\overline{0}}(c\mathbb{C}P(2);\mathbb{Q}) &= \begin{cases} \mathbb{Q} & \text{if } i = 0,\\ 0 & \text{if } i > 0.\end{cases}\\
H^i_{\overline{2}}(c\mathbb{C}P(2);\mathbb{Q}) &= \begin{cases} \mathbb{Q} & \text{if } i = 0,\ 2,\\ 0 & \text{otherwise.}\end{cases}\\
H^i_{\overline{4}}(c\mathbb{C}P(2);\mathbb{Q}) &= \begin{cases} \mathbb{Q} & \text{if } i = 0,\ 2,\ 4\\ 0 & \text{otherwise.}\end{cases}
\end{aligned}$$

In this example, we perform a computation of perverse cohomologies with filtrations that do not correspond to any notion of dimension and, as expected, the results coincide with the classical determination of the perverse cohomology of a cone. If we want to obtain these cohomologies with the same filtration, from the Goresky and MacPherson complexes, we need to choose perversities $\overline{p}$ connected to the previous $\overline{q}$'s by $\overline{p}+\overline{q}=-1$. We find: $H^*_{\mathrm{GM},-\overline{1}}(\mathbb{C}P(2);\mathbb{Q}) = H^*_{\overline{0}}(\mathbb{C}P(2);\mathbb{Q})$, $H^*_{\mathrm{GM},-\overline{3}}(\mathbb{C}P(2);\mathbb{Q}) = H^*_{\overline{2}}(\mathbb{C}P(2);\mathbb{Q})$ and $H^*_{\mathrm{GM},-\overline{5}}(\mathbb{C}P(2);\mathbb{Q}) = H^*_{\overline{4}}(\mathbb{C}P(2);\mathbb{Q})$.

Example 8.10. The suspension of the face set $\underline{S}$ is the filtered face set $\Sigma\underline{S} = \{\vartheta_1,\vartheta_2\} * \underline{S} \in \mathbf{\Delta}^{[1]}_{\mathcal{F}}$. By definition, the strict perverse CDGA of the blow-up of Sullivan's forms is $\widetilde{A}_{PL}(\Sigma\underline{S}) = \left(A_{PL}(\Delta^1)\oplus A_{PL}(\Delta^1)\right)_{t_1=t_2=0} \otimes A_{PL}(\underline{S})$, where $\left(A_{PL}(\Delta^1)\oplus A_{PL}(\Delta^1)\right)_{t_1=t_2=0}$ is the subspace of the direct sum generated by polynomials, $P(t_1,dt_1)+Q(t_2,dt_2)$, such that $P(0,0)=Q(0,0)$. As the polynomials coincide on one side of the intervals, we can replace the previous complex by $C = A_{PL}(\Delta^1)\otimes A_{PL}(\underline{S})$, with perverse degrees:

- if $f \in A^0_{PL}(\Delta^1)$, then $\|f\otimes\omega\| = -\infty$ if $f_{|\partial\Delta^1} = 0$ or $\omega = 0$ and $\|f\otimes\omega\| = |\omega|$ otherwise,
- if $\alpha\in A^1_{PL}(\Delta^1)$, then $\|\alpha\otimes\omega\| = -\infty$.

By definition of the elements of $\overline{q}$-intersection, we have,

$$C_{\overline{q}} = A_{PL}^{<\overline{q}(1)}(\underline{S}) \oplus \mathcal{Z}A_{PL}^{\overline{q}(1)} \oplus s(A_{PL}^{>\overline{q}(1)}(\underline{S})).$$

A verification shows that the cochain map, $\varphi\colon C_{\overline{q}} \to \widetilde{A}_{PL,\overline{q}}(\Sigma\underline{S})$, defined by $\varphi(\eta) = 1\otimes\eta$ if $|\eta|\leq\overline{q}(1)$ and $\varphi(\eta) = (dt,-dt)\otimes\eta$, if $\eta\in s(A_{PL}^{>\overline{q}(1)}(\underline{S}))$, is a quasi-isomorphism. In the case of $\mathbb{C}P(2)$, only three values suffice for the description of the various intersection cohomologies:

$$\begin{array}{rcl}
H^i_{\overline{0}}(\Sigma\mathbb{C}P(2);\mathbb{Q}) & = & \left\{\begin{array}{ll}\mathbb{Q} & \text{if } i=0,\ 3,\ 5,\\ 0 & \text{otherwise.}\end{array}\right.\\
H^i_{\overline{2}}(\Sigma\mathbb{C}P(2);\mathbb{Q}) & = & \left\{\begin{array}{ll}\mathbb{Q} & \text{if } i=0,\ 2,\ 5,\\ 0 & \text{otherwise.}\end{array}\right.\\
H^i_{\overline{4}}(\Sigma\mathbb{C}P(2);\mathbb{Q}) & = & \left\{\begin{array}{ll}\mathbb{Q} & \text{if } i=0,\ 2,\ 4,\\ 0 & \text{otherwise.}\end{array}\right.
\end{array}$$

In a similar computation done with GM-cochains, the perversities $\overline{0}$, $\overline{2}$ and $\overline{4}$ are replaced by $-\overline{1}$, $-\overline{3}$ and $-\overline{5}$, respectively.

9. Homotopy of filtered face sets

In this section, we introduce the notion of product of a filtered face set with a face set and use it for a definition of homotopy between filtered face sets maps. Kan fibrations are also defined in the category of filtered face sets maps and connected to their analogue in the category of face sets maps.

Let $\Delta^N = \Delta^{j_0}*\cdots*\Delta^{j_n}$ be a filtered simplex and Δ^k be a simplex. We describe a filtered face set $(\Delta^{j_0}*\cdots*\Delta^{j_n})\otimes\Delta^k$ by its vertices, as follows. We choose an order, $z_0<\cdots<z_k$, on the vertices of Δ^k and an order $a^0_\ell<\cdots<a^{j_\ell}_\ell$, on the vertices of Δ^{j_ℓ}, for any $\ell\in\{0,\ldots,n\}$.

The vertices of $(\Delta^{j_0}*\cdots*\Delta^{j_n})\otimes\Delta^k$ are the couples (a^s_ℓ, z_i) of vertices of Δ^N and Δ^k. The filtered simplices of the product are now defined by strictly increasing sequences of vertices, for the alphabetic order, the filtration degree being given by the index ℓ. For instance, the maximal simplices are the sequences of $(k+N+1)$ couples. We observe that this product is still of dimension n, as filtered face set, and that this construction is compatible with face operators. This justifies the next definition.

Definition 9.1. Let $\underline{K} \in \mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets$ be a filtered face set and $\underline{T}$ be a face set. The product of $\underline{K}$ and $\underline{T}$ is the filtered face set defined by

$$\underline{K} \otimes \underline{T} = \operatorname*{colim}_{\Delta^{j_0} * \cdots * \Delta^{j_n} \to \underline{K}} \left(\operatorname*{colim}_{\Delta^k \to \underline{T}} (\Delta^{j_0} * \cdots * \Delta^{j_n}) \otimes \Delta^k \right).$$

This product is clearly associative, i.e.,

$$\underline{K} \otimes (\underline{T} \otimes \underline{S}) = (\underline{K} \otimes \underline{T}) \otimes \underline{S},$$

extends naturally in a bifunctor,

$$\otimes \colon \mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets \times \mathbf{\Delta}-Sets \to \mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets,$$

and commutes with colimits by definition. For handling the product of two filtered face sets, one must consider filtered face sets of different dimensions, $[n]$; we do not need this generalization in the sequel.

We define now some structures on the sets of morphisms. We keep the previous notation: $\underline{K}$, $\underline{L}$ are filtered face sets and $\underline{T}$ is a face set. We define a filtered face set, $\underline{K}^{\underline{T}}$, by

$$(\underline{K}^{\underline{T}})_{j_0,\dots,j_n} = \hom_{\mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets}((\Delta^{j_0} \otimes \cdots \otimes \Delta^{j_n}) \otimes \underline{T}, \underline{K}),$$

and a face set, $\hom^{\mathbf{\Delta}}(\underline{K},\underline{L})$, by

$$\hom^{\mathbf{\Delta}}(\underline{K},\underline{L})_k = \hom_{\mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets}(\underline{K} \otimes \Delta^k, \underline{L}).$$

The following formulae of adjonctions derive directly from the definitions.

Proposition 9.2. *If $\underline{K}$, $\underline{L}$ are two filtered face sets and $\underline{T}$ is a face set, we have natural bijections,*

$$\begin{aligned} \hom_{\mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets}(\underline{K} \otimes \underline{T}, \underline{L}) &\cong \hom_{\mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets}(\underline{K}, \underline{L}^{\underline{T}}) \\ &\cong \hom_{\mathbf{\Delta}-Sets}(\underline{T}, \hom^{\mathbf{\Delta}}(\underline{K},\underline{L})). \end{aligned}$$

The two canonical maps, ι_0, $\iota_1 \colon \Delta^0 \to \Delta^1$, give a notion of homotopy in the category $\mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets$.

Definition 9.3. Let f, $g \colon \underline{K} \to \underline{L}$ be two maps of filtered face sets. They are *homotopic* if there exists a map of filtered face sets, $F \colon \underline{K} \otimes \Delta^1 \to \underline{L}$, such that $F \circ (\mathrm{id} \otimes \iota_0) = f$ and $F \circ (\mathrm{id} \otimes \iota_1) = g$. We denote this relation by $f \simeq g$.

In particular, the two injections ι_0, ι_1 are homotopic. As in the case of face sets, for getting an equivalence relation, we have to impose some restrictions. Let $m \geq 1$. We denote by $\Delta^{m,k}$ the face subset of $\partial\Delta^m$ obtained by removing the k-th $(m-1)$-face.

Definition 9.4. A map $f\colon \underline{K} \to \underline{K}'$ of filtered face sets is a *Kan fibration* if, for each filtered face set $\underline{L}$ and each commutative diagram of filtered face sets maps,

$$\begin{tikzcd}
\underline{L} \otimes \Delta^{m,k} \arrow[r, "\varphi"] \arrow[d] & \underline{K} \arrow[d, "f"] \\
\underline{L} \otimes \Delta^{m} \arrow[r, "\psi"] \arrow[ur, dashed, "g"] & \underline{K}'
\end{tikzcd}$$

there exists a filtered face sets map, $g\colon \underline{L} \otimes \Delta^m \to \underline{K}$, extending φ and lifting ψ. In the case $\underline{K}' = \underline{K}'_+ = \Delta^0$, the filtered face set $\underline{K}$ is called a *Kan filtered face set.*

This definition is connected to the notion of Kan fibration in the category of face sets maps, see [30].

Proposition 9.5. *A filtered face sets map, $f\colon \underline{K} \to \underline{K}'$, is Kan if, and only if, the corresponding face sets map, $\hat{f}\colon \hom^{\mathbf{\Delta}}(\underline{L},\underline{K}) \to \hom^{\mathbf{\Delta}}(\underline{L},\underline{K}')$, is a Kan fibration for any filtered face set $\underline{L}$.*

Proof. With Proposition 9.2, the existence of a filtered face sets map g, making commutative the following diagram,

$$\begin{tikzcd}
L \otimes \Delta^{m,k} \arrow[r, "\varphi"] \arrow[d] & \underline{K} \arrow[d, "f"] \\
L \otimes \Delta^{m} \arrow[r, "\psi"] \arrow[ur, dashed, "g"] & \underline{K}',
\end{tikzcd}$$

is equivalent to the existence of a face sets map, $\hat{g}$, making commutative the following diagram,

$$\begin{tikzcd}
\Delta^{m,k} \arrow[r, "\hat{\varphi}"] \arrow[d] & \hom^{\mathbf{\Delta}}(\underline{L},\underline{K}) \arrow[d, "\hat{f}"] \\
\Delta^{m} \arrow[r, "\hat{\psi}"] \arrow[ur, dashed, "\hat{g}"] & \hom^{\mathbf{\Delta}}(\underline{L},\underline{K}'),
\end{tikzcd}$$

where $\hat{\varphi}$, $\hat{\psi}$ correspond to φ, ψ in the adjonction, respectively. This last property is the definition of a Kan fibration in $\mathbf{\Delta}-Sets$, see [30]. □

Corollary 9.6. *Let $\underline{K}$ be a Kan filtered face set and $(\underline{T},\underline{S})$ be a pair of face sets. Then the map $\underline{K}^{\underline{T}} \to \underline{K}^{\underline{S}}$ is a Kan fibration in $\mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets$.*

Proof. Let $\underline{L}$ be a filtered face set. We have to find a dotted arrow making commutative the following diagram in $\mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets$,

$$\begin{tikzcd}
\underline{L} \otimes \Delta^{m,k} \arrow[r] \arrow[d] & \underline{K}^{\underline{T}} \arrow[d] \\
\underline{L} \otimes \Delta^{m} \arrow[r] \arrow[ur, dashed] & \underline{K}^{\underline{S}},
\end{tikzcd}$$

which is equivalent to the existence of a dotted arrow, in $\mathbf{\Delta}-Sets$, making commutative,

$$\begin{tikzcd}
\Delta^{m,k} \arrow[r] \arrow[d] & \hom^{\Delta}(\underline{L}, \underline{K}^{\underline{T}}) \arrow[d] \\
\Delta^{m} \arrow[r] \arrow[ur, dashed] & \hom^{\Delta}(\underline{L}, \underline{K}^{\underline{S}}).
\end{tikzcd}$$

This problem of existence can also be rewritten as the following lifting problem in $\mathbf{\Delta}-Sets$,

$$\begin{tikzcd}
\Delta^{m} \otimes \underline{S} \cup_{\Delta^{m,k}\otimes \underline{S}} \Delta^{m,k} \otimes \underline{T} \arrow[r] \arrow[d] & \hom^{\Delta}(\underline{L}, \underline{K}). \\
\Delta^{m} \otimes \underline{T} \arrow[ur, dashed] &
\end{tikzcd}$$

The face set $\hom^{\Delta}(\underline{L}, \underline{K})$ being Kan, by Proposition 9.5, this last dotted arrow exists. ☐

Two maps, $f, g \colon \underline{K} \to \underline{L}$ in $\mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets$, can be viewed as elements of $\hom^{\mathbf{\Delta}}(\underline{K}, \underline{L})_0$. From Definition 9.3, they are homotopic in $\mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets$ if, and only if, there exists $F \in \hom^{\mathbf{\Delta}}(\underline{K}, \underline{L})_1$ such that $\partial_0 F = f$ and $\partial_1 F = g$, where ∂_0, ∂_1 are face maps of the face set $\hom^{\mathbf{\Delta}}(\underline{K}, \underline{L})$. This observation implies the next result, by definition of homotopy groups of face sets.

Corollary 9.7. *If $\underline{L}$ is a Kan filtered face set, the relation of homotopy between maps of filtered face sets, of codomain $\underline{L}$, is an equivalence relation and the set of equivalence classes verifies*

$$[\underline{K}, \underline{L}]_{\mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets} = \pi_0 \hom^{\mathbf{\Delta}}(\underline{K}, \underline{L}).$$

We study now the behavior of the product of a filtered face set and a face set with intersection cohomology, in some particular cases which will be considered in Section 11.

Proposition 9.8. *Let $\underline{K}$ be a filtered face set and $f \colon \underline{X} \to \underline{Y}$ be a simplicial map between simplicial sets. Then, the map $\mathrm{id}\otimes f \colon \underline{K}\otimes\underline{X} \to \underline{K}\otimes\underline{Y}$ induces a chain map*

$$(\mathrm{id} \otimes f)_* \colon C_*^{\mathrm{GM},\overline{p}}(\underline{K} \otimes \underline{X}) \to C_*^{\mathrm{GM},\overline{p}}(\underline{K} \otimes \underline{Y}),$$

for any loose perversity $\overline{p}$. Moreover, if f is homotopic to g, then we have $H_*^{\mathrm{GM},\overline{p}}(\mathrm{id}\otimes f) = H_*^{\mathrm{GM},\overline{p}}(\mathrm{id}\otimes g)$. In particular, if f is a homotopy equivalence, the chain map $(\mathrm{id} \otimes f)_*$ is a quasi-isomorphism.

Proof. The map $\mathrm{id} \otimes f$ induces a chain map $C_*(\underline{K} \otimes \underline{X}) \to C_*(\underline{K} \otimes \underline{Y})$ and we have only to determine its behavior with the perverse degree. As it is a local computation, we suppose $f \colon \Delta^k \to \Delta^{k'}$ and consider the map $\mathrm{id} \otimes f \colon (\Delta^{j_0} * \cdots * \Delta^{j_n}) \otimes \Delta^k \to (\Delta^{j_0} * \cdots * \Delta^{j_n}) \otimes \Delta^{k'}$.

Let $\sigma \colon \nabla \to (\Delta^{j_0} * \cdots * \Delta^{j_n}) \otimes \Delta^k$ be a simplex. The simplex ∇ has a perverse degree induced by the perverse degree of a product defined at the beginning of this section. Thus, ∇ has a decomposition such that $\|\sigma\|_\ell = \dim\left(\nabla^{k_0} * \cdots * \nabla^{k_{n-\ell}}\right)$. Recall also that ∇ can be described by its vertices and these last ones are couples of a vertex of $\Delta^{j_0} * \cdots * \Delta^{j_n}$ and a vertex of

Δ^k. In this context, the integer $1+\|\sigma\|_\ell$ is the cardinal of the set of vertices whose first component belongs to $\Delta^{j_0}*\cdots*\Delta^{j_{n-\ell}}$. The image $((\mathrm{id}\otimes f)\circ\sigma)(\nabla)$ is a simplex whose vertices are the images of the vertices of ∇ by $\mathrm{id}\otimes f$. Thus the number of vertices whose first components are in $\Delta^{j_0}*\cdots*\Delta^{j_{n-\ell}}$ cannot increase and we get an induced map $C_*^{\mathrm{GM},\overline{p}}((\Delta^{j_0}*\cdots*\Delta^{j_n})\otimes\Delta^k)\to C_*^{\mathrm{GM},\overline{p}}((\Delta^{j_0}*\cdots*\Delta^{j_n})\otimes\Delta^{k'})$.

For the second part of the statement, it is sufficient to prove that the two injections ι_0, $\iota_1\colon \underline{K}\to\underline{K}\otimes\Delta^1$ induce the same map in homology. The argument used in the proof of Proposition 2.11 can be, word for word, adapted to this situation. □

10. Perverse differential graded algebras

> We collect here some results of Hovey [22] concerning the closed model structure on the category of perverse CDGA's and precise the notion of homotopy. A definition of minimal model is given.

With Example 8.10, it is clear that the cohomology of a strict perverse CDGA is not a strict perverse algebra. For the study of formality for instance, it is important that cochain complexes and their homology are objects of the same category. Therefore, we need to enlarge the notion of strict perverse CDGA's and we work with the perverse CDGA's introduced by M. Hovey, see [22]. Intuitively, starting from a strict perverse CDGA, A, we replace A by the data $A_\bullet$ of the $A_{\overline{p}}$ for the perversities $\overline{p}$, see Definitions 10.1 and 10.3. In fact, [22] is written in the case of Goresky-MacPherson perversities and we enlarge this setting by considering the lattice $\hat{\mathcal{P}}^n=\mathcal{P}^n\cup\{\infty\}$.

If $\overline{p}$ and $\overline{q}$ are elements of $\hat{\mathcal{P}}^n$, we define $\overline{p}\oplus\overline{q}$ as the smallest element, $\overline{r}$, of $\hat{\mathcal{P}}_n$ such that $\overline{p}+\overline{q}\leq\overline{r}$, see [16]. (In particular, if $\overline{p}(i)+\overline{q}(i)>i-2$ for some i, then $\overline{p}\oplus\overline{q}=\overline{\infty}$.) We may construct $\overline{p}\oplus\overline{q}$ by induction (see [12]), starting from $(\overline{p}\oplus\overline{q})(n)=\overline{p}(n)+\overline{q}(n)$. The inductive step is given by

$$\begin{array}{lll}\text{if } \overline{p}(k)+\overline{q}(k)<(\overline{p}\oplus\overline{q})(k+1), & \text{one sets} & (\overline{p}\oplus\overline{q})(k)=(\overline{p}\oplus\overline{q})(k+1)-1,\\ \text{if } \overline{p}(k)+\overline{q}(k)=(\overline{p}\oplus\overline{q})(k+1), & \text{one sets} & (\overline{p}\oplus\overline{q})(k)=(\overline{p}\oplus\overline{q})(k+1).\end{array}$$

This law is commutative, associative and has the null perversity as neutral element.

In the last sections of this work, we are only interested by algebraic structures defined on the field of rational numbers.

Definition 10.1. A *perverse vector space* is a functor from $\hat{\mathcal{P}}_n$ to the category of $\mathbb{Q}$-vector spaces, $\overline{p}\mapsto M_{\overline{p}}$. *Perverse linear maps* are natural transformations between perverse vector spaces.

As in [22], we observe that a perverse vector space, M, is *projective* if the map $\mathrm{colim}_{\overline{q}<\overline{p}}M_{\overline{q}}\to M_{\overline{p}}$ is injective and we define a tensor product of perverse vector spaces by

$$(M\otimes N)_{\overline{r}}=\operatorname*{colim}_{\overline{p}+\overline{q}\leq\overline{r}}M_{\overline{p}}\otimes_{\mathbb{Q}}N_{\overline{q}}. \tag{10}$$

If V is a $\mathbb{Q}$-vector space and $\overline{p}\in\hat{\mathcal{P}}_n$, we define a perverse vector space $V_{[\overline{p}]}$ by $(V_{[\overline{p}]})_{\overline{r}}=V$ if $\overline{r}\geq\overline{p}$ and 0 otherwise. The projective perverse vector spaces are direct factors of sums of $V_{[\overline{p}]}$, in which the vector space V and the perversity $\overline{p}$ vary.

Denote by $\mathrm{Ch}(\hat{\mathcal{P}}_n)$ the category of chain complexes, $\mathbb{Z}$-graded, formed of perverse graded vector spaces provided with a differential; we call them *perverse chain complexes.* A *perverse chain map* is a perverse linear map compatible with the differentials. The projective elements are the perverse chain complexes formed of projective perverse vector spaces, A, so that each perverse chain map, $A \to B$, into an exact complex is chain homotopic to zero. In particular, bounded below perverse chain complexes formed of projective perverse vector spaces are projective.

Theorem 10.2 (cf. [7], [22]). *The category* $\mathrm{Ch}(\hat{\mathcal{P}}_n)$ *is a closed model category for the following classes of objects.*

- *The* weak equivalences *are the perverse chain maps* $f\colon A \to B$ *such that*
$$H_*(f)(\overline{p})\colon H_*(A_{\overline{p}}) \to H_*(B_{\overline{p}})$$
is an isomorphism for any $\overline{p} \in \hat{\mathcal{P}}_n$.
- *The* fibrations *are the surjections.*
- *The* cofibrations *are the injective maps with a projective cokernel in* $\mathrm{Ch}(\hat{\mathcal{P}}_n)$.

Moreover, this structure is monoidal.

In particular, all objects of $\mathrm{Ch}(\hat{\mathcal{P}}_n)$ are fibrant.

Definition 10.3. A *perverse* DGA is a monoid in the category $\mathrm{Ch}(\hat{\mathcal{P}}_n)$. We denote by $\mathrm{CDGA}_{\mathcal{F}}$ the category of perverse CDGA's.

To any strict perverse CDGA, A, we associate a perverse CDGA defined by
$$A_{\overline{p}} = \{\omega \mid \|\omega\|_i \leq \overline{p}(i) \text{ and } \|d\omega\|_i \leq \overline{p}(i)\}.$$

The free preverse DGA on a perverse module X is defined by $T(X) = \oplus_{k\geq 0} X^{\otimes k}$, where the tensor product is defined in (10). The commutative free perverse CDGA on a perverse module X is the quotient $\wedge X := T(X)/\Sigma_n$ of the free perverse DGA by the action of the symmetric group, i.e.,
$$x_0\otimes\cdots\otimes x_i\otimes\cdots\otimes x_j\otimes\cdots\otimes x_n \sim (-1)^{|x_i|\,|x_j|}x_0\otimes\cdots\otimes x_j\otimes\cdots\otimes x_i\otimes\cdots\otimes x_n.$$

Remark 10.4. Two strict perverse CDGA's can be isomorphic as perverse CDGA's but not as strict perverse CDGA's, as

- $A = \wedge(e,x)$ with $|e| = \|e\| = 2$, $|x| = 3$, $\|x\| = 4$ and $dx = e^2$,
- $B = \wedge(e,y)$ with $|e| = \|e\| = 2$, $|y| = \|y\| = 3$, and $dy = e^2$.

Theorem 10.5 (cf. [22]). *The category* $\mathrm{CDGA}_{\mathcal{F}}$ *is a closed model category for the following classes of objects.*

- *The* weak equivalences *are the morphisms which are weak-equivalences in* $\mathrm{Ch}(\hat{\mathcal{P}}_n)$.
- *The* fibrations *are the surjections.*
- *The* cofibrations *are defined by the lifting property relatively to trivial fibrations.*

In the free perverse CGA, $B = \wedge \oplus_{\overline{p}} V_{[\overline{p}]}$, the elements $x \in V_{[\overline{p}]}$ have a cohomological degree, denoted by $|x|$, and a perverse degree denoted by $\|x\| = \overline{p}$. This degree is extended to the free CGA by $\|x_1 x_2\| = \|x_1\| + \|x_2\|$. We denote by $B_{[\overline{p}]}$ the vector space of elements of perverse graduation $\overline{p}$ in B and by $B_{\overline{p}} = \oplus_{\overline{r} \leq \overline{p}} B_{[\overline{r}]}$.

Definition 10.6. A perverse CDGA $(\wedge \oplus_{\overline{p}} V_{[\overline{p}]}, d)$ is a *Sullivan algebra* if it is free as perverse CGA, and if there exists a basis $(x_i)_{i \in I}$ of $\oplus_{\overline{p}} V_{[\overline{p}]}$ such that

(i) the set I is well-ordered, i.e., any non-empty subset has a smaller element,
(ii) the inequality $\|x_\alpha\| < \|x_\beta\|$ implies $\alpha < \beta$,
(iii) the differential of x_α belongs to $\wedge x_{<\alpha}$, where $\wedge x_{<\alpha}$ denotes the sub-CDGA of $\wedge \oplus_{\overline{p}} V_{[\overline{p}]}$ generated by the elements $(x_\beta)_{\beta < \alpha}$ of the basis.

A *Sullivan algebra is minimal* if, for all perversities $\overline{p}$ or $\overline{p} = \overline{\infty}$, we have

$$d(V_{[\overline{p}]}) \subset \left(\wedge \oplus_{\overline{r} < \overline{p}} V_{[\overline{r}]}\right)^{+} \otimes \wedge V_{[\overline{p}]} \oplus \left(\wedge \oplus_{\overline{r} < \overline{p}} V_{[\overline{r}]}\right) \otimes \wedge^{\geq 2} V_{[\overline{p}]}.$$

If $A_\bullet$ is a perverse CDGA, a *Sullivan model* of $A_\bullet$ is a quasi-isomorphism of $\mathrm{CDGA}_{\mathcal{F}}$, $\rho \colon (\wedge \oplus_{\overline{p}} V_{[\overline{p}]}, d) \to A_\bullet$, whose domain is a Sullivan algebra. If the domain is also minimal, we say that ρ is a *minimal model* of $A_\bullet$.

Observe that, in a Sullivan algebra, $(\wedge \oplus_{\overline{p}} V_{[\overline{p}]}, d)$, the differential decreases the perverse degree.

Proposition 10.7. *Sullivan perverse algebras are cofibrant in* $\mathrm{CDGA}_{\mathcal{F}}$.

Proof. Let $(\wedge \oplus_{\overline{p}} V_{[\overline{p}]}, d)$ be a Sullivan algebra. We consider the following diagram in $\mathrm{CDGA}_{\mathcal{F}}$

$$\begin{tikzcd} & E_\bullet \arrow[d, "\rho", "\sim"'] \\ (\wedge \oplus_{\overline{p}} V_{[\overline{p}]}, d)_\bullet \arrow[ur, dashed, "g"] \arrow[r, "f"'] & B_\bullet, \end{tikzcd}$$

where ρ is a surjective weak equivalence and the morphism $g \in \mathrm{CDGA}_{\mathcal{F}}$ has to be constructed such that $\rho \circ g = f$. On $\wedge V_{\overline{0}}$, that is the usual lifting lemma for CDGA's. We consider a basis $(x_\alpha)_{\alpha \in I}$ of $\oplus_{\overline{p}} V_{[\overline{p}]}$.

Consider x_α with $\|x_\alpha\| = \overline{p}$ and suppose that g is defined on $\wedge x_{<\alpha}$. This implies $\rho(g(dx_\alpha)) = f(dx_\alpha) = df(x_\alpha)$. We choose $y_\alpha \in E_{\overline{p}}$ such that $\rho(y_\alpha) = f(x_\alpha)$. We have

$$\left\{ \begin{array}{lclcl} \rho(g(dx_\alpha) - dy_\alpha) & = & df(x_\alpha) - df(x_\alpha) & = & 0 \\ d(g(dx_\alpha) - dy_\alpha) & = & g(d^2 x_\alpha) - d^2 y_\alpha & = & 0. \end{array} \right.$$

As $g(dx_\alpha) - dy_\alpha$ is a cocycle of $\operatorname{Ker} \rho$, by acyclicity, there exists $z_\alpha \in (\operatorname{Ker} \rho)_{\overline{p}}$ with $dz_\alpha = g(dx_\alpha) - dy_\alpha$. We set $g(x_\alpha) = y_\alpha + z_\alpha$ which satisfies

$$\left\{ \begin{array}{lclcl} \rho(g(x_\alpha)) & = & \rho(y_\alpha) + \rho(z_\alpha) & = & f(x_\alpha) \\ \rho(g(dx_\alpha)) & = & \rho(dy_\alpha) + \rho(dz_\alpha) & = & d\rho(y_\alpha) \\ & = & df(x_\alpha) & = & d\rho(g(x_\alpha)). \end{array} \right.$$

$\square$

As in the classical case ([35, Theorem 5.1]), minimal models are unique up to isomorphisms.

Proposition 10.8. *Any quasi-isomorphism between Sullivan minimal perverse algebras is an isomorphism.*

Proof. Let $\varphi\colon B=(\wedge\oplus_{\overline{p}}V_{[\overline{p}]},d)\to C=(\wedge\oplus_{\overline{p}}W_{[\overline{p}]},d)$ be a quasi-isomorphism between two Sullivan minimal algebras. We adapt to the perverse setting a proof made by Antonio Gómez-Tato in [14], by building a map $\psi\colon C\to B$ such that $\varphi\circ\psi=\mathrm{id}$.

First, we observe that the restriction $\varphi\colon(\wedge V_{[\overline{0}]},d)\to(\wedge W_{[\overline{0}]},d)$ is a quasi-isomorphism between classical minimal models therefore it is an isomorphism and the map ψ is defined on $\wedge W_{[\overline{0}]}$. Let $\overline{p}$ be a perversity. The conditions on the basis and the minimality property imply that there exists a filtration of $W^n_{[\overline{p}]}$ as $W^n_{[\overline{p}]}(i)=W^n_{[\overline{p}]}(i-1)\oplus W^n_{[\overline{p}]}[i]$ such that

$$d(W^n_{[\overline{p}]}[i])\subset\left(\wedge\oplus_{\overline{r}<\overline{p}}W_{[\overline{r}]}\right)\otimes\wedge W^{\leq n}_{[\overline{p}]}(i-1).$$

Set $M=\left(\wedge\oplus_{\overline{r}<\overline{p}}W_{[\overline{r}]}\right)\otimes\wedge W^{\leq n}_{[\overline{p}]}(i-1)$. We suppose that ψ is defined on M and the construction of ψ on C is reduced to its extension to $W^n_{[\overline{p}]}[i]$. With the 5-lemma, we get a quasi-isomorphism

$$\overline{\varphi}\colon\;(B/\psi(M))_{\overline{p}}\to(C/M)_{\overline{p}}.$$

Let (w_i) be a basis of $W^n_{[\overline{p}]}[i]$. Any w_i is a cocycle in $(C/M)_{\overline{p}}$ which cannot be a coboundary. We deduce the existence of $v_i\in B_{\overline{p}}$, $m_i\in M_{\overline{p}}$, $m'_i\in M_{\overline{p}}$ such that

$$\varphi(v_i)=w_i+m_i \text{ and } dv_i=\psi(m'_i).$$

We set $\psi(w_i)=v_i-\psi(m_i)$ and we check:

$$\begin{array}{rcl} dw_i &=& d\varphi(v_i)-dm_i=\varphi(\psi(m'_i))-dm_i=m'_i-dm_i,\\ \psi(dw_i) &=& \psi(m'_i)-\psi(dm_i)=dv_i-d\psi(m_i)=d\psi(w_i),\\ \varphi(\psi(w_i)) &=& \varphi(v_i)-\varphi(\psi(m_i))=w_i. \end{array}$$

From the equality $\varphi\circ\psi=\mathrm{id}$, we deduce that ψ is a quasi-isomorphism as well and the same construction applied to ψ brings a morphism φ' such that $\psi\circ\varphi'=\mathrm{id}$. Therefore ψ is an isomorphism and $\varphi=\psi^{-1}$ is one also. □

Definition 10.9. A *path object* of an element $B_\bullet$ of $\mathrm{CDGA}_{\mathcal{F}}$ is given by

$$(B\otimes\wedge(t,dt))_\bullet \underset{\pi_1}{\overset{\pi_0}{\rightrightarrows}} B_\bullet$$

where $|t|=0$, $|dt|=1$, $(B\otimes\wedge(t,dt))_{\overline{p}}=B_{\overline{p}}\otimes\wedge(t,dt)$ and the maps π_i are defined by the identity on $B_\bullet$ and the evaluations $\pi_i(t)=i$, for $i=0,1$.

Two morphisms, f, $g\colon A_\bullet\to B_\bullet$ of $\mathrm{CDGA}_{\mathcal{F}}$, are *homotopic* if there exists a diagram in $\mathrm{CDGA}_{\mathcal{F}}$,

$$\begin{tikzcd} A_\bullet \arrow[r, shift left, "f"] \arrow[r, shift right, "g"'] \arrow[dr, "\varphi"'] & B_\bullet \\ & (B\otimes\wedge(t,dt))_\bullet \arrow[u, shift left, "\pi_0"] \arrow[u, shift right, "\pi_1"'] \end{tikzcd}$$

such that $\pi_0\circ\varphi=f$ and $\pi_1\circ\varphi=g$.

The proofs of the following properties of homotopy are an easy extension of the classical case.

Proposition 10.10. *Let $A'_\bullet$ and $A_\bullet$ be cofibrant objects of* $\mathrm{CDGA}_{\mathcal{F}}$.

(1) *If*

$$\begin{tikzcd} A'_\bullet \arrow[r, "g"] & A_\bullet \arrow[r, shift left, "f_0"] \arrow[r, shift right, "f_1"'] & B_\bullet \arrow[r, "h"] & B'_\bullet \end{tikzcd}$$

is a diagram of morphisms of $\mathrm{CDGA}_{\mathcal{F}}$*, the next properties are satisfied.*

- (a) *If f_0 is homotopic to f_1, then $h \circ f_0 \circ g$ is homotopic to $h \circ f_1 \circ g$.*
- (b) *Homotopy is an equivalence relation on the set of morphisms from $A_\bullet$ to $B_\bullet$. We denote this relation by $\simeq$.*

(2) *Any weak equivalence, $h\colon B_\bullet \to B'_\bullet$ in* $\mathrm{CDGA}_{\mathcal{F}}$*, induces an isomorphism between the homotopy classes,*

$$h_\sharp\colon [A_\bullet, B_\bullet] \to [A_\bullet, B'_\bullet].$$

Definition 10.11. Let f_0, $f_1\colon B_\bullet \to B'_\bullet$ be two morphisms in $\mathrm{CDGA}_{\mathcal{F}}$ and $\varphi\colon A_\bullet \to B_\bullet$ be a cofibrant model. The morphisms f_0 and f_1 are *homotopic* if $f_0 \circ \varphi \simeq f_1 \circ \varphi$.

Proposition 10.10 implies that this definition is independant of the choice of the model, φ, of $B_\bullet$.

11. FILTERED FACE SETS AND PERVERSE DIFFERENTIAL GRADED ALGEBRAS

We construct two contravariant functors between the category of perverse CDGA's and the category of filtered face sets. We prove that this adjunction is compatible with homotopy classes (Theorem E) and establish the representability of the rational intersection cohomology of filtered face sets.

Definition 11.1. The *filtered realization* of a perverse CDGA, $A_\bullet$, is the filtered face set $\langle A_\bullet \rangle$, defined by

$$\langle A_\bullet \rangle_{(j_0,\ldots,j_n)} = \hom_{\mathrm{CDGA}_{\mathcal{F}}}(A_\bullet, \widetilde{A}_{PL,\bullet}(\Delta^{j_0} * \cdots * \Delta^{j_n})).$$

Let $\underline{K}$ be a filtered face set. The blow-up of Sullivan's forms gives a perverse CDGA, $\mathcal{A}(\underline{K})_\bullet$, defined by

$$\mathcal{A}(\underline{K})_{\overline{q}} = \widetilde{A}_{PL,\overline{q}}(\underline{K}).$$

The constructions $\mathcal{A}$ and $\langle - \rangle$ are extended in functors which are adjoint in the following sense.

Proposition 11.2. *The filtered realization and the blow-up of Sullivan's forms give a pair of contravariant adjoint functors, $\mathcal{A}\colon \mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets \to$* $\mathrm{CDGA}_{\mathcal{F}}$ *and $\langle - \rangle\colon$* $\mathrm{CDGA}_{\mathcal{F}} \to \mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets$*, i.e., for any $\underline{K} \in \mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets$ and any $A_\bullet \in$* $\mathrm{CDGA}_{\mathcal{F}}$*, there is a bijection*

$$\hom_{\mathbf{\Delta}^{[n]}_{\mathcal{F}}-Sets}(\underline{K}, \langle A_\bullet \rangle) \cong \hom_{\mathrm{CDGA}_{\mathcal{F}}}(A_\bullet, \mathcal{A}(\underline{K})_\bullet).$$

Proof. By definition of the functors and Yoneda lemma, we have

$$\begin{aligned}\hom_{\Delta^{[n]}_{\mathcal{F}}-Sets}(\underline{K},\langle A_\bullet\rangle) &= \hom_{\Delta^{[n]}_{\mathcal{F}}-Sets}(\operatorname*{colim}_{\underline{K}}\Delta^{j_0}*\cdots*\Delta^{j_n},\langle A_\bullet\rangle)\\ &= \lim_{\underline{K}}\hom_{\Delta^{[n]}_{\mathcal{F}}-Sets}(\Delta^{j_0}*\cdots*\Delta^{j_n},\langle A_\bullet\rangle)\\ &= \lim_{\underline{K}}\langle A_\bullet\rangle_{(j_0,\dots,j_n)}\end{aligned}$$

and

$$\begin{aligned}\hom_{\mathrm{CDGA}_{\mathcal{F}}}(A_\bullet,\mathcal{A}(\underline{K})_\bullet) &= \hom_{\mathrm{CDGA}_{\mathcal{F}}}(A_\bullet,(\widetilde{A}_{PL,\bullet}(\operatorname*{colim}_{\underline{K}}\Delta^{j_0}*\cdots*\Delta^{j_n})))\\ &= \hom_{\mathrm{CDGA}_{\mathcal{F}}}(A_\bullet,\lim_{\underline{K}}(\widetilde{A}_{PL,\bullet}(\Delta^{j_0}*\cdots*\Delta^{j_n})))\\ &= \lim_{\underline{K}}\hom_{\mathrm{CDGA}_{\mathcal{F}}}(A_\bullet,\widetilde{A}_{PL,\bullet}(\Delta^{j_0}*\cdots*\Delta^{j_n}))\\ &= \lim_{\underline{K}}\langle A_\bullet\rangle_{(j_0,\dots,j_n)}.\end{aligned}$$

□

The proof of the next theorem is postpone at the end of this section.

Theorem E. *Let $A_\bullet$ be a cofibrant perverse* CDGA*. Then the isomorphisms of Proposition 11.2 induce a bijection between homotopy classes, i.e.,*

$$[\underline{K},\langle A_\bullet\rangle]_{\Delta^{[n]}_{\mathcal{F}}-Sets}\cong[A_\bullet,\mathcal{A}(\underline{K})_\bullet]_{\mathrm{CDGA}_{\mathcal{F}}}.$$

Denote by $S^k_{\overline{p}}$ the free perverse vector space of dimension 1, with a generator of degree k in perverse degree $\overline{p}\in\hat{\mathcal{P}}_n$. If $\underline{K}$ is a filtered face set, Theorem E implies

$$\begin{aligned}H^k_{\overline{p}}(\underline{K};\mathbb{Q}) &= [S^k_{\overline{p}},\mathcal{A}(\underline{K})_\bullet]_{\mathrm{Ch}(\hat{\mathcal{P}}_n)}\\ &\cong [(\wedge S^k_{\overline{p}})_\bullet,\mathcal{A}(\underline{K})_\bullet]_{\mathrm{CDGA}_{\mathcal{F}}}\\ &\cong [\underline{K},\langle(\wedge S^k_{\overline{p}})_\bullet\rangle]_{\Delta^{[n]}_{\mathcal{F}}-Sets}.\end{aligned}$$

This proves the following statement which extends the representability of the usual cohomology of face sets.

Corollary 11.3. *The functor $\underline{K}\mapsto H^k_{\overline{p}}(\underline{K};\mathbb{Q})$ is representable; i.e., for any filtered face set $\underline{K}$ and any loose perversity $\overline{p}\in\hat{\mathcal{P}}_n$, we have an isomorphism,*

$$H^n(\underline{K};\mathbb{Q})\cong[\underline{K},\langle(\wedge S^k_{\overline{p}})_\bullet\rangle]_{\Delta^{[n]}_{\mathcal{F}}-Sets}.$$

Definition 11.4. The filtered face set $\langle(\wedge S^k_{\overline{p}})_\bullet\rangle$ is called the *Eilenberg-MacLane perverse filtered face set* and denoted $K(\mathbb{Q},k,\overline{p})$.

The set of filtered simplices of $K(\mathbb{Q},k,\overline{p})$ coincides with a set of cocycles of the Sullivan's forms on a blow-up. More precisely, we have:

$$\begin{aligned}K(\mathbb{Q},k,\overline{p})_{(j_0,\dots,j_n)} &= \hom_{\mathrm{CDGA}_{\mathcal{F}}}(((\langle\wedge S^k_{\overline{p}}\rangle)_\bullet,\widetilde{A}_{PL,\bullet}(\Delta^{j_0}*\cdots*\Delta^{j_n}))\\ &\cong \hom_{\mathrm{Ch}(\hat{\mathcal{P}}_n)}(S^k_{\overline{p}},\widetilde{A}_{PL,\bullet}(\Delta^{j_0}*\cdots*\Delta^{j_n}))\\ &= \mathcal{Z}\widetilde{A}_{PL,\overline{p}}(\Delta^{j_0}*\cdots*\Delta^{j_n}).\end{aligned}$$

We state and prove now three lemmas that we need for the proof of Theorem E. We denote by $\Delta^{m,k}$ the face subset of the boundary $\partial\Delta^m$ obtained by removing the k-th $(m-1)$-face.

Lemma 11.5. *Let* $\underline{K} \in \mathbf{\Delta}^{[n]}_{\mathcal{F}}$*−Sets. For any* $m \geq 1$ *and any* k*,* $0 \leq k \leq m+1$*, the canonical inclusion* $\Delta^{m,k} \to \Delta^m$ *induces a trivial fibration*

$$\varphi\colon \mathcal{A}(\underline{K} \otimes \Delta^m)_\bullet \to \mathcal{A}(\underline{K} \otimes \Delta^{m,k})_\bullet.$$

Proof. The fact that φ is a quasi-isomorphism is a direct consequence of Proposition 9.8 and Corollary 7.3. We are reduced to prove the surjectivity. We proceed in two steps.

- We establish first the surjectivity of $\mathcal{A}(\underline{K} \otimes \Delta^m)_\bullet \to \mathcal{A}(\underline{K} \otimes \partial\Delta^m)_\bullet$. Any simplex $\sigma \in \underline{K} \otimes \Delta^m \backslash \underline{K} \otimes \partial\Delta^m$ owns a face $\sigma' \in \underline{K} \otimes \partial\Delta^m$, i.e.,

$$\begin{tikzcd}
\Delta^{j_0} * \cdots * (\Delta^{j_i} * \{v\}) * \cdots * \Delta^{j_n} \arrow[r, "\sigma"] & \underline{K} \otimes \Delta^m \\
\Delta^{j_0} * \cdots * \Delta^{j_i} * \cdots * \Delta^{j_n}. \arrow[u] \arrow[ur, "\sigma'"'] &
\end{tikzcd}$$

  The surjectivity of the map induced by any face operator, $A_{PL}(c(\Delta^{j_i} * \{v\})) \to A_{PL}(c\Delta^{j_i})$, implies the surjectivity of the cochain map $A_{PL}(\underline{K} \otimes \Delta^m) \to A_{PL}(\underline{K} \otimes \partial\Delta^m)$. By definition of the perverse degree, this surjectivity inherits to the restriction $A_{PL,\overline{q}}(\underline{K} \otimes \Delta^m) \to A_{PL,\overline{q}}(\underline{K} \otimes \partial\Delta^m)$, for any $\overline{q} \in \hat{\mathcal{P}}^n$.
- Then, we consider the following pushout of filtered face sets,

$$\begin{tikzcd}
\underline{K} \otimes \partial\Delta^{m-1} \arrow[r] \arrow[d] & \underline{K} \otimes \Delta^{m,k} \arrow[d] \\
\underline{K} \otimes \Delta^{m-1} \arrow[r] & \underline{K} \otimes \partial\Delta^m.
\end{tikzcd}$$

  As the functor $\mathcal{A}$ satisfies adjunction properties (Proposition 11.2), the image of this pushout is a pullback and we have

$$\mathcal{A}(\underline{K} \otimes \partial\Delta^m)_\bullet = \mathcal{A}(\underline{K} \otimes \Delta^{m,k})_\bullet \times_{\mathcal{A}(\underline{K} \otimes \partial\Delta^{m-1})_\bullet} \mathcal{A}(\underline{K} \otimes \Delta^{m-1})_\bullet.$$

  The surjectivity of $\mathcal{A}(\underline{K} \otimes \partial\Delta^m)_\bullet \to \mathcal{A}(\underline{K} \otimes \Delta^{m,k})_\bullet$ follows from the surjectivity of $\mathcal{A}(\underline{K} \otimes \Delta^{m-1})_\bullet \to \mathcal{A}(\underline{K} \otimes \partial\Delta^{m-1})_\bullet$ already established.

□

Lemma 11.6. *If* $A_\bullet \in \mathrm{CDGA}_{\mathcal{F}}$ *is a cofibrant object, its realization,* $\langle A_\bullet \rangle$*, is a Kan filtered face set.*

Proof. Let $\underline{K}$ be a filtered face set. From Proposition 11.2 we know that the existence of a dotted arrow making commutative the following diagram of filtered face sets maps

$$\begin{tikzcd}
\underline{K} \otimes \Delta^{m,k} \arrow[r] \arrow[d] & \langle A_\bullet \rangle \\
\underline{K} \otimes \Delta^m \arrow[ur, dashed] &
\end{tikzcd}$$

is equivalent to the existence of a dotted arrow making commutative the following diagram in $\mathrm{CDGA}_{\mathcal{F}}$,

$$\begin{tikzcd}
A_\bullet \arrow[r] \arrow[dr, dashed, "g"'] & \mathcal{A}(\underline{K}\otimes\Delta^{m,k})_\bullet \\
 & \mathcal{A}(\underline{K}\otimes\Delta^{m})_\bullet . \arrow[u]
\end{tikzcd}$$

In this last diagram, the map g exists because $A_\bullet$ is cofibrant and the map $\mathcal{A}(\underline{K}\otimes\Delta^{m})_\bullet \to \mathcal{A}(\underline{K}\otimes\Delta^{m,k})_\bullet$ is a trivial fibration, by Lemma 11.5. $\square$

Lemma 11.7. *A path object in* $\mathrm{CDGA}_{\mathcal{F}}$ *can be built from the two injections,* ι_0, $\iota_1\colon \Delta^0 \to \Delta^1$, *as:*

$$\mathcal{A}(\underline{K})_\bullet \longrightarrow \mathcal{A}(\underline{K}\otimes\Delta^1)_\bullet \xrightarrow{\iota_0^*+\iota_1^*} \mathcal{A}(\underline{K})_\bullet \oplus \mathcal{A}(\underline{K})_\bullet .$$

Proof. The map $\iota_0^*+\iota_1^*$ coincides with the map $\mathcal{A}(\underline{K}\otimes\Delta^1)_\bullet \to \mathcal{A}(\underline{K}\otimes\partial\Delta^1)_\bullet$ induced by the canonical inclusion. This is a surjective map thus a fibration.

The projection $\Delta^1 \to \Delta^0$ induces a map $p\colon \underline{K}\otimes\Delta^1 \to \underline{K}$, which is not in $\mathbf{\Delta}^{[n]}_{\mathcal{F}}-$*Sets* but which induces a quasi-isomorphism of perverse CDGA's, $\mathcal{A}(\underline{K})_\bullet \to \mathcal{A}(\underline{K}\otimes\Delta^1)_\bullet$, according to Proposition 9.8 and Corollary 7.3.

By construction, the sequence of the statement is the map induced by the diagonal. With the previous properties, this sequence is a path object in the category $\mathrm{CDGA}_{\mathcal{F}}$, by definition. $\square$

Proof of Theorem E. First, we claim that, if f, $g\colon \underline{K} \to \underline{K}'$ are homotopic in $\mathbf{\Delta}^{[n]}_{\mathcal{F}}-$*Sets*, then, for any $B_\bullet \in \mathrm{CDGA}_{\mathcal{F}}$ cofibrant, we have

$$f^* = g^*\colon [B_\bullet, \mathcal{A}(\underline{K}')_\bullet]_{\mathrm{CDGA}_{\mathcal{F}}} \to [B_\bullet, \mathcal{A}(\underline{K})_\bullet]_{\mathrm{CDGA}_{\mathcal{F}}}.$$

It suffices to prove it for the two canonical injections, ι_0, $\iota_1\colon \Delta^0 \to \Delta^1$. The perverse CDGA, $B_\bullet$, being cofibrant, according to Lemma 11.7, Corollary 7.3 and (2) of Proposition 10.10, we have a bijection,

$$[B_\bullet, \mathcal{A}(\underline{K})_\bullet]_{\mathrm{CDGA}_{\mathcal{F}}} \xrightarrow[\cong]{} [B_\bullet, \mathcal{A}(\underline{K}\otimes\Delta^1)_\bullet]_{\mathrm{CDGA}_{\mathcal{F}}},$$

induced by $p\colon \underline{K}\otimes\Delta^1 \to \underline{K}$. From $p\circ\iota_0 = p\circ\iota_1 = \mathrm{id}$, we deduce $\iota_0^* = \iota_1^*\colon [B_\bullet, \mathcal{A}(\underline{K}\otimes\Delta^1)_\bullet] \to [B_\bullet, \mathcal{A}(\underline{K})_\bullet]$.

In a second step, we consider two morphisms, f, $g\colon B_\bullet \to C_\bullet$, in $\mathrm{CDGA}_{\mathcal{F}}$ and a homotopy between them, $\varphi\colon B_\bullet \to (C\otimes\wedge(t,dt))_\bullet$. We want to construct a homotopy, $\langle C_\bullet\rangle\otimes\Delta^1 \to \langle B_\bullet\rangle$, of filtered face sets, between $\langle f\rangle$ and $\langle g\rangle$. We first have a map, involving the hom functor in the category $\mathrm{CDGA}_{\mathcal{F}}$,

$$\begin{tikzcd}
\Big(\mathrm{hom}(C_\bullet, \widetilde{A}_{PL,\bullet}(\Delta^{k_0}*\cdots*\Delta^{k_n}))\otimes\Delta^1\Big)_{(j_0,\dots,j_n)} \arrow[d, "\Phi_{(j_0,\dots,j_n)}"] \\
\Big(\mathrm{hom}(C_\bullet, \widetilde{A}_{PL,\bullet}(\Delta^{k_0}*\cdots*\Delta^{k_n}))\otimes\mathrm{hom}(B_\bullet, C_\bullet\otimes A_{PL}(\Delta^1))\Big)_{(j_0,\dots,j_n)}
\end{tikzcd}$$

defined as the tensor product of the identity map on the first component with the map given by the homotopy φ on the second component. A second map, involving also the hom functor in the category $\mathrm{CDGA}_{\mathcal{F}}$,

$$\begin{tikzcd}
\Big(\hom(C_\bullet,\widetilde{A}_{PL,\bullet}(\Delta^{k_0}*\cdots*\Delta^{k_n}))\otimes\hom(B_\bullet,C_\bullet\otimes A_{PL}(\Delta^1))\Big)_{(j_0,\dots,j_n)} \arrow[d, "\Phi'_{(j_0,\dots,j_n)}"] \\
\hom(B_\bullet,\widetilde{A}_{PL,\bullet}(\Delta^{j_0}*\cdots*\Delta^{j_n}))
\end{tikzcd}$$

is defined by composition as follows: let $\varphi\otimes\psi$ in the domain of $\Phi'_{(j_0,\dots,j_n)}$, we first consider the composition

$$B_\bullet\xrightarrow{\psi}C_\bullet\otimes A_{PL}(\Delta^1)\xrightarrow{\varphi\otimes\mathrm{id}}\widetilde{A}_{PL,\bullet}(\Delta^{k_0}*\cdots*\Delta^{k_n})\otimes A_{PL}(\Delta^1),$$

and compose it with the map induced by the two projections of the product,

$$\widetilde{A}_{PL,\bullet}(\Delta^{k_0}*\cdots*\Delta^{k_n})\otimes A_{PL}(\Delta^1)\to\widetilde{A}_{PL,\bullet}((\Delta^{k_0}*\cdots*\Delta^{k_n})\otimes\Delta^1),$$

followed by the restriction map

$$\widetilde{A}_{PL,\bullet}((\Delta^{k_0}*\cdots*\Delta^{k_n})\otimes\Delta^1)\to\widetilde{A}_{PL,\bullet}(\Delta^{j_0}*\cdots*\Delta^{j_n}).$$

(To be complete, we add, at the end of this proof, the justification of the existence of this last map.) The map $\Phi'_{(j_0,\dots,j_n)}\circ\Phi_{(j_0,\dots,j_n)}$ gives the wanted homotopy between $\langle f\rangle$ and $\langle g\rangle$.

Denote by $\Phi\colon A_\bullet\to\mathcal{A}(\langle A_\bullet\rangle)_\bullet$, $\Psi\colon\underline{K}\to\langle\mathcal{A}(\underline{K})_\bullet\rangle$ the morphisms of adjunction and by $\chi:\hom_{\mathrm{CDGA}_{\mathcal{F}}}(A_\bullet,\mathcal{A}(\underline{K})_\bullet)\cong\hom_{\Delta^{[n]}_{\mathcal{F}}-Sets}(\underline{K},\langle A_\bullet\rangle)$ the isomorphism of Proposition 11.2. Let $f\colon A_\bullet\to\mathcal{A}(\underline{K})_\bullet$. The composite map

$$\underline{K}\xrightarrow{\Psi}\langle\mathcal{A}(\underline{K})_\bullet\rangle\xrightarrow{\langle f\rangle}\langle A_\bullet\rangle$$

is equal to $\chi(f)$. Moreover, the composition

$$A_\bullet\xrightarrow{\Phi}\mathcal{A}(\langle A_\bullet\rangle)_\bullet\xrightarrow{\mathcal{A}(\chi(f))}\mathcal{A}(\underline{K})_\bullet$$

is equal to f. With these remarks, the theorem is a direct consequence of the first two steps of this proof.

Lastly, we develop the promised technical point. Consider two simplices,

$$\Delta=\Delta^{k_0}*\cdots*\Delta^{k_{i-1}}*\Delta^{k_i+1}*\Delta^{k_{i+1}}*\cdots*\Delta^{k_n}$$

and

$$\nabla=\Delta^{k_0}*\cdots*\Delta^{k_{i-1}}*\Delta^{k_i}*\Delta^{k_{i+1}}*\cdots*\Delta^{k_n},$$

and a map $(f,g)\colon\Delta\to\nabla\times\Delta^1$ such that $f=\mathrm{id}*\cdots*f_i*\cdots*\mathrm{id}$, with $f_i\colon\Delta^{k_i+1}\to\Delta^{k_i}$ and $g\colon\Delta\to\Delta^1$ linear. Recall that $\mu_\Delta\colon\widetilde{\Delta}\to\Delta$ and $\mu_\nabla\colon\widetilde{\nabla}\to\nabla$ are the blow-up's, see Definition 5.7. We define $cf_i\colon c\Delta^{k_i+1}\to c\Delta^{k_i}$ by $cf_i([x,t]=[f_i(x),t]$ and $(\tilde{f},\tilde{g})=(\mathrm{id}\times\cdots\times cf_i\times\cdots\times\mathrm{id},g\circ\mu_\Delta)$. As $\tilde{f}$ is linear, the following diagram commutes,

$$\begin{tikzcd}
\widetilde{\Delta} \arrow[r, "(\tilde{f},\tilde{g})"] \arrow[d, "\mu_\Delta"'] & \widetilde{\nabla}\times\Delta^1 \arrow[d, "\mu_\nabla\times\mathrm{id}"] \\
\Delta \arrow[r, "(f,g)"] & \nabla\times\Delta^1.
\end{tikzcd}$$

As $\tilde{g} = g \circ \mu_\Delta$, with g linear and μ_Δ induces a CDGA's map $\mu_\Delta^* \colon A_{PL}(\Delta) \to A_{PL}(\widetilde{\Delta})$, there is an induced CDGA's map $\tilde{g}^* \colon A_{PL}(\Delta^1) \to A_{PL}(\widetilde{\Delta})$, and thus a CDGA's map, $A_{PL}(\widetilde{\nabla} \times \Delta^1) \to A_{PL}(\widetilde{\Delta})$.

Finally, the compatibility with perverse degrees comes from the inclusion

$$(\tilde{f}, \tilde{g})(\Delta^{k_i+1} \times \{1\}) \subset \Delta^{k_i} \times \{1\} \times \Delta^1.$$

□

12. Minimal models

In this section, we construct a minimal model of some perverse CDGA's, unique up to isomorphism. This is valuable for the algebra of perverse forms on a filtered face set $\underline{K}$ and for its cohomology. Moreover, in the case $\underline{K}$ is coming from a locally cone like space, the model we have built is a topological invariant. A notion of formality is defined and examples are given.

We are interested here by *non-negatively graded* Sullivan models in the sense of Definition 10.6. Recall that $\mathcal{P}^n$ is the lattice of GM-perversities and $\hat{\mathcal{P}}^n = \mathcal{P}^n \cup \{\overline{\infty}\}$. We use the lattice structure of $\hat{\mathcal{P}}_n$ for the construction of such models and begin first by a description of the predecessors of a given GM-perversity $\overline{p}$.

Definition 12.1. Let $\overline{p} \in \mathcal{P}_n$ be a GM-perversity. A *peak* of $\overline{p}$ is an integer $j \in \{2, \ldots, n-1\}$ such that $\overline{p}(j+1) = \overline{p}(j) > \overline{p}(j-1)$. The integer n is a peak if $\overline{p}(n) > \overline{p}(n-1)$. We order the peaks $(\mathtt{n}_1, \ldots, \mathtt{n}_s)$ of $\overline{p}$ by $1 \leq \mathtt{n}_1 < \mathtt{n}_2 < \ldots < \mathtt{n}_s \leq n$. For each peak $\mathtt{n}_i$ of $\overline{p}$, we define a perversity $\overline{p}_i$ by

$$\begin{cases} \overline{p}_i(k) = \overline{p}(k), \text{ if } k \neq \mathtt{n}_i, \\ \overline{p}_i(\mathtt{n}_i) = \overline{p}(\mathtt{n}_i) - 1. \end{cases}$$

We check easily the following properties.

(i) The previous perversities $(\overline{p}_1, \ldots, \overline{p}_s)$ are in $\mathcal{P}_n$.
(ii) The perversities $(\overline{p}_1, \ldots, \overline{p}_s)$ are the predecessors of $\overline{p}$. *We order them by the corresponding peaks.*
(iii) We have $\overline{p}(\mathtt{n}_i) \geq i$ and $\overline{p}_i(\mathtt{n}_i) \geq i-1$.
(iv) If $(\overline{p}_1, \ldots, \overline{p}_s)$ are the ordered predecessors of $\overline{p}$, then

$$(\inf(\overline{p}_1, \overline{p}_2), \inf(\overline{p}_1, \overline{p}_3), \ldots, \inf(\overline{p}_1, \overline{p}_s))$$

are ordered predecessors of $\overline{p}_1$ and the corresponding ordered peaks are $(\mathtt{n}_2, \ldots, \mathtt{n}_s)$.

We first build minimal models of cohomologically connected regular perverse CDGA's, in the following sense.

Definition 12.2. Let $A_\bullet$ be a perverse CDGA, then:

(a) $A_\bullet$ is *cohomologically connected* if $H^0(A_{\overline{p}}) = \mathbb{Q}$, for all $\overline{p} \in \hat{\mathcal{P}}_n$,
(b) $A_\bullet$ is *regular* if it is a strict perverse algebra whose elements satisfy $\|\omega\|_i \leq |\omega|$, for $i \in \{1, \ldots, n\}$.

By construction, the perverse CDGA of forms on a filtered face set is regular.

Theorem F (Construction of a minimal model). *Let $A_\bullet$ be a cohomologically connected, regular perverse* CDGA *or the cohomology of it. Then, there exists a positively graded, Sullivan minimal model of $A_\bullet$,*

$$\rho_\bullet \colon B_\bullet = (\wedge \oplus_{\overline{p}} V_{[\overline{p}]}, d)_\bullet \to A_\bullet,$$

i.e., $\rho_{\overline{p}}(B_{\overline{p}}) \subset A_{\overline{p}}$, the restriction $\rho_{\overline{p}} \colon B_{\overline{p}} \to A_{\overline{p}}$ is a quasi-isomorphism for any $\overline{p} \in \hat{\mathcal{P}}_n$ and the elements of $V_{[\overline{p}]}$ have a strictly positive degree. This model is unique up to isomorphism.

In Example 4.5, to any locally conelike space X (Definition 3.1) whose regular part is connected, we associate a filtered face set, $\underline{L} = \mathrm{ISing}^{\mathcal{F}}_*(X)$, such that $\mathcal{A}(\underline{L})_\bullet$ is a cohomologically connected, regular perverse CDGA.

Definition 12.3. A filtered face set, $\underline{K}$, is *intersection-formal* if there is an isomorphism between the minimal models of $\mathcal{A}(\underline{K})_\bullet$ and $H(\mathcal{A}(\underline{K})_\bullet)$.

A locally conelike space, X, whose regular part is connected, is *intersection-formal* if its associated filtered face set, $\underline{L} = \mathrm{ISing}^{\mathcal{F}}_*(X)$, is formal.

The next result is the topological invariance of the minimal model of $\underline{L}$.

Corollary 12.4. *Let X be a locally conelike space whose regular part is connected. Then the minimal model of the associated filtered face set, $\underline{L}$, does not depend on the stratification of X, in perversity degrees strictly less than $\overline{\infty}$.*

Remark 12.5. We emphasize that there is no stratum of codimension one in a locally conelike space and that the perversity $\overline{\infty}$ is not a topological invariant. For instance, we may observe that:

- a circle S^1, with one stratum of dimension 0 (codimension 1), has a trivial $\overline{0}$-cohomology in degree 1,
- a sphere S^2 with two strata of dimension 0 has a $\overline{1}$-cohomology isomorphic to the cohomology a circle.

We can construct minimal models in each of these two cases but they are not covered by Corollary 12.4, the first one having a stratum of codimension 1 and, for the second one, the perversity $\overline{1}$ being strictly greater than the top perversity.

Before giving the proofs of Theorem F and Corollary 12.4, we need to develop some properties of regular perverse CDGA's and of Sullivan algebras.

Proposition 12.6. *Let $A_\bullet$ be a regular perverse* CDGA *and $(\overline{p}_1, \dots, \overline{p}_j)$ be any sequence of ordered predecessors of a GM-perversity $\overline{p}$, associated to the peaks $(\mathtt{n}_1, \dots, \mathtt{n}_j)$. Then the following properties are satisfied.*

(1) For any loose perversities, $\overline{q}_1$ and $\overline{q}_2$, including $\overline{q}_2 = \overline{\infty}$, such that $0 \leq \overline{q}_1 \leq \overline{q}_2$, the canonical map $A_{\overline{q}_1} \hookrightarrow A_{\overline{q}_2}$ induces an injection $H^1(A_{\overline{q}_1}) \to H^1(A_{\overline{q}_2})$.

(2) For all $m < \overline{p}_j(\mathtt{n}_j)$, we have

  (i) $A_{\overline{p}_1} \cap A_{\overline{p}_2} = A_{\inf(\overline{p}_1, \overline{p}_2)}$,

  (ii) $A^m_{\overline{p}_k} \subset A^m_{\overline{p}_j}$, for all $k \in \{1, \dots, j\}$,

  (iii) $A^m_{\overline{p}_1} + \cdots + A^m_{\overline{p}_j} = A^m_{\overline{p}_j}$,

  (iv) $A^m_{\overline{p}_1} \cap (A^m_{\overline{p}_2} + \cdots + A^m_{\overline{p}_j}) = (A^m_{\overline{p}_1} \cap A^m_{\overline{p}_2}) + \cdots + (A^m_{\overline{p}_1} \cap A^m_{\overline{p}_j})$,

(v) $H^{m+1}((A_{\overline{p}_1}\cap A_{\overline{p}_2})+\cdots+(A_{\overline{p}_1}\cap A_{\overline{p}_j}))$ *injects in* $H^{m+1}(A_{\overline{p}_1}\cap(A_{\overline{p}_2}+\cdots+A_{\overline{p}_j}))$,

(vi) $H^{m+2}(A_{\overline{p}_j})\to H^{m+2}(A_{\overline{p}})$ *is injective,*

(vii) $H^{m+1}(A_{\overline{p}_j})\to H^{m+1}(A_{\overline{p}})$ *is surjective.*

(3) $H^1(A_{\overline{p}_1}+\cdots+A_{\overline{p}_j})\to H^1(A_{\overline{p}})$ *is injective,*

Proof. (1) Let $\omega\in A^1_{\overline{q}_1}$ such that $\omega=df$ with $f\in A^0_{\overline{q}_2}$. For any j, we have, by regularity of A,

$$\max(\|f\|_j,\|df\|_j)\le\max(0,\overline{q}_1(j))\le\overline{q}_1(j)$$

and $f\in A^0_{\overline{q}_1}$. The same argument works if $\overline{q}_2=\overline{\infty}$.

Property (2)(i) is a consequence of the definition of strict perverse CDGA.

(ii) Let $\omega\in A^m_{\overline{p}_k}$. We consider two cases:

- if $\ell\neq \mathtt{n}_j$, then $\|\omega\|_\ell\le\overline{p}_k(\ell)\le\overline{p}_j(\ell)$,
- if $\ell=\mathtt{n}_j$, then $\max(\|\omega\|_{\mathtt{n}_j},\|d\omega\|_{\mathtt{n}_j})\le m+1\le\overline{p}_j(\mathtt{n}_j)$.

Therefore $\omega\in A^m_{\overline{p}_j}$.

(iii) This is a direct consequence of (ii).

(iv) The inclusion $\supseteq$ is immediate. From property (ii), we deduce

$$\begin{array}{rcl}A^m_{\overline{p}_1}\cap(A^m_{\overline{p}_2}+\cdots+A^m_{\overline{p}_j}) & \subseteq & A^m_{\overline{p}_1}\cap A^m_{\overline{p}_j}\\ & \subseteq & (A^m_{\overline{p}_1}\cap A^m_{\overline{p}_2})+\cdots+(A^m_{\overline{p}_1}\cap A^m_{\overline{p}_j}).\end{array}$$

As for the assertion (v), let $\omega\in(A^{m+1}_{\overline{p}_1}\cap A^{m+1}_{\overline{p}_2})+\cdots+(A^{m+1}_{\overline{p}_1}\cap A^{m+1}_{\overline{p}_j})$ such that $d\omega=0$ and $\omega=d\alpha$, with $\alpha\in A^m_{\overline{p}_1}\cap(A^m_{\overline{p}_2}+\cdots+A^m_{\overline{p}_j})$. From assertion (iv), we deduce that ω is a coboundary also in $(A^{m+1}_{\overline{p}_1}\cap A^{m+1}_{\overline{p}_2})+\cdots+(A^{m+1}_{\overline{p}_1}\cap A^{m+1}_{\overline{p}_j})$.

(vi) Let $\omega\in A^{m+2}_{\overline{p}_j}$ and $\alpha\in A^{m+1}_{\overline{p}}$ with $d\alpha=\omega$. We consider two cases:

- if $\ell\neq\mathtt{n}_j$, then $\|\alpha\|_\ell\le\overline{p}(\ell)=\overline{p}_j(\ell)$,
- if $\ell=\mathtt{n}_j$, then $\|\alpha\|_{\mathtt{n}_j}\le m+1\le\overline{p}_j(\mathtt{n}_j)$.

This implies $\alpha\in A^m_{\overline{p}_j}$ and the injectivity.

(vii) Let $\omega\in A^{m+1}_{\overline{p}}$, $d\omega=0$. We consider two cases:

- if $\ell\neq\mathtt{n}_j$, then $\|\omega\|_\ell\le\overline{p}(\ell)=\overline{p}_j(\ell)$,
- if $\ell=\mathtt{n}_j$, then $\|\omega\|_{\mathtt{n}_j}\le m+1\le\overline{p}_j(\mathtt{n}_j)$.

This implies $\omega\in A_{\overline{p}_j}$.

(3) If $j=1$, it is proved in (1). If $j=2$, then, if $\omega\in A_{\overline{p}_1}+A_{\overline{p}_2}$ is such that $\omega=df$ with $f\in A_{\overline{p}}$, we have

$$\max(\|f\|_{\mathtt{n}_2},\|df\|_{\mathtt{n}_2})\le 1\le\overline{p}_2(\mathtt{n}_2).$$

As $\overline{p}$ and $\overline{p}_2$ differ only in $\mathtt{n}_2$, we have $f\in A_{\overline{p}_2}$ and the injectivity is proved. If $j\ge 3$, then $A^1_{\overline{p}_1}+\cdots+A^1_{\overline{p}_j}=A^1_{\overline{p}_j}$, by (2)(iii) and $1<2\le j-1\le\overline{p}_j(\mathtt{n}_j)$. The injectivity comes now from (1). ☐

We want to establish similar properties for the cohomology of $A_\bullet$. In this case, the sum $H_{\overline{p}_1}(A_\bullet)+H_{\overline{p}_2}(A_\bullet)$ is not defined and we first define the sum of cochain complexes as a pushout.

Definition 12.7. Suppose given two cochain maps

$$\begin{tikzcd} C_1 & C \arrow[l, "f_1"'] \arrow[r, "f_2"] & C_2 \end{tikzcd}$$

The sum $(C_1, f_1) \dotplus (C_2, f_2)$ is defined as the push-out of f_1 and f_2, i.e., its elements are the equivalence classes of couples $(c_1, c_2) \in C_1 \times C_2$ for the relation $(c_1, c_2) \sim (c_1', c_2')$ if, and only if, there exists $c \in C$ with $f_1(c) = c_1' - c_1$ and $-f_2(c) = c_2' - c_2$. We denote by $\langle c_1, c_2\rangle$ the class of (c_1, c_2).

We apply this notion of sum to any perverse CDGA, $C_\bullet$, as follows. Let $(\overline{p}_1, \dots, \overline{p}_j)$ be a sequence of ordered predecessors of a perversity $\overline{p} \in \mathcal{P}^n$.

(1) We denote by $C_{\overline{p}_1} \dotplus C_{\overline{p}_2}$ the sum associated to the diagram

$$\begin{tikzcd} C_{\overline{p}_1} & C_{\inf(\overline{p}_1, \overline{p}_2)} \arrow[l] \arrow[r] & C_{\overline{p}_2}. \end{tikzcd}$$

(2) Working by induction, we denote by $C_{\overline{p}_1} \dotplus \cdots \dotplus C_{\overline{p}_j}$ the sum associated to the diagram

$$\begin{tikzcd} C_{\overline{p}_1} & C_{\inf(\overline{p}_1, \overline{p}_2)} \dotplus \cdots \dotplus C_{\inf(\overline{p}_1, \overline{p}_j)} \arrow[l] \arrow[r] & C_{\overline{p}_2} \dotplus \cdots \dotplus C_{\overline{p}_j}. \end{tikzcd}$$

Remark 12.8. If $C_\bullet = A_\bullet$ is a regular perverse CDGA, the two sums, $+$ and $\dotplus$, are distinct. More precisely, we have $A_{\overline{p}_1} + A_{\overline{p}_2} = A_{\overline{p}_1} \dotplus A_{\overline{p}_2}$ but for a sum of three terms or more they do not coincide. For instance, in the case of three terms, the two sums correspond to the next short exact sequences,

$$0 \longrightarrow A_{\overline{p}_1} \cap (A_{\overline{p}_2} + A_{\overline{p}_3}) \longrightarrow A_{\overline{p}_1} \oplus (A_{\overline{p}_2} + A_{\overline{p}_3}) \longrightarrow A_{\overline{p}_1} + A_{\overline{p}_2} + A_{\overline{p}_3} \longrightarrow 0$$

and

$$0 \to (A_{\overline{p}_1} \cap A_{\overline{p}_2}) + (A_{\overline{p}_1} \cap A_{\overline{p}_3}) \to A_{\overline{p}_1} \oplus (A_{\overline{p}_2} + A_{\overline{p}_3}) \to A_{\overline{p}_1} \dotplus A_{\overline{p}_2} \dotplus A_{\overline{p}_3} \to 0.$$

The two left hand side terms are not isomorphic; we compare them in Proposition 12.6, in low degrees.

Proposition 12.9. *Let $A_\bullet$ be a regular perverse* CDGA, *$(\overline{p}_1, \dots, \overline{p}_j)$ be a sequence of ordered predecessors of a perversity $\overline{p} \in \mathcal{P}^n$, associated to the peaks $(\mathtt{n}_1, \dots, \mathtt{n}_j)$ and $f_i^* \colon H^*(A_{\overline{p}_i}) \to H^*(A_{\overline{p}})$ be the induced maps for $i \in \{1, \dots, j\}$. Then, the following properties are satisfied for all $m < \overline{p}_j(\mathtt{n}_j)$.*

(1) The map $\psi^{m+1} \colon H^{m+1}(A_{\overline{p}_j}) \to H^{m+1}(A_{\overline{p}_1}) \dotplus \cdots \dotplus H^{m+1}(A_{\overline{p}_j})$, which sends $[\omega]$ on $\langle 0, \langle 0, \dots, \langle 0, [\omega]\rangle \cdots \rangle$, is surjective.

(2) The cochain map $f^{m+1} \colon H^{m+1}(A_{\overline{p}_1}) \dotplus \cdots \dotplus H^{m+1}(A_{\overline{p}_j}) \to H^{m+1}(A_{\overline{p}})$,

$$f^{m+1}(\langle \omega_1, \langle \omega_2, \dots, \langle \omega_{j-1}, \omega_j \rangle \cdots \rangle) = f_1^{m+1}(\omega_1) + \cdots + f_j^{m+1}(\omega_j)$$

is well defined and injective.

(3) In particular, the cochain map $f^1 \colon H^1(A_{\overline{p}_1}) \dotplus \cdots \dotplus H^1(A_{\overline{p}_j}) \to H^1(A_{\overline{p}})$ is well defined and injective.

Proof. (1) Let

$$\Upsilon = \langle [\omega_1], \langle [\omega_2], \dots, \langle [\omega_{j-1}], [\omega_j]\rangle \cdots \rangle \in H^{m+1}(A_{\overline{p}_1}) \dotplus \cdots \dotplus H^{m+1}(A_{\overline{p}_j}).$$

As the induced map, $f^*_{k,j}\colon H^{m+1}(A_{\inf(\overline{p}_k,\overline{p}_j)})\to H^{m+1}(A_{\overline{p}_k})$ is surjective (cf. Proposition 12.6) for $k\in\{1,\ldots,j-1\}$, we have

$$\begin{aligned}\Upsilon &= \langle f^*_{1,j}[\omega'_1],\langle f^*_{2,j}[\omega'_2],\ldots,\langle f^*_{j-1,j}[\omega'_{j-1}],[\omega_j]\rangle\cdots\rangle\\ &= \langle 0,\langle 0,\ldots\langle 0,[\omega'_j]\rangle\cdots\rangle\\ &= \psi^{m+1}([\omega'_j]).\end{aligned}$$

(2) We consider the following commutative diagram

$$\begin{tikzcd} & H^{m+1}(A_{\overline{p}}) & \\ H^{m+1}(A_{\overline{p}_j}) \arrow[ur, "f_j^{m+1}"] \arrow[rr, "\psi^{m+1}"'] & & H^{m+1}(A_{\overline{p}_1})\dotplus\cdots\dotplus H^{m+1}(A_{\overline{p}_j}). \arrow[ul, "f^{m+1}"'] \end{tikzcd}$$

The first property implies the surjectivity of ψ^{m+1} and Proposition 12.6 gives the injectivity of f_j^{m+1}. Therefore, the map f^{m+1} is injective.

(3) For $j=1$, this is proved in Proposition 12.6. For $j\geq 2$, this is a direct consequence of (2) and $0<1\leq j-1\leq \overline{p}_j(\mathtt{n}_j)$. ☐

Proposition 12.10. *Let* $B=\wedge\oplus_{\overline{p}}V_{[\overline{p}]}$. *Recall that* $B_{[\overline{p}]}$ *is the vector space of elements of perverse graduation* $\overline{p}$ *and* $B_{\overline{p}}=\oplus_{\overline{r}\leq\overline{p}}B_{[\overline{r}]}$. *For any perversities,* $\overline{p}$, $\overline{q}_1,\ldots,\overline{q}_j$, *we have*

$$\begin{aligned}B_{\overline{q}_1}\cap B_{\overline{q}_2} &= B_{\inf(\overline{q}_1,\overline{q}_2)},\\ B_{\overline{p}}\cap(B_{\overline{q}_1}+\cdots+B_{\overline{q}_j}) &= (B_{\overline{p}}\cap B_{\overline{q}_1})+\cdots+(B_{\overline{p}}\cap B_{\overline{q}_j}).\end{aligned}$$

Proof. The first equality comes directly from definitions of $B_\bullet$. The inclusion $(B_{\overline{p}}\cap B_{\overline{q}_1})+\cdots+(B_{\overline{p}}\cap B_{\overline{q}_j})\subset B_{\overline{p}}\cap(B_{\overline{q}_1}+\cdots+B_{\overline{q}_j})$ being obvious, let $\omega\in B_{\overline{p}}\cap(B_{\overline{q}_1}+\cdots+B_{\overline{q}_j})$. Denote by

$$\mathcal{J}_i=\{\overline{r}\mid\overline{r}\leq\overline{q}_i\}\text{ and }\mathcal{J}=\{\overline{r}\mid\overline{r}\leq\overline{p}\}.$$

We write ω as a linear combination of elements of $B_{[\overline{r}]}$, $\overline{r}\in\cup_{i=1}^j\mathcal{J}_i$,

$$\omega=\sum_{i=1}^j\sum_{\overline{r}\in\mathfrak{I}_i}v_{i,[\overline{r}]},\text{ with }v_{i,[\overline{r}]}\in B_{[\overline{r}]}.$$

Decompose now each $\mathcal{J}_i$ in two disjoint subsets

$\mathfrak{I}_{i,1}=\mathcal{J}_i\cap\mathcal{J}=\{\overline{r}\mid\overline{r}\leq\overline{q}_i\text{ and }\overline{r}\leq\overline{p}\}$, $\mathfrak{I}_{i,2}=\mathcal{J}_i\backslash\mathcal{J}_{i,1}=\{\overline{r}\mid\overline{r}\leq\overline{q}_i\text{ and }\overline{r}\not\leq\overline{p}\}$.

Observe that

$$\sum_{i=1}^j\sum_{\overline{r}\in\mathfrak{I}_{i,2}}v_{i,[\overline{r}]}=\omega-\sum_{i=1}^j\sum_{\overline{r}\in\mathfrak{I}_{i,1}}v_{i,[\overline{r}]}\in B_{\overline{p}}.$$

Any element of $B_{\overline{p}}$ can be written as a sum of elements of $B_{[\overline{r}]}$ with $\overline{r}\in\mathcal{J}$. The intersections $\mathcal{J}\cap\mathcal{J}_{i,2}$, $B_{[\overline{r}]}\cap B_{[\overline{r}']}$ being the empty set if $\overline{r}\neq\overline{r}'$, we get $\sum_{i=1}^j\sum_{\overline{r}\in\mathfrak{I}_{i,2}}v_{i,[\overline{r}]}=0$ and

$$\omega=\sum_{i=1}^j\sum_{\overline{r}\in\mathfrak{I}_{i,1}}v_{i,[\overline{r}]}\in(B_{\overline{p}}\cap B_{\overline{q}_1})+\cdots+(B_{\overline{p}}\cap B_{\overline{q}_j}).$$

☐

The next results are the keys for the existence of a relative model, see [35, Theorem 5.1], the first one for a model of $A_\bullet$ and the second one for a model of $H(A_\bullet)$.

Proposition 12.11. *Let $A_\bullet$ be a cohomologically connected regular perverse* CDGA *and $\overline{p} \in \mathcal{P}^n$. Suppose we have a morphism of* CDGA*'s,*

$$\rho\colon B = (\wedge \oplus_{\overline{r}<\overline{p}} V_{[\overline{r}]}, d) \to A_{\overline{\infty}},$$

satisfying the next properties, for any $\overline{r} < \overline{p}$,

(i) $V^{\leq 0}_{[\overline{r}]} = 0$,

(ii) $\rho(B_{\overline{r}}) \subset A_{\overline{r}}$ and the restriction $\rho_{\overline{r}}\colon B_{\overline{r}} \to A_{\overline{r}}$ is a quasi-isomorphism.

Then the map $H^1_{\overline{p}}(\rho)\colon H^1(B_{\overline{p}}) \to H^1(A_{\overline{p}})$ is injective.

Proof. Let $(\overline{p}_1, \dots, \overline{p}_s)$ be the predecessors of $\overline{p}$. We first establish the property

$$B^1_{\overline{p}} = B^1_{\overline{p}_1} + \cdots + B^1_{\overline{p}_s}. \tag{11}$$

The inclusion $B^1_{\overline{p}_1} + \cdots + B^1_{\overline{p}_s} \subset B^1_{\overline{p}}$ being obvious, we consider $\omega \in B^1_{\overline{p}}$. This element can be written as a sum $\omega = \sum_i \lambda_i\, \omega_i$, with $|\lambda_i| = 0$ and $\omega_i \in \oplus_{\overline{r}<\overline{p}} V^1_{[\overline{r}]}$. As $V^0_{[\overline{r}]} = 0$ and $V^1_{[\overline{p}]} = 0$, we have $\lambda_i \in \mathbb{Q}$ and $\omega_i \in B_{\overline{p}_i}$ for a certain predecessor, $\overline{p}_i$, of $\overline{p}$. Regrouping these elements, we write ω as a sum of elements of $(B_{\overline{p}_i})_{1\leq i\leq s}$ and the equality (11) is proved. As we have a cochain complex, we deduce from (11)

$$H^1(B^1_{\overline{p}}) = H^1(B^1_{\overline{p}_1} + \cdots + B^1_{\overline{p}_s}).$$

For getting the injectivity of $H^1_{\overline{p}}(\rho)\colon H^1(B_{\overline{p}}) \to H^1(A_{\overline{p}})$, we decompose it in two maps

$$H^1(B_{\overline{p}_1} + \cdots + B_{\overline{p}_s}) \xrightarrow{f} H^1(A_{\overline{p}_1} + \cdots + A_{\overline{p}_s}) \xrightarrow{g} H^1(A_{\overline{p}}). \tag{12}$$

The injectivity of g has been proved in Proposition 12.6 and we consider now the map f.

Let $(\overline{q}_1, \dots, \overline{q}_k)$ be any ordered set of k predecessors of a perversity $\overline{q} \in \hat{\mathcal{P}}^n$, $\overline{q} \leq \overline{p}$. We denote by $\mathtt{n}_i$ the peak associated to $\overline{q}_i$ and by $\varphi_k\colon B_{\overline{q}_1} + \cdots + B_{\overline{q}_k} \to A_{\overline{q}_1} + \cdots + A_{\overline{q}_k}$ the map induced by ρ. For $k = 1$, φ_1 is a quasi-isomorphism by hypothesis. We prove by induction on k that

$$H^i(\varphi_k) \quad \begin{array}{l} \text{is an isomorphism for } i < \overline{q}_k(\mathtt{n}_k) - k + 2, \\ \text{is injective for } i = \overline{q}_k(\mathtt{n}_k) - k + 2. \end{array} \tag{13}$$

As $\overline{q}_k(\mathtt{n}_k) - k + 2 \geq 1$, the statement will be established. We suppose that (13) is true for any ordered set of k predecessors $(\overline{q}_1, \dots, \overline{q}_k)$ of any perversity $\overline{q} \in \hat{\mathcal{P}}^n$, $\overline{q} \leq \overline{p}$, and we prove it for an ordered set of $(k+1)$ predecessors $(\overline{q}_1, \dots, \overline{q}_{k+1})$. We use the following morphism of short exact sequences, where the map φ'_k is induced by ρ.

$$\begin{tikzcd}
0 \arrow[r] & B_{\overline{q}_1} \cap (B_{\overline{q}_2} + \cdots + B_{\overline{q}_{k+1}}) \arrow[r] \arrow[d, "\Psi_k"] & B_{\overline{q}_1} \oplus (B_{\overline{q}_2} + \cdots + B_{\overline{q}_{k+1}}) \arrow[r] \arrow[d, "\varphi_1 \oplus \varphi'_k"] & B_{\overline{q}_1} + \cdots + B_{\overline{q}_{k+1}} \arrow[r] \arrow[d, "\varphi_{k+1}"] & 0 \\
0 \arrow[r] & A_{\overline{q}_1} \cap (A_{\overline{q}_2} + \cdots + A_{\overline{q}_{k+1}}) \arrow[r] & A_{\overline{q}_1} \oplus (A_{\overline{q}_2} + \cdots + A_{\overline{q}_{k+1}}) \arrow[r] & A_{\overline{q}_1} + \cdots + A_{\overline{q}_{k+1}} \arrow[r] & 0
\end{tikzcd}$$

If $k = 1$, the maps Ψ_k and $\varphi_1 \oplus \varphi'_k$ reduce in $\Psi_1\colon B_{\inf(\overline{q}_1,\overline{q}_2)} \to A_{\inf(\overline{q}_1,\overline{q}_2)}$ and $\varphi_1 \oplus \varphi'_1\colon B_{\overline{q}_1} \oplus B_{\overline{q}_2} \to A_{\overline{q}_1} \oplus A_{\overline{q}_2}$. They are quasi-isomorphisms and the map φ_2 is a quasi-isomorphism by application of the 5-Lemma. We suppose now $k \geq 2$.

Recall that $B_{\overline{q}_1} \cap (B_{\overline{q}_2} + \cdots + B_{\overline{q}_{k+1}}) = B_{\inf(\overline{q}_1,\overline{q}_2)} + \cdots + B_{\inf(\overline{q}_1,\overline{q}_{k+1})}$. The map Ψ_k is the composition of

$$\psi_k \colon B_{\inf(\overline{q}_1,\overline{q}_2)} + \cdots + B_{\inf(\overline{q}_1,\overline{q}_{k+1})} \to A_{\inf(\overline{q}_1,\overline{q}_2)} + \cdots + A_{\inf(\overline{q}_1,\overline{q}_{k+1})}$$

followed by

$$\psi'_k \colon A_{\inf(\overline{q}_1,\overline{q}_2)} + \cdots + A_{\inf(\overline{q}_1,\overline{q}_{k+1})} \to A_{\overline{q}_1} \cap (A_{\overline{q}_2} + \cdots + A_{\overline{q}_{k+1}}).$$

Applying the induction to the k predecessors, $(\inf(\overline{q}_1,\overline{q}_2),\ldots,\inf(\overline{q}_1,\overline{q}_{k+1}))$, of $\overline{q}_1$, we have that $H^i(\psi_k)$ is an isomorphism for $i < \overline{q}_{k+1}(\mathtt{n}_{k+1}) - k + 2$ and an injection in degree $\overline{q}_{k+1}(\mathtt{n}_{k+1}) - k + 2$. Using Proposition 12.6, we get that the map $H^i(\psi'_k)$ is an isomorphism if $i < \overline{q}_{k+1}(\mathtt{n}_{k+1})$ and an injection if $i = \overline{q}_{k+1}(\mathtt{n}_{k+1})$. As $k \geq 2$, we have $\overline{q}_{k+1}(\mathtt{n}_{k+1}) - k + 2 \leq \overline{q}_{k+1}(\mathtt{n}_{k+1})$ and the composition $H^i(\Psi_k)$ is an isomorphism for $i < \overline{q}_{k+1}(\mathtt{n}_{k+1}) - k + 2$ and an injection in degree $\overline{q}_{k+1}(\mathtt{n}_{k+1}) - k + 2$. With the induction applied to the predecessors $(\overline{q}_2,\ldots,\overline{q}_{k+1})$ of $\overline{q}$, we obtain that $H^i(\varphi_1 \oplus \varphi'_k)$ verifies the same properties than $H^i(\Psi_k)$. An application of the four lemma (see [26]) ends the induction by giving that $H^i(\varphi_{k+1})$ is an isomorphism for $i < \overline{q}_{k+1}(\mathtt{n}_{k+1}) - (k+1) + 2$ and an injection for $i = \overline{q}_{k+1}(\mathtt{n}_{k+1}) - (k+1) + 2$. □

Proposition 12.12. *Let $A_\bullet$ be a cohomologically connected regular perverse* CDGA *and $\overline{p} \in \mathcal{P}^n$. Suppose we have a morphism of* CDGA*'s,*

$$\rho \colon B = (\wedge \oplus_{\overline{r}<\overline{p}} V_{[\overline{r}]}, d) \to H(A_{\overline{\infty}}),$$

satisfying the next properties, for any $\overline{r} < \overline{p}$,

(i) $V^{\leq 0}_{[\overline{r}]} = 0$,

(ii) $\rho(B_{\overline{r}}) \subset H(A_{\overline{r}})$ *and the restriction* $\rho_{\overline{r}} \colon B_{\overline{r}} \to H(A_{\overline{r}})$ *is a quasi-isomorphism.*

Then the map $H^1_{\overline{p}}(\rho) \colon H^1(B_{\overline{p}}) \to H^1(A_{\overline{p}})$ *is injective.*

Proof. Let $(\overline{q}_1,\ldots,\overline{q}_k)$ be any ordered set of k predecessors of a perversity $\overline{q} \in \hat{\mathcal{P}}^n$, $\overline{q} \leq \overline{p}$. We denote by $\mathtt{n}_i$ the peak associated to $\overline{q}_i$. We set $B^*_{\overline{q}_{1,k}} = B^*_{\inf(\overline{q}_1,\overline{q}_k)}$, $H^*_{\overline{q}_k} = H^*(A_{\overline{q}_k})$ and $H^*_{\overline{q}_{1,k}} = H^*(A_{\inf(\overline{q}_1,\overline{q}_k)})$ and denote by $\varphi_k \colon B_{\overline{q}_1} + \cdots + B_{\overline{q}_k} \to H_{\overline{q}_1} \dot{+} \cdots \dot{+} H_{\overline{q}_k}$ the map induced by ρ. As in the proof of Proposition 12.11, we reduce the problem to the proof that

$$H^i(\varphi_k) \text{ is an isomorphism for } i \leq \overline{q}_k(\mathtt{n}_k) - k + 2, \tag{14}$$

for any ordered set of k predecessors, $(\overline{q}_1,\ldots,\overline{q}_k)$, of any perversity $\overline{q} \in \hat{\mathcal{P}}^n$, $\overline{q} \leq \overline{p}$. As $\overline{q}_k(\mathtt{n}_k) - k + 2 \geq 1$, the statement will be established.

We do an induction on k by supposing that (14) is true for any ordered set of k predecessors, $(\overline{q}_1,\ldots,\overline{q}_k)$, of any perversity $\overline{q} \in \hat{\mathcal{P}}^n$, $\overline{q} \leq \overline{p}$, and we prove it for an ordered set of $(k+1)$ predecessors $(\overline{q}_1,\ldots,\overline{q}_{k+1})$. By hypothesis, $H(\varphi_1)$ is an isomorphism and we may suppose $k \geq 2$. From (2) of Proposition 12.9, we obtain a commutative diagram where φ'_k is induced

by ρ and whose lines are exact if $* \leq \overline{q}_{k+1}(\mathtt{n}_{k+1})$,
(15)

$$\begin{tikzcd}
0 \arrow[r] & B^*_{\overline{q}_{1,2}} + \cdots + B^*_{\overline{q}_{1,k+1}} \arrow[r] \arrow[d, "\Psi_k"] & B^*_{\overline{q}_1} \oplus (B^*_{\overline{q}_2} + \cdots + B^*_{\overline{q}_{k+1}}) \arrow[r] \arrow[d, "\varphi_1 \oplus \varphi'_k"] & B^*_{\overline{q}_1} + \cdots + B^*_{\overline{q}_{k+1}} \arrow[r] \arrow[d, "\varphi_{k+1}"] & 0 \\
0 \arrow[r] & H^*_{\overline{q}_{1,2}} \dot{+} \cdots \dot{+} H^*_{\overline{q}_{1,k+1}} \arrow[r] & H^*_{\overline{q}_1} \oplus (H^*_{\overline{q}_2} \dot{+} \cdots \dot{+} H^*_{\overline{q}_{k+1}}) \arrow[r] & H^*_{\overline{q}_1} \dot{+} \cdots \dot{+} H^*_{\overline{q}_{k+1}} \arrow[r] & 0
\end{tikzcd}$$

If $k = 1$, from (2)(vi) of Proposition 12.6, we know that $H^{\overline{q}_2(\mathtt{n}_2)+1}_{\overline{q}_{1,2}} \to H^{\overline{q}_2(\mathtt{n}_2)+1}_{\overline{q}_1}$ is injective and the morphism between the long exact sequences induced in cohomology can be extended on the right by the next diagram whose lines are exact and whose columns are isomorphisms by hypothesis, except $H^{\overline{q}_2(\mathtt{n}_2)}(\varphi_2)$,

$$\begin{tikzcd}
H^{\overline{q}(\mathtt{n}_2)}(B_{\overline{q}_{1,2}}) \arrow[r] \arrow[d] & H^{\overline{q}(\mathtt{n}_2)}(B_{\overline{q}_1}) \oplus H^{\overline{q}(\mathtt{n}_2)}(B_{\overline{q}_2}) \arrow[r] \arrow[d] & H^{\overline{q}(\mathtt{n}_2)}(B_{\overline{q}_1} \oplus B_{\overline{q}_2}) \arrow[r] \arrow[d, "H^{\overline{q}_2(\mathtt{n}_2)}(\varphi_2)"] & {} \\
H^{\overline{q}(\mathtt{n}_2)}_{\overline{q}_{1,2}} \arrow[r] & H^{\overline{q}(\mathtt{n}_2)}_{\overline{q}_1} \dot{+} H^{\overline{q}(\mathtt{n}_2)}_{\overline{q}_2} \arrow[r] & H^{\overline{q}(\mathtt{n}_2)}_{\overline{q}_1} \dot{+} H^{\overline{q}(\mathtt{n}_2)}_{\overline{q}_2} \arrow[r] & {}
\end{tikzcd}$$

$$\begin{tikzcd}
H^{\overline{q}(\mathtt{n}_2)+1}(B_{\overline{q}_{1,2}}) \arrow[r] \arrow[d] & H^{\overline{q}(\mathtt{n}_2)+1}(B_{\overline{q}_1}) \oplus H^{\overline{q}(\mathtt{n}_2)+1}(B_{\overline{q}_2}) \arrow[d] \\
H^{\overline{q}(\mathtt{n}_2)+1}_{\overline{q}_{1,2}} \arrow[r] & H^{\overline{q}(\mathtt{n}_2)+1}_{\overline{q}_1} \dot{+} H^{\overline{q}(\mathtt{n}_2)+1}_{\overline{q}_2}
\end{tikzcd}$$

From the 5-Lemma, we deduce that

$$H^i(\varphi_2)\colon H^i(B_{\overline{q}_1} + B_{\overline{q}_2}) \to H^i_{\overline{q}_1} \dot{+} H^i_{\overline{q}_2}$$

is an isomorphism for $i \leq \overline{q}_2(\mathtt{n}_2)$. Suppose now $k \geq 2$. By the induction hypothesis,

- $H^i(\Psi_k)$ is an isomorphism for $i \leq \inf(\overline{q}_1, \overline{q}_{k+1})(\mathtt{n}_{k+1}) - k + 2 = \overline{q}_{k+1}(\mathtt{n}_{k+1}) - k + 2$,
- $H^i(\varphi'_k)$ is an isomorphism for $i \leq \overline{q}_{k+1}(\mathtt{n}_{k+1}) - k + 2$.

As $\overline{q}_{k+1}(\mathtt{n}_{k+1}) - k + 2 \leq \overline{q}_{k+1}(\mathtt{n}_{k+1})$, we may apply the 5-Lemma to the diagram (15) and deduce that $H^i(\varphi_{k+1})$ is an isomorphism for $i \leq \overline{q}_{k+1}(\mathtt{n}_{k+1}) - k + 2 - 1 = \overline{q}_{k+1}(\mathtt{n}_{k+1}) - (k+1) + 2$. □

Proof of Theorem F. The uniqueness up to isomorphism is a direct consequence of Proposition 10.8 and (2) of Proposition 10.10.

The construction of this model begins with a classical minimal model $(\wedge V_{[\overline{0}]}, d)$ of the CDGA $A_{\overline{0}}$. We write the proof for the regular perverse CDGA, $A_\bullet$, the same argument gives the result for $H(A_\bullet)$, replacing Proposition 12.11 by Proposition 12.12.

Let $\overline{p} \in \hat{\mathcal{P}}^n$ be a fixed perversity and suppose we have a morphism of CDGA's,

$$\rho\colon B = (\wedge \oplus_{\overline{r} < \overline{p}} V_{[\overline{r}]}, d) \to A_{\overline{\infty}},$$

satisfying the next properties, for any $\overline{r} < \overline{p}$,

(i) $V^{\leq 0}_{[\overline{r}]} = 0$,
(ii) $\rho(B_{\overline{r}}) \subset A_{\overline{r}}$ and the restriction $\rho_{\overline{r}}\colon B_{\overline{r}} \to A_{\overline{r}}$ is a quasi-isomorphism.

For having a model for the perversity $\overline{p}$, we suppose that we have extended ρ in a morphism of CDGA's,

$$\rho\colon B' = (B \otimes \wedge V_{[\overline{p}]}^{<m}, d) = (\wedge(\oplus_{\overline{r}<\overline{p}} V_{[\overline{r}]} \oplus V_{[\overline{p}]}^{<m}), d) \to A_{\overline{\infty}},$$

such that,

(iii) $V_{\overline{p}}^{\leq 0} = 0$,

(iv) $\rho(V_{[\overline{p}]}^{<m}) \subset A_{\overline{p}}$ and the restriction $\rho_{\overline{p}}\colon B'_{\overline{p}} \to A_{\overline{p}}$ verifies,

   a) $H^i(\rho_{\overline{p}})$ is an isomorphism for any $i \leq m-1$,

   b) $H^i(\rho_{\overline{p}})$ is injective if $i = m$.

Observe that Proposition 12.11 implies these properties for $m = 1$ in the case $\overline{p} \in \mathcal{P}^n$ and Proposition 12.6 in the case $\overline{p} = \overline{\infty}$. We show now that the morphism ρ can be extended in a morphism of CDGA's, $\rho\colon B' \otimes \wedge V_{[\overline{p}]}^m \to A_{\overline{\infty}}$, so that properties (iii) and (iv) are satisfied for $m+1$.

We set

$$\left\{\begin{array}{rcl} Y_{[\overline{p}]}^m(0) & = & \operatorname{Coker} H^m(\rho_{\overline{p}})\colon H^m(B'_{\overline{p}}) \to H^m(A_{\overline{p}}), \\ Z_{[\overline{p}]}^m(0) & = & \operatorname{Ker} H^{m+1}(\rho_{\overline{p}})\colon H^{m+1}(B'_{\overline{p}}) \to H^{m+1}(A_{\overline{p}}), \\ V_{[\overline{p}]}^m(0) & = & Y_{\overline{p}}^m(0) \oplus Z_{\overline{p}}^m(0). \end{array}\right.$$

We extend the differential d and the morphism ρ by:

- if $y \in Y_{[\overline{p}]}^m(0)$, then $dy = 0$ and $\rho(y) \in A_{\overline{p}}^m$ is a cocycle representing the class in $\operatorname{Coker} H^m(\rho_{\overline{p}})$ corresponding to y,
- if $z \in Z_{[\overline{p}]}^m(0)$, then $dz \in B'_{\overline{p}}$ is a cocycle in the class corresponding to z in $\operatorname{Ker} H^{m+1}(\rho_{\overline{p}})$. The element $\rho(z) \in A_{\overline{p}}^m$ is defined by the equality $\rho \circ d = d \circ \rho$.

By construction, we have

$$d(V_{[\overline{p}]}^m(0)) \subset B'_{\overline{p}} = (B \otimes \wedge V_{[\overline{p}]}^{<m})_{\overline{p}} \text{ and } \rho(V_{[\overline{p}]}^m(0)) \subset A_{\overline{p}}.$$

Let j be a fixed integer. Suppose we have defined, for all $i \in \{0, \ldots, j\}$, $V_{[\overline{p}]}^m(i)$, ρ and d on $B' \otimes \wedge \oplus_{i \leq j} V_{\overline{p}}^m(i)$ so that

$$\left\{\begin{array}{l} d(V_{[\overline{p}]}^m(k)) \in (B' \otimes \wedge \oplus_{i \leq k-1} V_{[\overline{p}]}^m(i))_{\overline{p}}, \text{ for all } k \leq j, \\ \rho \circ d = d \circ \rho \text{ and } \rho(V_{[\overline{p}]}^m(k)) \subset A_{\overline{p}}. \end{array}\right.$$

We set

(16) $V_{[\overline{p}]}^m(j+1) = \operatorname{Ker} H^{m+1}(\rho_{\overline{p}})\colon H^{m+1}(B' \otimes \wedge \oplus_{i \leq j} V_{[\overline{p}]}^m(i))_{\overline{p}} \to H^{m+1}(A_{\overline{p}})$.

We extend the differential d by choosing a representing cocycle for each element of $\operatorname{Ker} H^{m+1}(\rho_{\overline{p}})$ and define ρ so that $\rho \circ d = d \circ \rho$. Finally, we set

$$V_{[\overline{p}]}^m = \oplus_i V_{[\overline{p}]}^m(i).$$

We verify now properties (i), (ii), (iii), (iv) for $\rho\colon (B' \otimes \wedge V_{[\overline{p}]}^m, d) \to A_{\overline{\infty}}$. Properties (i) and (iii) are satisfied by construction. If $\overline{r} < \overline{p}$, then we have $(B' \otimes \wedge V_{[\overline{p}]}^m)_{\overline{r}} = B'_{\overline{r}}$ and $H^i(\rho_{\overline{r}})$ *is an isomorphism for all* i. Thus (ii) is satisfied also. We are therefore reduced to the perversity $\overline{p}$.

- If $i < m$, then $(B' \otimes \wedge V_{[\overline{p}]}^m)_{\overline{p}}^i = B'^i_{\overline{p}}$ and $H^i(\rho_{\overline{p}})$ *is an isomorphism.*
- *The morphism* $H^m(\rho_{\overline{p}})$ *is surjective* by construction of $Y_{\overline{p}}^m(0)$.

- *The morphism $H^m(\rho_{\overline{p}})$ is injective.* To prove that, we consider $\omega \in (B' \otimes \wedge V^m_{[\overline{p}]})^m_{\overline{p}}$ such that $d\omega = 0$ and $\rho(\omega)$ is a boundary in $A^m_{\overline{p}}$. There exists j with $\omega \in B' \otimes \wedge \oplus_{i \leq j+1} V^m_{[\overline{p}]}(i)$. We set $\hat{B} = B' \otimes \wedge \oplus_{i \leq j} V^m_{[\overline{p}]}(i)$ and decompose ω in
$$\omega = \omega_1 + \omega_2 \in (\hat{B}^0 \otimes V^m_{[\overline{p}]}(j+1)) \oplus \hat{B}^m.$$
As, by induction, $\hat{B}^0 = \mathbb{Q}$, we have $\omega_1 \in V^m_{[\overline{p}]}(j+1)$. The equality $d\omega = 0$ implies $d\omega_1 \in d(\hat{B}^m)$. By construction of $V^m_{[\overline{p}]}(j+1)$, see (16), this implies $\omega_1 = 0$ and $\omega \in B' \otimes \wedge \oplus_{i \leq j} V^m_{[\overline{p}]}(i)$. We iterate this process and, at the end, we get $\omega \in B'_{\overline{p}}$. Now hypothesis (iv) b) implies the nullity in $\hat{B}$ of the cohomology class associated to ω.
- *The morphism $H^{m+1}(\rho_{\overline{p}})$ is injective.* Let $\omega \in (B' \otimes \wedge V^m_{[\overline{p}]})^{m+1}_{\overline{p}}$ such that $d\omega = 0$ and $\rho(\omega)$ is a boundary in $A_{\overline{p}}$. There exist an integer j with $\omega \in (B \otimes \wedge \oplus_{i \leq j} V^m_{[\overline{p}]}(i))^{m+1}_{\overline{p}}$ and, by construction of $V^m_{[\overline{p}]}(j+1)$, an element $v \in V^m_{[\overline{p}]}(j+1)$ such that $\omega - d(1 \otimes v)$ is a boundary in $B' \otimes \wedge \oplus_{i \leq j} V^m_{[\overline{p}]}(i)$. Thus ω is a boundary in $B' \otimes \wedge V^m_{[p]}$.

This model is minimal by construction. □

Proof of Corollary 12.4. In [23, Page 150], King associates to any locally conelike space, X, a locally conelike space, X^*, which is an intrinsic coarsest stratification of X. In particular, the identity map induces a stratified map, $X \to X^*$, in the sense of Definition 1.5. We begin by proving that the regular part of X^* is connected.

Let $R = X \backslash X_{n-1}$ and $R^* = X^* \backslash X^*_{n-1}$ be the regular parts of X and X^*, respectively. If R^* is the union of two disjoint open sets, $R^* = U_1 \sqcup U_2$, then we have $R = (R \cap U_1) \sqcup (R \cap U_2)$. As R is connected, by hypothesis, and open, we must have $U_1 \cap R = \emptyset$ or $U_2 \cap R = \emptyset$. The open set R being dense in X, we obtain $U_1 = \emptyset$ or $U_2 = \emptyset$, which implies the connectivity of R^*.

We denote by $\underline{L}$ and $\underline{L}^*$ the filtered face sets associated to X and X^*, respectively, and consider the following diagram whose elements are detailed below,

$$\begin{tikzcd}
\widetilde{A}_{PL,\overline{q}}(\underline{L}^*) \arrow[r, dashed, "\varphi_1"] \arrow[d, "f_{\underline{L}^*}"'] & \widetilde{A}_{PL,\overline{q}}(\underline{L}) \arrow[d, "f_{\underline{L}}"] \\
\widetilde{C}^*_{\overline{q}}(\underline{L}^*) \arrow[d, "g_{\underline{L}^*}"'] & \widetilde{C}^*_{\overline{q}}(\underline{L}) \arrow[d, "g_{\underline{L}}"] \\
C^*_{\mathrm{GM},\overline{p}}(\underline{L}^*) = \hom(C^{\overline{p}}_*(X^*), \mathbb{Q}) \arrow[r, "\varphi_2"] \arrow[d, "h_{\underline{L}^*}"'] & C^*_{\mathrm{GM},\overline{p}}(\underline{L}) = \hom(C^{\overline{p}}_*(X), \mathbb{Q}) \arrow[d, "h_{\underline{L}}"] \\
\hom(K^{\overline{p}}_*(X^*), \mathbb{Q}) \arrow[r, "\varphi_3"] & \hom(K^{\overline{p}}_*(X), \mathbb{Q})
\end{tikzcd}$$

The perversities $\overline{p}$ and $\overline{q}$ are elements of $\mathcal{P}^n$ such that $\overline{p}(k) + \overline{q}(k) = k - 2$, if $k \geq 2$. All vertical maps are quasi-isomorphisms, as this is proved in

- Corollary 7.3 for $f_{\underline{L}}$ and $f_{\underline{L}^*}$,

- Theorem D for $g_{\underline{L}^*}$ and $g_{\underline{L}}$,
- Proposition 3.4 for $h_{\underline{L}^*}$ and $h_{\underline{L}}$.

The proof that φ_3 is a quasi-isomorphism being done in [23, Theorem 9], the proof of Corollary 12.4 is reduced to the construction of φ_1 making the diagram commutative.

From Theorem A (see also Remark 2.4), we know that the stratified map, $X \to X^*$, induces an amalgamation of the singular simplices, generated by elementary amalgamations of the type $\Delta^p * \Delta^q \mapsto \emptyset * \Delta^{p+q+1}$. In Lemma 12.13, we show that these amalgamations induce a morphism of perverse CDGA's, $\varphi_1 \colon \widetilde{A}_{PL,\bullet}(\underline{L}^*) \to \widetilde{A}_{PL,\bullet}(\underline{L})$ making commutative the previous diagram and this ends the proof. ☐

Lemma 12.13. *Denote by* Φ_i *the identity map on a simplex* Δ*, corresponding to the elementary amalgamation*

$$\Delta^{j_0} * \cdots * \Delta^{j_n} \mapsto \Delta^{k_0} * \cdots * \Delta^{k_n},$$

with

$$\begin{aligned} k_a &= j_a, & \text{if} \quad & a < i \text{ or } a > i+1, \\ k_i &= -1, & \text{and} \quad & k_{i+1} = j_i + j_{i+1}. \end{aligned}$$

Let $\mu \colon \widetilde{\Delta}_1 = c\Delta^{j_0} \times \cdots \times \Delta^{j_n} \to \Delta$ *and* $\mu' \colon \widetilde{\Delta}_2 = c\Delta^{k_0} \times \cdots \times \Delta^{k_n} \to \Delta$ *be the projections defining the two blow-up's. Then, for all* $i \in \{0, \ldots, n-1\}$*, the map* Φ_i *lifts in a map* $\widetilde{\Phi}_i$ *between the two blow-up's, compatible with the face operators of* Δ *and such that* $\mu' \circ \widetilde{\Phi}_i = \mu$*. The induced cochain map,* $\widetilde{\Phi}_i^*$*, is such that, the following diagram commutes,*

$$\begin{tikzcd}
A_{PL}(c\Delta^{j_0}) \otimes \cdots \otimes A_{PL}(\Delta^{j_n}) \arrow[r, "\int"] & \mathbb{Q} \\
A_{PL}(c\Delta^{k_0}) \otimes \cdots \otimes A_{PL}(\Delta^{k_n}) \arrow[u, "\widetilde{\Phi}_i^*"] \arrow[ru, "\int"'] &
\end{tikzcd}$$

where $\int \omega = \int_{\mathring{\Delta}} (\mu^{-1})^* \omega$ *if* $\omega \in A_{PL}(c\Delta^{j_0}) \otimes \cdots \otimes A_{PL}(\Delta^{j_n})$ *and a similar formula for the second integral map. Moreover, the map* $\widetilde{\Phi}_i^*$ *verifies*

$$\widetilde{\Phi}_i^*(\widetilde{A}_{PL,\overline{q}}(\Delta^{k_0} * \cdots * \Delta^{k_n})) \subset \widetilde{A}_{PL,\overline{q}}(\Delta^{j_0} * \cdots * \Delta^{j_n}),$$

for all positive loose perversity $\overline{q}$*.*

Proof. Recall, from Remark 5.8, that the two blow-up's can be described in barycentric coordinates by

$$\mu((x_0,t_0), \ldots, (x_{n-1},t_{n-1}), x_n) = (x_0, t_0x_1, t_0t_1x_2, \ldots, t_0 \cdots t_{n-1}x_n)$$

with $x_i \in \mathbb{R}^{j_i+1}$, $t_i \in \mathbb{R}$ for all i, and a similar formula for μ'. For the study of Φ_i, we have two cases, depending if $i+1 = n$ or not.

1) We begin with $i+1 \neq n$. We construct a map $\widetilde{\Phi}_i \colon c\Delta^{j_0} \times \cdots \times \Delta^{j_n} \to c\Delta^{k_0} \times \cdots \times \Delta^{k_n}$ by $\widetilde{\Phi}_i = \mathrm{id} \times f_i \times \mathrm{id}$, where

$$f_i \colon c\Delta^{j_i} \times c\Delta^{j_{i+1}} \to c\emptyset \times c(\Delta^{j_i} * \Delta^{j_{i+1}})$$

is defined in barycentric coordinates by

$$f_i((x_i,t_i),(x_{i+1},t_{i+1})) = ((\vartheta),(x_i, t_ix_{i+1}, t_it_{i+1})),$$

with $\{\vartheta\}=c\emptyset$. We check easily from the definition that the map $\widetilde{\Phi}_i$ verifies $\mu'\circ\widetilde{\Phi}_i=\mu$ and is compatible with face operators. Thus it induces a CDGA's map $\widetilde{\Phi}_i^*\colon A_{PL}(c\Delta^{k_0})\otimes\cdots\otimes A_{PL}(\Delta^{k_n})\to A_{PL}(c\Delta^{j_0})\otimes\cdots\otimes A_{PL}(\Delta^{j_n})$.

Let $\omega=\omega_0\otimes\cdots\otimes\omega_n\in A_{PL}(c\Delta^{k_0})\otimes\cdots\otimes A_{PL}(\Delta^{k_n})$. In the next equalities, we use the fact that μ and μ' are diffeomorphisms on the interior of the integration domains,

$$\begin{aligned}\int_{\widetilde{\Delta}_2}\omega &= \int_{\mathring{\Delta}}({\mu'}^{-1})^*\omega=\int_{\mathring{\Delta}}\left((\mu^{-1})^*\circ\widetilde{\Phi}_i^*\right)(\omega)\\ &= \int_{\widetilde{\Delta}_1}\widetilde{\Phi}_i^*(\omega).\end{aligned}$$

This proves the equality $\int\circ\widetilde{\Phi}_i^*=\int$. We show now the compatibility with perversities, i.e.,

$$\|\omega\|_{n-\ell}\leq\overline{q}(n-\ell)\Rightarrow\|\widetilde{\Phi}_i^*\omega\|_{n-\ell}\leq\overline{q}(n-\ell),$$

for all $\ell\in\{0,\dots,n-1\}$. Consider first $\ell=0$, with $0<i$. The commutativity of the next diagram,

$$\begin{tikzcd}
\big(\Delta^{j_0}\times\{1\}\big)\times\cdots\times c\Delta^{j_i}\times c\Delta^{j_{i+1}}\times\cdots\times\Delta^{j_n} \arrow[r, "\mathrm{pr}"] \arrow[d, "\widetilde{\Phi}_i"] & \Delta^{j_0}\times\{1\} \\
\big(\Delta^{j_0}\times\{1\}\big)\times\cdots\times c\emptyset\times c(\Delta^{j_i}*\Delta^{j_{i+1}})\times\cdots\times\Delta^{j_n}, \arrow[ur, "\mathrm{pr}"'] &
\end{tikzcd}$$

implies

$$\|\widetilde{\Phi}_i^*(\omega)\|_n\leq\|\omega\|_n.$$

The same argument works for all the perverse degrees, except in $n-i$ and $n-i-1$.

We continue with $\ell=i+1$. In this case, the restriction of the map $\widetilde{\Phi}_i=\mathrm{id}\times f_i\times\mathrm{id}$,

$$\begin{tikzcd}
c\Delta^{j_0}\cdots\times c\Delta^{j_i}\times\big(\Delta^{j_{i+1}}\times\{1\}\big)\times\cdots\times\Delta^{j_n} \arrow[d, "\widetilde{\Phi}_i=\mathrm{id}\times f_i\times\mathrm{id}"'] \\
c\Delta^{j_0}\cdots\times c\emptyset\times\big((\Delta^{j_i}*\Delta^{j_{i+1}})\times\{1\}\big)\times\cdots\times\Delta^{j_n},
\end{tikzcd}$$

is defined by $f_i((x,t),(y,0))=(\vartheta,(x,ty,0))$. (Observe that the face $\Delta^{j_i}\times\{1\}$ of $c\Delta^{j_i}$, used in Example 5.9, corresponds to $t=0$ in terms of barycentric coordinates.) We denote by pr_1 the projection of the domain of $\widetilde{\Phi}_i$ on $c\Delta^{j_0}\cdots\times c\Delta^{j_i}\times\big(\Delta^{j_{i+1}}\times\{1\}\big)$ and by pr_2 the projection of the codomain on $c\Delta^{j_0}\cdots\times c\emptyset\times\big((\Delta^{j_i}*\Delta^{j_{i+1}})\times\{1\}\big)$. From the equality $\mathrm{pr}_2\circ\widetilde{\Phi}_i=\widetilde{\Phi}_i\circ\mathrm{pr}_1$ and the definition of f_i, we deduce that the map $\widetilde{\Phi}_i$ induces the identity between the fibers of these projections. This implies

$$\|\widetilde{\Phi}_i^*(\omega)\|_{n-i-1}\leq\|\omega\|_{n-i-1}.$$

The last perversity we have to study corresponds to $\ell=i$. The restriction of $\widetilde{\Phi}_i$ to $\widetilde{S}_i=c\Delta^{j_0}\times\cdots\times\big(\Delta^{j_i}\times\{1\}\big)\times c\Delta^{j_{i+1}}\times\cdots\times\Delta^{j_n}$ is defined by

$f_i((x,0),(y,s)) = (\vartheta,(x,0,0))$. Thus we have

$$\begin{aligned} \widetilde{\Phi}_i(\widetilde{S}_i) &= c\Delta^{j_0}\times\cdots\times c\emptyset\times\left(\Delta^{j_i}\times\{1\}\right)\times c\Delta^{j_{i+2}}\times\cdots\times\Delta^{j_n} \\ &\subset c\Delta^{j_0}\times\cdots\times c\emptyset\times(\Delta^{j_i}*\Delta^{j_{i+1}}\times\{1\})\times c\Delta^{j_{i+2}}\times\cdots\times\Delta^{j_n}. \end{aligned}$$

We denote by pr_1 the projection of $\widetilde{S}_i$ on $c\Delta^{j_0}\times\cdots\times c\emptyset\times\left(\Delta^{j_i}\times\{1\}\right)$ and by pr_2 the projection of $c\Delta^{j_0}\times\cdots\times c\emptyset\times(\Delta^{j_i}*\Delta^{j_{i+1}}\times\{1\})\times c\Delta^{j_{i+2}}\times\cdots\times\Delta^{j_n}$ on $c\Delta^{j_0}\times\cdots\times c\emptyset\times(\Delta^{j_i}*\Delta^{j_{i+1}}\times\{1\})$. The projection pr_1 is used for the determination of $\|\widetilde{\Phi}_i^*\omega\|_{n-i}$ and the projection pr_2 for the determination of $\|\omega\|_{n-i-1}$. The equality $\widetilde{\Phi}_i\circ\mathrm{pr}_1=\mathrm{pr}_2\circ\widetilde{\Phi}_i$ implies

$$\|\widetilde{\Phi}_i^*(\omega)\|_{n-i}\leq\|\omega\|_{n-i-1}\leq\overline{q}(n-i-1)\leq\overline{q}(n-i).$$

2) We study now the case $i+1=n$. The map $\widetilde{\Phi}_{n-1}=\mathrm{id}\times f_{n-1}$, with $f_{n-1}\colon c\Delta^{j_{n-1}}\times\Delta^{j_n}\to c\emptyset\times\left(\Delta^{j_{n-1}}*\Delta^{j_n}\right)$, is defined by $f_{n-1}((x,t),y)=(\vartheta,(x,ty))$. The proof goes like in the previous case, except for $\ell=n-1$. The restriction of $\widetilde{\Phi}_{n-1}$ to $\widetilde{S}_{n-1}=c\Delta^{j_0}\times\cdots\times\left(\Delta^{j_{n-1}}\times\{1\}\right)\times\Delta^{j_n}$ is defined by $f_{n-1}((x,0),y)=(\vartheta,(x,0))$ and we have $\widetilde{\Phi}_{n-1}(\widetilde{S}_{n-1})=c\Delta^{j_0}\times\cdots\times c\emptyset\times\Delta^{j_{n-1}}$. In the next diagram, the map h is defined by $h(x,0)=(\vartheta,(x,0))$,

$$\begin{tikzcd}
c\Delta^{j_0}\times\cdots\times\left(\Delta^{j_{n-1}}\times\{1\}\right)\times\Delta^{j_n} \arrow[r, "\mathrm{pr}"] \arrow[d, "\widetilde{\Phi}_{n-1}=\mathrm{id}\times f_{n-1}"'] & c\Delta^{j_0}\times\cdots\times\left(\Delta^{j_{n-1}}\times\{1\}\right) \arrow[dl, "\mathrm{id}\times h"] \\
c\Delta^{j_0}\times\cdots\times c\emptyset\times\left(\Delta^{j_{n-1}}*\Delta^{j_n}\right) &
\end{tikzcd}$$

The commutativity of this diagram implies

$$\|\widetilde{\Phi}_{n-1}^*(\omega)\|_1=\|\mathrm{pr}^*\circ(\mathrm{id}\times h)^*\omega\|_1\leq 0.$$

□

The construction of the minimal model of a cone follows directly from Example 8.9.

Corollary 12.14. *Let $\underline{S}$ be a connected face set of Sullivan minimal model $(\wedge V,d)$. Then the perverse minimal model of the cone, $c\underline{S}$, is $(\wedge V,d)$, where the perverse degree coincides with the homological degree.*

Therefore, the cone, $c\underline{S}$, is an intersection-formal filtered face set if, and only if, $\underline{S}$ is a formal face set.

The treatment we have used for the study of intersection cohomology allows spaces of infinite dimension. For instance, a perverse model of $c\mathbb{C}P(\infty)$ is $(\wedge x,0)$, with $|x|=\|x\|=2$, $n=1$.

For getting a filtered face set which is not intersection-formal, it is sufficient to take the cone on a non-formal face set. For instance, the cone on the pullback of the Hopf fibration, $S^3\to S^7\to S^4$, along a map $S^2\times S^2\to S^4$ of degree one, is not intersection-formal, (see Example 2.91 of [9] for more details on this space).

Remark 12.15. In the description of the complex C of Example 8.10, we may replace $A_{PL}(\underline{S})$ by a model, $(\wedge V,d)$, of $\underline{S}$. Thus a perverse model of $\Sigma\underline{S}$ is a perverse model of $\wedge(t,dt)\otimes(\wedge V,d)$, with the perverse degrees defined as in Example 8.10.

Example 12.16. Consider a lattice of perversities containing

$$\begin{tikzcd}
& \overline{p}_1 & \\
\overline{p}_2 \arrow[ur] & \overline{p}_3 \arrow[u] & \overline{p}_4 \arrow[ul] \\
\overline{p}_5 \arrow[u] \arrow[ur, dashed] & \overline{p}_6 \arrow[ul] \arrow[ur] & \overline{p}_7 \arrow[u] \arrow[ul, dashed] \\
& \overline{p}_8 \arrow[ul] \arrow[u] \arrow[ur] &
\end{tikzcd}$$

Suppose that $A_\bullet$ is a perverse cdga such that $A_{\overline{p}_8} = \mathbb{Q}\, e$ with $|e| = 2$ and $H^+(A_{\overline{p}_i}) = 0$ if $i < 8$. Then the minimal model will contain elements,

- of degree 1, x of perversity $\overline{p}_5$, y of perversity $\overline{p}_6$ and z of perversity $\overline{p}_7$ such that $dx = dy = dz = e$,
- of degree 0, f of perversity $\overline{p}_2$, g of perversity $\overline{p}_3$, h of perversity $\overline{p}_4$ such that $df = x - y$, $dg = x - z$, $dh = y - z$,
- of degree -1, α of perversity $\overline{p}_1$ such that $d\alpha = f - g + h$.

This example shows that we cannot have a model with generators of positive degree for any perverse cdga. Observe that this perverse cdga cannot be the cohomology of a filtered face set because the map $H^1(A_{\overline{p}_6} + A_{\overline{p}_7}) \to H^1(A_{\overline{p}_4})$ is not injective. This is in contradiction with the injectivity of such a map established in the proof of Proposition 12.11, see formula (12).

Example 12.17. Proposition 12.6 contains the fact that $H^1(A_{\overline{t}}) = 0$ implies $H^1(A_{\overline{p}}) = 0$, for any perversity $\overline{p}$. This example shows that the reverse way is not true.

We choose $A_{\overline{p}} = \mathcal{A}(\underline{K})_{\overline{p}}$, where $\underline{K} = c(S^1 \times S^1)$ is the cone on the torus, stratified by its summit. As a cone is a contractible space, we have $H^0(A_{\overline{0}}) = H^0(\underline{K}; \mathbb{Q}) = \mathbb{Q}$ and $H^1(A_{\overline{0}}) = H^1(\underline{K}; \mathbb{Q}) = 0$. The filtered face set $\underline{K}$ has two strata and a perversity consists of an integer. For the perversity $\overline{1}$, constant on 1, we have

$$H^i_{\overline{1}}(\underline{K}; \mathbb{Q}) = \begin{cases} H^i(S^1 \times S^1; \mathbb{Q}) & \text{if} \quad i \leq 1, \\ 0 & \text{if} \quad i > 2. \end{cases}$$

Thus we get $H^1(A_{\overline{0}}) = 0$ and $H^1(A_{\overline{1}}) = \mathbb{Q}$.

References

1. Markus Banagl, *Topological invariants of stratified spaces*, Springer Monographs in Mathematics, Springer, Berlin, 2007. MR 2286904 (2007j:55007)
2. ———, *Intersection spaces, spatial homology truncation, and string theory*, Lecture Notes in Mathematics, vol. 1997, Springer-Verlag, Berlin, 2010. MR 2662593
3. A. Borel and et al., *Intersection cohomology*, Modern Birkhäuser Classics, Birkhäuser Boston Inc., Boston, MA, 2008, Notes on the seminar held at the University of Bern, Bern, 1983, Reprint of the 1984 edition. MR 2401086 (2009k:14046)
4. Jean-Paul Brasselet, Gilbert Hector, and Martin Saralegi, *Théorème de de Rham pour les variétés stratifiées*, Ann. Global Anal. Geom. **9** (1991), no. 3, 211–243. MR 1143404 (93g:55009)

5. Jean-Luc Brylinski, *Equivariant intersection cohomology*, Kazhdan-Lusztig theory and related topics (Chicago, IL, 1989), Contemp. Math., vol. 139, Amer. Math. Soc., Providence, RI, 1992, pp. 5–32. MR 1197827 (94c:55010)
6. Bohumil Cenkl, Gilbert Hector, and Martin Saralegi, *Cohomologie d'intersection modérée. Un théorème de de Rham*, Pacific J. Math. **169** (1995), no. 2, 235–289. MR 1346255 (96k:55007)
7. J. Daniel Christensen and Mark Hovey, *Quillen model structures for relative homological algebra*, Math. Proc. Cambridge Philos. Soc. **133** (2002), no. 2, 261–293. MR 1912401 (2003f:18012)
8. Yves Félix, Stephen Halperin, and Jean-Claude Thomas, *Rational homotopy theory*, Graduate Texts in Mathematics, vol. 205, Springer-Verlag, New York, 2001. MR 1802847 (2002d:55014)
9. Yves Félix, John Oprea, and Daniel Tanré, *Algebraic models in geometry*, Oxford Graduate Texts in Mathematics, vol. 17, Oxford University Press, Oxford, 2008. MR 2403898 (2009a:55006)
10. Yves Félix and Daniel Tanré, *Topologie algébrique*, Sciences SUP. Dunod, Paris, 2010.
11. Greg Friedman, *Stratified fibrations and the intersection homology of the regular neighborhoods of bottom strata*, Topology Appl. **134** (2003), no. 2, 69–109. MR 2009092 (2004g:55008)
12. ______, *On the chain-level intersection pairing for PL pseudomanifolds*, Homology, Homotopy Appl. **11** (2009), no. 1, 261–314. MR 2529162 (2010m:55004)
13. Rudolf Fritsch and Marek Golasiński, *Topological, simplicial and categorical joins*, Arch. Math. (Basel) **82** (2004), no. 5, 468–480. MR 2061452 (2005a:55012)
14. Antonio Gómez-Tato, *Modèles minimaux résolubles*, J. Pure Appl. Algebra **85** (1993), no. 1, 43–56. MR 1207067 (94c:55017)
15. Antonio Gómez-Tato, Stephen Halperin, and Daniel Tanré, *Rational homotopy theory for non-simply connected spaces*, Trans. Amer. Math. Soc. **352** (2000), no. 4, 1493–1525. MR 1653355 (2000i:55035)
16. Mark Goresky and Robert MacPherson, *Intersection homology theory*, Topology **19** (1980), no. 2, 135–162. MR 572580 (82b:57010)
17. ______, *Intersection homology. II*, Invent. Math. **72** (1983), no. 1, 77–129. MR 696691 (84i:57012)
18. ______, *Stratified Morse theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 14, Springer-Verlag, Berlin, 1988. MR 932724 (90d:57039)
19. R. Mark Goresky, *Intersection homology operations*, Comment. Math. Helv. **59** (1984), no. 3, 485–505. MR 761809 (86i:55008)
20. Werner Greub, Stephen Halperin, and Ray Vanstone, *Connections, curvature, and cohomology. Vol. I: De Rham cohomology of manifolds and vector bundles*, Academic Press, New York, 1972, Pure and Applied Mathematics, Vol. 47. MR 0336650 (49 #1423)
21. S. Halperin, *Lectures on minimal models*, Mém. Soc. Math. France (N.S.) (1983), no. 9-10, 261. MR 736299 (85i:55009)
22. Mark Hovey, *Intersection homological algebra*, New topological contexts for Galois theory and algebraic geometry (BIRS 2008), Geom. Topol. Monogr., vol. 16, Geom. Topol. Publ., Coventry, 2009, pp. 133–150. MR 2544388 (2010g:55009)
23. Henry C. King, *Topological invariance of intersection homology without sheaves*, Topology Appl. **20** (1985), no. 2, 149–160. MR 800845 (86m:55010)
24. Frances Kirwan and Jonathan Woolf, *An introduction to intersection homology theory*, second ed., Chapman & Hall/CRC, Boca Raton, FL, 2006. MR 2207421 (2006k:32061)
25. Steven L. Kleiman, *The development of intersection homology theory*, Pure Appl. Math. Q. **3** (2007), no. 1, Special Issue: In honor of Robert D. MacPherson. Part 3, 225–282. MR 2330160 (2008e:55006)
26. Saunders Mac Lane, *Homology*, Classics in Mathematics, Springer-Verlag, Berlin, 1995, Reprint of the 1975 edition. MR 1344215 (96d:18001)
27. Daniel Quillen, *Rational homotopy theory*, Ann. of Math. (2) **90** (1969), 205–295. MR 0258031 (41 #2678)

28. Frank Quinn, *Intrinsic skeleta and intersection homology of weakly stratified sets*, Geometry and topology (Athens, Ga., 1985), Lecture Notes in Pure and Appl. Math., vol. 105, Dekker, New York, 1987, pp. 233–249. MR 873296 (88g:57022)
29. ———, *Homotopically stratified sets*, J. Amer. Math. Soc. **1** (1988), no. 2, 441–499. MR 928266 (89g:57050)
30. C. P. Rourke and B. J. Sanderson, *$\triangle$-sets. I. Homotopy theory*, Quart. J. Math. Oxford Ser. (2) **22** (1971), 321–338. MR 0300281 (45 #9327)
31. Martin Saralegi, *Un théorème de dualité entre l'homologie d'intersection et la cohomologie $\mathcal{L}^2$*, Ph.D. thesis, Lille, 1988.
32. Martintxo Saralegi-Aranguren, *de Rham intersection cohomology for general perversities*, Illinois J. Math. **49** (2005), no. 3, 737–758 (electronic). MR 2210257 (2006k:55013)
33. Edwin H. Spanier, *Algebraic topology*, McGraw-Hill Book Co., New York, 1966. MR 0210112 (35 #1007)
34. Paul Stienne, *Modèles minimaux et cohomologie d'intersection*, Lille, 2002.
35. Dennis Sullivan, *Infinitesimal computations in topology*, Inst. Hautes Études Sci. Publ. Math. (1977), no. 47, 269–331 (1978). MR 0646078 (58 #31119)
36. Daniel Tanré, *Homotopie rationnelle: modèles de Chen, Quillen, Sullivan*, Lecture Notes in Mathematics, vol. 1025, Springer-Verlag, Berlin, 1983. MR 764769 (86b:55010)
37. Andrei Verona, *Stratified mappings—structure and triangulability*, Lecture Notes in Mathematics, vol. 1102, Springer-Verlag, Berlin, 1984. MR 771120 (86k:58010)

Département de Mathematiques, UMR 8524 et Fédération CNRS Nord-Pas-de-Calais FR 2956, Université de Lille 1, 59655 Villeneuve d'Ascq Cedex, France
E-mail address: `David.Chataur@math.univ-lille1.fr`

Laboratoire de Mathématiques de Lens, EA 2462 et Fédération CNRS Nord-Pas-de-Calais FR 2956, Université d'Artois, SP18, rue Jean Souvraz, 62307 Lens Cedex, France
E-mail address: `saralegi@euler.univ-artois.fr`

Département de Mathematiques, UMR 8524 et Fédération CNRS Nord-Pas-de-Calais FR 2956, Université de Lille 1, 59655 Villeneuve d'Ascq Cedex, France
E-mail address: `Daniel.Tanre@univ-lille1.fr`
URL: `http://www.tanre.org/Pro`